✓ Complete

O



Matrix-Matrix Multiplication

We multiply two matrices by breaking it into several vector multiplications and concatenating the result.

$$\begin{bmatrix} a & b \\ c & d \\ e & f \end{bmatrix} * \begin{bmatrix} w & x \\ y & z \end{bmatrix} = \begin{bmatrix} a*w+b*y & a*x+b*z \\ c*w+d*y & c*x+d*z \\ e*w+f*y & e*x+f*z \end{bmatrix}$$

An **m x n matrix** multiplied by an **n x o matrix** results in an **m x o** matrix. In the above example, a 3 x 2 matrix times a 2 x 2 matrix resulted in a 3 x 2 matrix.

To multiply two matrices, the number of **columns** of the first matrix must equal the number of **rows** of the second matrix.

For example:

```
% Initialize a 3 by 2 matrix
    A = [1, 2; 3, 4; 5, 6]
    % Initialize a 2 by 1 matrix
 5
    B = [1; 2]
     % We expect a resulting matrix of (3 \text{ by } 2)*(2 \text{ by } 1) = (3 \text{ by } 1)
                                                                                            Run
    mult_AB = A*B
                                                                                            Reset
    % Make sure you understand why we got that result
A =
   1
       2
B =
   1
mult_AB =
    5
   11
```

1 of 1 5/14/18, 4:44 AM