# QCD VBFMET samples MadGraph Studies

João Pela

Imperial College London

2015-08-03





2015-08-03

#### Introduction

## Summary of PPD meeting

- Presentation was made on our proposal for QCD VBF-MET samples and feedback was requested.
- Main concern points were:
  - $\bullet$  The huge amount of time required by obtaining events in the low pT hats (filter efficiency is  $5\times 10^{-6}$  for working point A at lowest pT hat)
  - $\bullet$  The amount of events going over DIGI (apparently this request would be  $\sim$  15% of the a last campaign)
- The size of the sample itself did not sound like a big problem.
- It was suggested to look into looking into MadGraph with some reasonable generator cuts in order to reduce filtering needs and processing time.

### Steps to preform

After the meeting in a chat within our group it was suggested

- ullet Looking into usable MadGraph variables, avoid  $\Delta\phi$  or even  $\Delta\eta$  cuts.
- Study the relationship between generator partons and generator jets
- Study filter efficiency
- (I would also like to) Look into parton versus HLT objects
- (I would also like to) Look into a mini sample of this versus data.

#### Introduction

The event production runs on MadGraph are controlled by a set of "card" files which contain all the parameters that are relevant. For the *run\_card.dat* here are the relevant parameters for our analysis.

## Relevant run\_card.dat parameters

- nevents: Number of unweighted events requested
- ptj: (default 20) minimum pt for the jets
- etaj: (default 5.0) max rap for the jets
- drjj: (default 0.4) min distance between jets
- mmjj: (default 0.0) min invariant mass of a jet pair
- ptj1min: (default 0.0) minimum pt for the leading jet in pt
- ptj2min: (default 0.0) minimum pt for the second jet in pt
- deltaeta: (default 0.0) minimum rapidity for two jets in the WBF case



# MadGraph process cross sections

I have tried several working points to check what would be the necessary number of events to produce.

## Madgraph working points

etaj	ptj1min	ptj2min	mmjj	Cross Section [pb]	Notes
5.0	0	0	0	$7.008 \times 10^8 \pm 6.648 \times 10^5$	MadGraph Default values
4.8	0	0	0	$6.879 \times 10^8 \pm 6.644 \times 10^5$	Reduncing jet eta range
4.8	40	40	0	$5.266 \times 10^7 \pm 4.772 \times 10^4$	Require 2 jets with $p_{\perp} >$ 40 GeV
4.8	40	40	800	$8.911 \times 10^5 \pm 653.2$	Working Point A
4.8	35	35	800	$1.21  imes 10^6 \pm 878.9$	WP A: dijet p⊥ -5 GeV
4.8	30	30	800	$1.699  imes 10^6 \pm 1304$	WP A: dijet $p_{\perp}$ -10 GeV
4.8	40	40	700	$1.234  imes 10^6 \pm 940$	WP A: dijet p⊥ -100 GeV
4.8	40	40	900	$6.611  imes 10^5 \pm 482.6$	WP A: dijet $p_{\perp}$ +100 GeV
4.8	40	40	1000	$5.009  imes 10^5 \pm 377$	WP A: dijet $p_{\perp}$ +200 GeV
4.8	50	50	1000	$2.948 \times 10^5 \pm 222.8$	Working Point B

- Cross section obtained from generating 100k events with MG5\_aMC\_v2.3.0 (from 2015-07-01)
- Process p p > j j where j = g u c d s u $\sim$  c $\sim$  d $\sim$  s $\sim$
- ullet Working Point A and be are basically the same as the previously proposed filters working points but without  $\Delta\phi$  or  $\Delta\eta$

The with pythia the total cross section of the pT hats to generate was:  $1.87 \times 10^8$  two orders of magnitude higher.



# CMSSW and Pythia8 hadronization

I have decided to proceed with Working Point A and with the help of Chayanit was able to interface CMSSW with the produced MadGraph events.

#### CMSSW details

- CMSSW\_7\_1\_18 version
- Hadronization done by Hadronizer\_TuneCUETP8M1\_13TeV\_MLM\_5f\_max4j\_LHE\_pythia8\_cff
- Generator jets are AK4 GenJetsNoNu

## Matching results

Events in	Events Pass	Event Eff [%]	xsec before [pb]	xsec match [pb]
100000	17077	$17.1 \pm 0.1$	$8.911 \times 10^5 \pm 6.532 \times 10^2$	$\boxed{1.522 \times 10^5 \pm 1.066 \times 10^3}$

#### Questions

- Which file to use: events.lhe.gz or unweighted\_events.lhe.gz (results with the unweighted events).
- Is this matching efficiency ok?
- What cross section to use? Pre or post matching?

# Matching Partons with Generator jets

### Parton selection

- For partons selecting status:
  - 23 outgoing
  - 24 outgoing, nonperturbatively kicked out in diffraction

### Pairing Partons and Generator Jets

- ullet Selecting all generator jets within  $\Delta R < 0.4$  (this may need to be a big bigger)
- ullet From those selecting the generator jet with the lowest  $p_{\perp}$  to the parton as a match.
  - ullet This avoids picking up the wrong jet from just picking lowest  $\Delta R$

### Pairing Results for 10k events

- Pythia8 matched events events: 1712
- Found genJet for 2 parton: 1489 (87.0%)
- Found genJet for 1 parton: 212 (12.4%)
- Found genJet for 2 parton: 11 (0.6%)
- Matched genJet was not the lowest  $\Delta R$ : 120

