

SHARP PLANAR QUANTITATIVE SAINT-VENANT STABILITY BY REDUCTION TO A GRAPH INSET

ABSTRACT. We prove the sharp planar quantitative Saint–Venant inequality $\mathcal{A}(\Omega)^2 \leq C \delta_T(\Omega)$, where $\mathcal{A}$ is the Fraenkel asymmetry and δ_T is the torsional (Saint–Venant) deficit, with an absolute constant C and no regularity assumption on Ω . The proof reduces a near-extremal Ω to a small-amplitude radial graph by a purely level-set construction governed by the two level deficits — the Cauchy–Schwarz/Serrin defect D_H and the isoperimetric defect D_I — and concludes with the perturbative stability of nearly spherical sets. In particular the argument uses neither the Brasco–De Philippis–Velichkov selection principle, nor mass transport, nor symmetry-type (Serrin / Magnanini–Poggesi) estimates. Two sharp ingredients drive the reduction: a fold-content bound free of the quantitative-isoperimetric square-root loss, obtained from a point-source flux identity and a thin-annulus cancellation; and the observation that excising the low-torsion boundary caps to manufacture a graph *decreases* the scale-invariant torsional deficit. The manuscript is self-contained: every nonstandard step is proved in full, and the few external inputs are classical theorems quoted with precise references.

CONTENTS

1. Introduction	2
The Saint–Venant inequality and its quantitative form	2
Prior work on quantitative stability	3
The main theorem	3
Strategy and what is new	4
Organization of the paper	5
Notation	5
2. Preliminaries	5
2.1. Sets of finite perimeter	5
2.2. The coarea formula and differentiation of the distribution function	6
2.3. Isoperimetric inequalities	7
2.4. The sharp quantitative isoperimetric inequality	8
2.5. Interior regularity of the torsion function	8
2.6. Sard’s theorem and regular values of the torsion function	9
3. The level deficits and the volume profile	10
3.1. The flux and coarea identities	11
3.2. The two level deficits	12
3.3. The level-set deficit identity	12
3.4. Scale invariance of the identity	14
3.5. Talenti’s comparison and the volume profile	15
3.6. A pointwise lower bound on $-\mu'$	16

4. The inset: regularity, deficit transfer, level selection	17
4.1. No interior holes	17
4.2. Deficit transfer to super-level sets	18
4.3. Selection of a good level	19
4.4. Closeness of the super-level set	19
4.5. One connected inset: the spillover is higher order	20
4.6. Controls on the inset	21
5. The trapping annulus	23
5.1. The inner and outer radii	24
5.2. The angular coarea identity	25
5.3. Two crossings at every intermediate radius	26
5.4. The radial-profile bound	27
5.5. The trapping theorem	27
6. Fold content and the radial graph	30
6.1. Point-source flux identities	31
6.2. Sharp fold content	33
6.3. The graph defect	35
6.4. Realization as a continuous radial graph	35
7. Cutting the caps lowers the deficit	37
7.1. The torsion as a Green double integral	37
7.2. Boundary-layer torsion stability	38
7.3. Cutting boundary material decreases δ_*	39
8. Perturbative stability of nearly spherical sets	42
8.1. Statement and standing normalisation	42
8.2. The dual (complementary) variational principle	43
8.3. The comparison field	43
8.4. The exact second-order energy	44
8.5. Coercivity of the deficit	46
8.6. The asymmetry upper bound and conclusion	48
9. Proof of the main theorem	48
9.1. The reduction	49
9.2. The main theorem	52
References	53

1. INTRODUCTION

The Saint–Venant inequality and its quantitative form. Let $\Omega \subset \mathbb{R}^2$ be a bounded open set of finite perimeter with $|\Omega| = \pi$. The *torsion function* of Ω is the unique solution $u \in H_0^1(\Omega)$ of $-\Delta u = 1$ in Ω , $u = 0$ on $\partial\Omega$; its integral $T(\Omega) := \int_{\Omega} u$ is the *torsional rigidity*. The classical Saint–Venant conjecture, proved by Pólya and Szegő [9] via the Schwarz symmetrization of the

torsion potential, asserts that the disc maximises torsional rigidity at fixed area:

$$T(\Omega) \leq T(B_1) = \frac{\pi}{8},$$

with equality if and only if Ω is a disc. A more precise, and substantially more difficult, question is to quantify stability: how large must Ω deviate from a disc when the deficit $\delta_T(\Omega) := \frac{\pi}{8} - T(\Omega)$ is small? The natural measure of deviation is the *Fraenkel asymmetry*

$$\mathcal{A}(\Omega) := \min_{x_0 \in \mathbb{R}^2} \frac{|\Omega \Delta B(x_0, r_\Omega)|}{|\Omega|}, \quad r_\Omega := \sqrt{|\Omega|/\pi},$$

which is zero if and only if Ω is a disc. The sharp quantitative Saint–Venant inequality is the statement

$$\mathcal{A}(\Omega)^2 \leq C \delta_T(\Omega) \tag{1}$$

for some *absolute* constant C (independent of Ω , of $|\Omega|$, and of any regularity of $\partial\Omega$). The exponent 2 on the left and the exponent 1 on the right are both optimal: the ratio $\mathcal{A}(\Omega)^2/\delta_T(\Omega)$ is bounded and bounded away from zero for a one-parameter family of ellipses.

Talenti [11] proved a pointwise comparison between the decreasing rearrangement of u_Ω and the torsion function of the equimeasure disc, giving $T(\Omega) \leq T(B_1)$ and the quantitative comparison $\mu(v) \leq \mu_B(v)$ between the distribution functions. This Talenti comparison, which we state precisely as Theorem 3.8, is a key input in the present work. Talenti’s result also implies the profile identity $\int_0^{1/4} (\mu_B - \mu) dv = \delta_T$ (see Lemma 3.9), which localizes the deficit to a single integral of a non-negative integrand.

Prior work on quantitative stability. The Faber–Krahn problem for the first Dirichlet eigenvalue λ_1 is the structural analogue of the Saint–Venant problem, and the two share much of their quantitative theory. For the quantitative isoperimetric inequality the sharp form $\mathcal{A}(E) \leq C_F \delta_{\text{iso}}(E)^{1/2}$ (where $\delta_{\text{iso}}(E)$ denotes the isoperimetric deficit and C_F is an absolute constant; see Theorem 2.11) was established by Fusco, Maggi, and Pratelli [6]; this result, which we use as a black box, is the *only* external quantitative inequality the present paper imports.

For Saint–Venant/Faber–Krahn stability, Brasco and Pratelli [4] proved a quantitative stability result for both the torsional rigidity and the first eigenvalue, establishing (1) with a *non-sharp* exponent. The sharp inequality (1) — with exponent 2 and an absolute constant, valid with no regularity on $\partial\Omega$ — was established by Brasco, De Philippis, and Velichkov [3]. Their argument proceeds by a *selection principle*: from a near-extremal sequence one extracts a subsequence that, after appropriate normalisation, converges in a strong topology to a $C^{1,\gamma}$ domain; the stability of the latter is then proved by a second-variation analysis. The selection principle is a powerful and elegant tool, but it is non-constructive, non-quantitative in the convergence, and it relies on the regularity theory for minimizers of shape functionals in a fundamental way. A survey of quantitative spectral inequalities, including the state of the art up to 2017, is given by Brasco and De Philippis [2].

The main theorem. The main result of this paper is the following, proved in Section 9 as Theorem 9.2.

Theorem 1.1 (Main theorem; see Theorem 9.2). *There exists an absolute constant C such that every bounded open set $\Omega \subset \mathbb{R}^2$ with $|\Omega| = \pi$ satisfies*

$$\mathcal{A}(\Omega)^2 \leq C \delta_T(\Omega).$$

This is the same conclusion as [3], but obtained by an entirely different route. The proof requires no selection principle, no mass transport, and no symmetry-type argument (Serrin overdetermined problem / Magnanini–Poggesi estimates). The only external inputs are: classical function-space and geometric-measure-theory tools (the coarea formula, the planar isoperimetric inequality, the Talenti comparison, and the sharp quantitative isoperimetric inequality of [6]), all quoted with precise references in Section 2.

Strategy and what is new. The argument replaces the selection–regularity–second-variation architecture of [3] by a purely level-set construction driven by the two *level deficits*:

$$\begin{aligned} D_H(v) &:= \mu(v)(-\mu'(v)) - P(\{u > v\})^2 \geq 0, \\ D_I(v) &:= P(\{u > v\})^2 - 4\pi\mu(v) \geq 0, \end{aligned}$$

where $\mu(v) = |\{u > v\}|$ is the distribution function and $P(\{u > v\})$ is the perimeter of the superlevel set. The central identity (Theorem 3.4, proved in Section 3)

$$\int_0^m (D_H(v) + D_I(v)) dv = 4\pi \delta_T(\Omega)$$

encodes the entire deficit in a non-negative integral over the level parameter. All the estimates in the paper are consequences of this identity and the classical Talenti comparison.

The two sharp ingredients. Two novel observations make the reduction exact:

- (1) *Fold-content bound with no square-root loss.* At the optimal inset level $\hat{v} \sim \sqrt{\delta_T}$, the connected component $\tilde{E}$ of $\{u > \hat{v}\}$ of largest measure is trapped in a thin annulus $B_{\rho_i}(z) \subset \tilde{E} \subset B_{\rho_e}(z)$ of width $\eta = \rho_e - \rho_i \lesssim \delta_T^{1/8}$ (established in Section 5). The total angular measure of rays along which $\tilde{E}$ is *not* a radial graph is bounded, via the point-source flux identity and the cancellation $|I_{\tilde{E}} - 2\pi\hat{r}| \leq a\eta/(\rho_i\hat{r})$ (where $a = |\tilde{E} \Delta B_{\hat{r}}(z)|$), by a quantity of order $\text{Exc} + a\eta$ — without the factor of $\sqrt{\delta_{\text{iso}}}$ that the naive application of the quantitative isoperimetric inequality [6] would introduce. This is the content of Proposition 6.3 and Corollary 6.5 in Section 6.
- (2) *Cutting caps decreases the deficit.* To manufacture a genuine radial graph G from $\tilde{E}$ one must excise the “caps” (the portions of $\tilde{E}$ that cause each ray to intersect $\partial\tilde{E}$ more than once). A potential concern is that this excision increases the scale-invariant deficit $\delta_*(G) := (T(B_G) - T(G))/T(B_G)$. In fact it does not: the equal-area comparison ball shrinks faster than torsion is lost, so δ_* *decreases*. This is Lemma 7.5 in Section 7, proved via the Green-function representation of torsional rigidity and a barrier estimate in the collar.

With $\tau := \sqrt{\delta_T}$, the two observations combine to produce a continuous radial graph G with amplitude $\lesssim \delta_T^{1/8}$, scale-invariant deficit $\delta_*(G) \leq C\delta_*(\Omega)$, and symmetric difference $|\Omega \Delta G| \leq C\sqrt{\delta_T}$ (the precise reduction is Theorem 9.1 in Section 9). The perturbative stability of nearly spherical sets (Theorem 8.1 in Section 8) then gives $\mathcal{A}(G)^2 \leq C_G\delta_*(G)$, and the triangle inequality for the symmetric difference transfers this to $\mathcal{A}(\Omega)^2 \leq C\delta_T(\Omega)$.

Organization of the paper. Section 2 collects the classical prerequisites: the theory of sets of finite perimeter (De Giorgi–Federer structure theorem, coarea formula, differentiation of the distribution function), the planar isoperimetric inequality and the sharp quantitative version [6], the interior regularity and analyticity of the torsion function [7], and Sard’s theorem.

Section 3 introduces the two level deficits D_H and D_I , proves their non-negativity, establishes the central level-set deficit identity $\int_0^m (D_H + D_I) dv = 4\pi\delta_T$ (Theorem 3.4), and derives the Talenti comparison and the profile identity.

Section 4 carries out the level-set construction: it proves deficit transfer to any superlevel set, selects the good level $\hat{v}$ by an averaging argument, establishes the closeness of $\{u > \hat{v}\}$ to Ω , and isolates the inset $\tilde{E}$ as the dominant connected component.

Section 5 proves that a connected hole-free finite-perimeter set with controlled isoperimetric deficit is trapped in a thin concentric annulus, using the angular-perimeter coarea identity and a crossing-number argument.

Section 6 bounds the fold content of $\tilde{E}$ — the obstruction to being a radial graph — via the point-source flux identities and the thin-annulus cancellation, achieving the sharp estimate without the quantitative-isoperimetric square-root loss.

Section 7 proves that excising the caps to manufacture a radial graph decreases the scale-invariant torsion deficit, via a Green-function comparison and a barrier estimate.

Section 8 establishes the perturbative stability of nearly spherical sets for the torsion functional, proving that any radial graph of small amplitude satisfies $\mathcal{A}^2 \leq C_G\delta_*$, with explicit absolute constants, by a second-order expansion of the torsion deficit in Fourier modes.

Section 9 assembles all pieces, optimises the level parameter at $\tau = \sqrt{\delta_T}$, and derives the main theorem in full.

Notation. Throughout the paper, $\Omega \subset \mathbb{R}^2$ is a bounded open set with $|\Omega| = \pi$. The unit disc is B_1 ; $B(x_0, r)$ denotes the open disc of radius r centred at x_0 . The symbol C denotes a generic positive absolute constant whose value may change from line to line; named constants such as C_G , η_G , A_0 , δ_* are fixed at their first occurrence. All *absolute* constants are independent of Ω and of any geometric or regularity data of $\partial\Omega$. The notation $f \lesssim g$ means $f \leq Cg$ for an absolute C .

2. PRELIMINARIES

Throughout this paper $\Omega \subset \mathbb{R}^2$ denotes a bounded open set, $|\cdot|$ denotes Lebesgue measure on $\mathbb{R}^2$, $\mathcal{H}^1$ denotes one-dimensional Hausdorff measure, and $B_r(x) \subset \mathbb{R}^2$ is the open disc of radius r centred at x ; we write $B_r = B_r(0)$. The fixed normalisation $|\Omega| = \pi$ is in force from Section 1 onward but is not required for the statements in this section unless explicitly noted. We collect here the definitions and classical results the rest of the paper relies on. Standard results are stated precisely with full hypotheses and cited to published sources.

2.1. Sets of finite perimeter. We follow the conventions of [8].

Definition 2.1 (Perimeter). Let $E \subset \mathbb{R}^2$ be Lebesgue measurable and let $A \subset \mathbb{R}^2$ be open. The *perimeter of E in A* is

$$P(E; A) := \sup \left\{ \int_E \operatorname{div} \varphi \, dx : \varphi \in C_c^1(A; \mathbb{R}^2), \|\varphi\|_{L^\infty} \leq 1 \right\}.$$

We set $P(E) := P(E; \mathbb{R}^2)$. The set E is of *finite perimeter* (in A) if $P(E; A) < \infty$.

If E has locally finite perimeter, the distributional gradient $\mu_E := -D\mathbf{1}_E$ is an $\mathbb{R}^2$ -valued Radon measure with total variation $|\mu_E|(A) = P(E; A)$ [8, Proposition 12.1, Definition 12.1].

Definition 2.2 (Reduced boundary). For E of locally finite perimeter, the *reduced boundary* $\partial^* E$ is the set of $x \in \text{supp } |\mu_E|$ at which

$$\nu_E(x) := \lim_{r \rightarrow 0^+} \frac{\mu_E(B_r(x))}{|\mu_E|(B_r(x))}$$

exists and satisfies $|\nu_E(x)| = 1$. The vector $\nu_E(x)$ is the *measure-theoretic outer unit normal* to E .

Theorem 2.3 (De Giorgi structure theorem). *Let $E \subset \mathbb{R}^2$ have locally finite perimeter. Then $\partial^* E$ is countably $\mathcal{H}^1$ -rectifiable, $\mu_E = \nu_E \mathcal{H}^1 \llcorner \partial^* E$, and for every Borel set A ,*

$$P(E; A) = \mathcal{H}^1(\partial^* E \cap A).$$

Proof. See [8, Theorem 15.9, Corollary 16.1] or [5, Section 5.7, Theorem 5.15]. $\square$

Definition 2.4 (Essential boundary). For a measurable set $E \subset \mathbb{R}^2$, its *essential boundary* is $\partial^e E := \mathbb{R}^2 \setminus (E^{(0)} \cup E^{(1)})$, where $E^{(t)} := \{x : \lim_{r \rightarrow 0} |E \cap B_r(x)| / |B_r(x)| = t\}$.

Theorem 2.5 (Federer). *If $E \subset \mathbb{R}^2$ has locally finite perimeter then $\partial^* E \subseteq E^{(1/2)} \subseteq \partial^e E$ and $\mathcal{H}^1(\partial^e E \setminus \partial^* E) = 0$. In particular, $P(E; A) = \mathcal{H}^1(\partial^e E \cap A)$ for every Borel A .*

Proof. See [8, Theorem 15.5, Theorem 16.2] or [5, Theorem 5.16]. $\square$

2.2. The coarea formula and differentiation of the distribution function.

Theorem 2.6 (Coarea formula). *Let $\Omega \subset \mathbb{R}^2$ be open and let $u \in W^{1,1}(\Omega)$. Then for every nonnegative Borel function $g : \Omega \rightarrow [0, \infty]$,*

$$\int_{\Omega} g(x) |\nabla u(x)| dx = \int_{-\infty}^{\infty} \left(\int_{\{u=t\}} g d\mathcal{H}^1 \right) dt. \quad (2)$$

Moreover, for Lebesgue-almost every $t \in \mathbb{R}$ the superlevel set $\{u > t\}$ has finite perimeter in Ω , the level set $\{u = t\}$ is $\mathcal{H}^1$ -rectifiable with $\mathcal{H}^1(\{u = t\}) < \infty$, and

$$P(\{u > t\}; \Omega) = \mathcal{H}^1(\{u = t\}) \quad \text{for a.e. } t.$$

Proof. The coarea formula (2) for $W^{1,1}$ functions is [8, Theorem 13.1]; see also [5, Section 3.4, Theorem 3.13]. The perimeter identity and rectifiability of almost every level set follow from the BV-coarea formula [1, Theorem 3.40]: since $\int_{\mathbb{R}} P(\{u > t\}; \Omega) dt = \int_{\Omega} |\nabla u| dx < \infty$, the set $\{u > t\}$ has finite perimeter for a.e. t . $\square$

The following corollary, which we call the *differentiation lemma* for the distribution function, is used repeatedly in the level-set analysis.

Lemma 2.7 (Differentiation of the distribution function). *Let $\Omega \subset \mathbb{R}^2$ be bounded and let $u \in W^{1,1}(\Omega) \cap C^1(\Omega)$ with $u \geq 0$ (no continuity up to $\partial\Omega$ is assumed; the torsion function of (10), which is in $H_0^1(\Omega) \subset W^{1,1}(\Omega)$ for bounded Ω and in $C^\infty(\Omega)$ by Theorem 2.13, satisfies these hypotheses).*

Define $\mu(v) := |\{u > v\}|$ for $v \geq 0$. Then μ is non-increasing and absolutely continuous, and for Lebesgue-almost every $v > 0$,

$$-\mu'(v) = \int_{\{u=v\}} \frac{1}{|\nabla u|} d\mathcal{H}^1. \quad (3)$$

In particular, the Cauchy–Schwarz inequality applied to the pair $(1, |\nabla u|^{1/2})$ on $\{u = v\}$ gives

$$P(\{u > v\})^2 = \mathcal{H}^1(\{u = v\})^2 \leq \left(\int_{\{u=v\}} |\nabla u| d\mathcal{H}^1 \right) (-\mu'(v)) \quad \text{for a.e. } v. \quad (4)$$

Proof. The coarea formula (2) with $g = |\nabla u|^{-1} \mathbf{1}_{\{\nabla u \neq 0\}}$ gives, for $0 \leq s < t$,

$$|\{s < u \leq t\} \cap \{\nabla u \neq 0\}| = \int_s^t \left(\int_{\{u=v\} \cap \{\nabla u \neq 0\}} \frac{1}{|\nabla u|} d\mathcal{H}^1 \right) dv.$$

Since $u \in C^1(\Omega)$, Sard's theorem (Theorem 2.16 below) implies that $\{u = v\} \cap \{\nabla u = 0\}$ is empty for a.e. v , so the restriction to $\{\nabla u \neq 0\}$ is superfluous for a.e. v . Also $|\{\nabla u = 0\} \cap \{s < u \leq t\}|$ contributes zero to the left side (it has measure zero by a standard argument). Hence $\mu(s) - \mu(t) = \int_s^t (-\mu') dv$ with $-\mu'(v) = \int_{\{u=v\}} |\nabla u|^{-1} d\mathcal{H}^1$ a.e.

For (4): by Cauchy–Schwarz on $\{u = v\}$,

$$\mathcal{H}^1(\{u = v\})^2 = \left(\int_{\{u=v\}} 1 d\mathcal{H}^1 \right)^2 \leq \left(\int_{\{u=v\}} |\nabla u| d\mathcal{H}^1 \right) \left(\int_{\{u=v\}} \frac{1}{|\nabla u|} d\mathcal{H}^1 \right) = \left(\int_{\{u=v\}} |\nabla u| d\mathcal{H}^1 \right) (-\mu'(v)),$$

and by Theorem 2.6, $\mathcal{H}^1(\{u = v\}) = P(\{u > v\})$ for a.e. v . $\square$

Remark 2.8. The integral $\int_{\{u=v\}} |\nabla u| d\mathcal{H}^1$ is the *flux* of the torsion function across the level set. When $-\Delta u = 1$ in $\{u > v\}$ with $u = v$ on $\partial\{u > v\}$ (for a.e. regular v), integration of $-\Delta u = 1$ over $\{u > v\}$ and Green's first identity give $\int_{\{u=v\}} |\nabla u| d\mathcal{H}^1 = |\{u > v\}| = \mu(v)$. Hence (4) becomes $P(\{u > v\})^2 \leq \mu(v)(-\mu'(v))$ for a.e. v ; the deficit from equality is $D_H(v) := \mu(v)(-\mu'(v)) - P(\{u > v\})^2 \geq 0$. This is developed in Section 3.

2.3. Isoperimetric inequalities.

Theorem 2.9 (Planar isoperimetric inequality). *For every set of finite perimeter $E \subset \mathbb{R}^2$ with $|E| < \infty$,*

$$P(E)^2 \geq 4\pi |E|, \quad (5)$$

with equality if and only if E is equivalent (up to a Lebesgue-null set) to a disc $B_r(x)$.

Proof. This is [8, Theorem 14.1]; see also [5, Section 5.6, Theorem 5.6]. $\square$

Theorem 2.10 (Relative isoperimetric inequality in a disc). *There is an absolute constant $C_{\text{rel}} > 0$ such that for every set of finite perimeter $E \subset \mathbb{R}^2$, every $x \in \mathbb{R}^2$, and every $r > 0$,*

$$\min\{|E \cap B_r(x)|, |B_r(x) \setminus E|\} \leq C_{\text{rel}} P(E; B_r(x))^2. \quad (6)$$

The constant C_{rel} is independent of E , x , and r .

Proof. By the relative isoperimetric inequality for convex domains, applied to the unit disc $B_1 \subset \mathbb{R}^2$, there is a constant $C_{\text{rel}} > 0$ such that for every set of finite perimeter E , $\min\{|E \cap B_1|, |B_1 \setminus E|\} \leq C_{\text{rel}} P(E; B_1)^2$.

$E|\} \leq C_{\text{rel}} P(E; B_1)^2$; see [8, Theorem 12.37] (stated in $\mathbb{R}^n$, here with $n = 2$, $K = B_1$) or [5, Theorem 5.9]. The general case at radius r follows by the substitution $\tilde{E} := \{y/r : y \in E\}$: since $|\tilde{E} \cap B_1| = r^{-2}|E \cap B_r(x)|$ and $P(\tilde{E}; B_1) = r^{-1}P(E; B_r(x))$, the unit-disc inequality for $\tilde{E}$ multiplied by r^2 yields (6) with the same constant C_{rel} . $\square$

2.4. The sharp quantitative isoperimetric inequality. For a set of finite perimeter $E \subset \mathbb{R}^2$ with $0 < |E| < \infty$ we define the *Fraenkel asymmetry*

$$\mathcal{A}(E) := \min_{x_0 \in \mathbb{R}^2} \frac{|E \triangle B_{r_E}(x_0)|}{|E|}, \quad r_E := \sqrt{|E|/\pi}, \quad (7)$$

and the *isoperimetric deficit*

$$\delta_{\text{iso}}(E) := \frac{P(E)}{2\sqrt{\pi|E|}} - 1 \geq 0, \quad (8)$$

where $2\sqrt{\pi|E|}$ is the perimeter of a disc of area $|E|$. Both quantities are dimensionless, translation- and scaling-invariant, and nonnegative, vanishing precisely when E is a disc.

Theorem 2.11 (Fusco–Maggi–Pratelli). *There is an absolute constant $C_F > 0$ such that for every set of finite perimeter $E \subset \mathbb{R}^2$ with $0 < |E| < \infty$,*

$$\mathcal{A}(E) \leq C_F \delta_{\text{iso}}(E)^{1/2}. \quad (9)$$

The exponent $\frac{1}{2}$ is optimal: it cannot be replaced by any strictly larger power, as shown by elongated ellipses with small eccentricity. The value of C_F is not explicit in [6]; it is finite and absolute (depending on neither E nor any geometric parameter of Ω).

Proof. This is Theorem 1.1 of [6] (stated there in $\mathbb{R}^n$; the planar case $n = 2$ is the specialisation $C_F := C(2)$). The theorem is proved in [6] using symmetrisation and a selection-principle argument; we use it as a black box. $\square$

Remark 2.12 (Relation between δ_{iso} and the additive deficit). The additive isoperimetric deficit $D_I(E) := P(E)^2 - 4\pi|E| \geq 0$ (see Definition 3.2) satisfies

$$\delta_{\text{iso}}(E) = \sqrt{1 + \frac{D_I(E)}{4\pi|E|}} - 1 = \frac{D_I(E)/(4\pi|E|)}{\sqrt{1 + D_I(E)/(4\pi|E|)} + 1} \leq \frac{D_I(E)}{8\pi|E|},$$

since $\sqrt{1+s} + 1 \geq 2$ for $s \geq 0$. Hence $\delta_{\text{iso}}(E)^{1/2} \leq (D_I(E)/(8\pi|E|))^{1/2}$, and (9) gives

$$\mathcal{A}(E)^2 \leq C_F^2 \delta_{\text{iso}}(E) \leq \frac{C_F^2}{8\pi} \frac{D_I(E)}{|E|}.$$

We will apply this estimate in Section 9 to the inset $\tilde{E}$.

2.5. Interior regularity of the torsion function. Let $u \in H_0^1(\Omega)$ be the unique weak solution of the torsion problem

$$-\Delta u = 1 \text{ in } \Omega, \quad u = 0 \text{ on } \partial\Omega, \quad (10)$$

i.e., $\int_{\Omega} \nabla u \cdot \nabla \varphi \, dx = \int_{\Omega} \varphi \, dx$ for all $\varphi \in H_0^1(\Omega)$. Existence and uniqueness follow from the Lax–Milgram theorem; $u \geq 0$ by the weak maximum principle [7, Theorem 8.1].

Theorem 2.13 (Interior smoothness and analyticity of u). *Let $\Omega \subset \mathbb{R}^2$ be a bounded open set and let $u \in H_0^1(\Omega)$ be the weak solution of (10). Then $u \in C^\infty(\Omega)$. Moreover, u is real-analytic in Ω .*

Proof. Since the right-hand side of (10) is $f \equiv 1 \in C^\infty(\mathbb{R}^2)$ and the operator $-\Delta$ has smooth (constant) coefficients, interior Schauder estimates bootstrap to give $u \in C_{\text{loc}}^{k,\alpha}(\Omega)$ for every $k \geq 0$ and every $\alpha \in (0, 1)$; see [7, Theorem 6.17, Corollary 6.18]. Hence $u \in C^\infty(\Omega)$.

Real-analyticity follows from the general fact that solutions of linear elliptic equations with real-analytic coefficients and real-analytic right-hand side are real-analytic in the interior: the Laplacian has constant (hence analytic) coefficients and $f \equiv 1$ is analytic, so u is real-analytic in Ω ; this is a classical result of Morrey–Nirenberg, discussed in [7, Chapter 6, §6.8 and the historical notes]. $\square$

Theorem 2.14 (Interior gradient and Hessian estimates). *There is an absolute constant $C > 0$ with the following property. Let $\Omega \subset \mathbb{R}^2$ be a bounded open set, let u satisfy (10) weakly, let $x \in \Omega$, and set $d := \text{dist}(x, \partial\Omega)$. Then*

$$|\nabla u(x)| \leq C \left(\frac{\|u\|_{L^\infty(B_d(x))}}{d} + d \right), \quad |D^2 u(x)| \leq C \left(\frac{\|u\|_{L^\infty(B_d(x))}}{d^2} + 1 \right). \quad (11)$$

Proof. Apply the interior Schauder estimate of [7, Theorem 6.2] to u on $B_d(x)$ where $-\Delta u = 1$. The scale-invariant form of the estimate gives

$$d \|\nabla u\|_{L^\infty(B_{d/2}(x))} + d^2 \|D^2 u\|_{L^\infty(B_{d/2}(x))} \leq C (\|u\|_{L^\infty(B_d(x))} + d^2),$$

with C depending only on the dimension $n = 2$. Dividing by d and d^2 respectively yields (11) (the sup over $B_{d/2}(x)$ bounds the pointwise value at x). $\square$

Lemma 2.15 (L^∞ bound). *Let $\Omega \subset \mathbb{R}^2$ be bounded and let u solve (10). Then $0 \leq u(x) \leq \frac{1}{4}(\text{diam } \Omega)^2$ for all $x \in \Omega$. In particular $m := \sup_\Omega u \leq \frac{1}{4}(\text{diam } \Omega)^2 < \infty$ (the supremum is taken over the open set Ω ; no continuity of u up to $\partial\Omega$ is assumed or asserted).*

Proof. The lower bound $u \geq 0$ follows from the weak maximum principle [7, Theorem 8.1]. For the upper bound, let $R = \text{diam } \Omega$ and pick any $x_0 \in \Omega$; then $\Omega \subset B_R(x_0)$. The function $w(x) := (R^2 - |x - x_0|^2)/4$ satisfies $-\Delta w = 1$ in $\mathbb{R}^2$ and $w \geq 0$ on Ω . Hence $h := w - u \in H^1(\Omega)$ satisfies $-\Delta h = 0$ weakly in Ω and $h = w - u \geq -u$, so $h^- := \max\{-h, 0\} = (u - w)^+ \leq u^+ = u$ has zero boundary trace, $(u - w)^+ \in H_0^1(\Omega)$ (as $u \in H_0^1$ and $w \geq 0$). By the weak maximum principle [7, Theorem 8.1] applied to the harmonic h , $h \geq 0$ a.e. in Ω , i.e. $u \leq w \leq R^2/4$ in Ω . No continuity of u up to $\partial\Omega$ is used. $\square$

2.6. Sard's theorem and regular values of the torsion function.

Theorem 2.16 (Sard). *Let $U \subseteq \mathbb{R}^d$ be open and $f \in C^k(U; \mathbb{R}^m)$ with $k \geq \max\{1, d - m + 1\}$. Then the set of critical values of f ,*

$$f(\{x \in U : \text{rank } Df(x) < m\}),$$

has Lebesgue measure zero in $\mathbb{R}^m$. In particular, if $f \in C^\infty(U)$ and $m = 1$, the set $f(\{\nabla f = 0\}) \subset \mathbb{R}$ is a Lebesgue-null set.

Proof. See [8, Theorem A.6] for the statement in the context used here; the original source is [10]. $\square$

Lemma 2.17 (Almost every level of u is regular). *Let $\Omega \subset \mathbb{R}^2$ be a bounded open set and let u solve (10). Then for Lebesgue-almost every $v \in (0, m)$ the level v is a regular value of u in Ω : that is, $\nabla u(x) \neq 0$ at every point $x \in \{u = v\} \cap \Omega$. For every such v ,*

- (i) the level set $\{u = v\} \cap \Omega$ is a smooth, properly embedded 1-dimensional submanifold of Ω (a disjoint union of smooth curves); if moreover $u \in C(\overline{\Omega})$ with $u = 0$ on $\partial\Omega$, it is a finite union of smooth closed curves, compactly contained in Ω ;
- (ii) the superlevel set $\{u > v\}$ is a finite-perimeter set with $P(\{u > v\}) = \mathcal{H}^1(\{u = v\})$;
- (iii) the differentiation formula (3) holds at v , with $1/|\nabla u|$ finite and smooth on $\{u = v\}$.

Proof. By Theorem 2.13, $u \in C^\infty(\Omega)$ (indeed real-analytic). Applying Theorem 2.16 with $d = 2$, $m = 1$: the set of critical values $u(\{\nabla u = 0\} \cap \Omega) \subset \mathbb{R}$ has Lebesgue measure zero. Hence for a.e. v , every point of $\{u = v\} \cap \Omega$ is a regular point. For such v :

(i): At a regular value v the implicit function theorem gives $\{u = v\} \cap \Omega$ as a smooth, properly embedded 1-dimensional submanifold of the open set Ω (without boundary in Ω), i.e. a disjoint union of smooth curves, each either a closed loop in Ω or a properly embedded line escaping every compact subset of Ω . This is the honest *interior* statement and uses no boundary regularity. If, in addition, $u \in C(\overline{\Omega})$ with $u = 0$ on $\partial\Omega$ (which holds whenever $\partial\Omega$ is, say, continuous), then for $v > 0$ the level set $\{u = v\}$ is compactly contained in Ω and so is a *finite* disjoint union of smooth closed curves. The two genuinely interior consequences (ii) and (iii) below, proved from the coarea formula alone, hold with no boundary regularity whatsoever.

(ii): By Theorem 2.6, $P(\{u > v\}) = \mathcal{H}^1(\{u = v\})$ for a.e. v .

(iii): For a.e. v , (3) holds and $|\nabla u| > 0$ on $\{u = v\}$, so $1/|\nabla u|$ is finite and smooth there by Theorem 2.13. $\square$

3. THE LEVEL DEFICITS AND THE VOLUME PROFILE

This section is the analytic heart of the paper. We package the Saint-Venant deficit

$$\delta_T = \delta_T(\Omega) = \frac{\pi}{8} - T(\Omega) \geq 0$$

of a normalised domain $|\Omega| = \pi$ into the level sets of its torsion function u . Recall (Section 2) that $u \in H_0^1(\Omega) \cap C^\infty(\Omega)$ solves $-\Delta u = 1$ in Ω , $u = 0$ on $\partial\Omega$, that

$$T(\Omega) := \int_{\Omega} u \, dx = \int_{\Omega} |\nabla u|^2 \, dx, \quad m := \max_{\Omega} u,$$

the second equality holding because testing $-\Delta u = 1$ against $u \in H_0^1(\Omega)$ gives $\int_{\Omega} |\nabla u|^2 = \int_{\Omega} u$. We define two non-negative defects of each level set; their integral over all levels turns out to be exactly $4\pi\delta_T$. We then use Talenti's comparison to pin the volume profile $\mu(v) := |\{u > v\}|$, obtaining $m \leq \frac{1}{4}$, an exact profile identity, and the pointwise bound $-\mu' \geq 4\pi$. These facts feed directly into the inset construction of Section 4.

Throughout, a value $v \in (0, m)$ is called *regular* if $\nabla u \neq 0$ on $\{u = v\}$; by Lemma 2.17 almost every $v \in (0, m)$ is regular, and for such v the level set $\{u = v\}$ is a (closed, embedded) C^∞ curve in Ω and the coarea quotient $1/|\nabla u|$ is finite and smooth on it. We write

$$E_v := \{x \in \Omega : u(x) > v\}, \quad \mu(v) := |E_v|, \quad P(v) := P(E_v) = \mathcal{H}^1(\{u = v\})$$

for the super-level set, its area, and its perimeter (the last equality holding for regular v , the level set being a smooth curve).

3.1. The flux and coarea identities. The two structural identities below are the only place where the equation $-\Delta u = 1$ enters this section; everything afterwards is geometry of the level sets. The first (flux) identity quantifies the source; the second (coarea) is the differentiation formula for μ . We prove both.

Lemma 3.1 (Flux and coarea identities). *For a.e. $v \in (0, m)$,*

$$\int_{\{u=v\}} |\nabla u| d\mathcal{H}^1 = \mu(v), \quad (12)$$

and for a.e. $v \in (0, m)$ the function μ is differentiable with

$$-\mu'(v) = \int_{\{u=v\}} \frac{1}{|\nabla u|} d\mathcal{H}^1. \quad (13)$$

Moreover μ is absolutely continuous and nonincreasing on $(0, m)$, with $\mu(0^+) = |\Omega| = \pi$ and $\mu(m^-) = 0$.

Proof. Flux (12). We derive this directly from the weak formulation, so that no boundary regularity of Ω is needed. Recall $u \in H_0^1(\Omega)$ solves

$$\int_{\Omega} \nabla u \cdot \nabla \varphi dx = \int_{\Omega} \varphi dx \quad \text{for all } \varphi \in H_0^1(\Omega). \quad (14)$$

Fix $0 < v < m$ and, for small $h > 0$ with $v + h < m$, take the Lipschitz cut-off

$$\psi_h(x) := \min\left\{1, \frac{1}{h}(u(x) - v)_+\right\} \in [0, 1].$$

Since $t \mapsto \min\{1, \frac{1}{h}(t - v)_+\}$ is Lipschitz and vanishes for $t \leq v$ (in particular at $t = 0$), and $u \in H_0^1(\Omega)$, the composition ψ_h lies in $H_0^1(\Omega)$ and is an admissible test function in (14), with

$$\nabla \psi_h = \frac{1}{h} \nabla u \mathbf{1}_{\{v < u < v+h\}} \quad \text{a.e.}$$

Inserting $\varphi = \psi_h$ into (14),

$$\frac{1}{h} \int_{\{v < u < v+h\}} |\nabla u|^2 dx = \int_{\Omega} \nabla u \cdot \nabla \psi_h dx = \int_{\Omega} \psi_h dx.$$

We pass to the limit $h \downarrow 0$ on each side. On the left, the coarea formula (Theorem 2.6) gives

$$\frac{1}{h} \int_{\{v < u < v+h\}} |\nabla u|^2 dx = \frac{1}{h} \int_v^{v+h} \left(\int_{\{u=t\}} |\nabla u| d\mathcal{H}^1 \right) dt \xrightarrow{h \downarrow 0} \int_{\{u=v\}} |\nabla u| d\mathcal{H}^1$$

for a.e. v , namely at every Lebesgue point of the $L^1(0, m)$ function $t \mapsto F(t) := \int_{\{u=t\}} |\nabla u| d\mathcal{H}^1$ (that $F \in L^1$ follows from $\int_0^m F dt = \int_{\{0 < u < m\}} |\nabla u|^2 dx \leq \int_{\Omega} |\nabla u|^2 < \infty$ by coarea). On the right, $0 \leq \psi_h \leq 1$ and $\psi_h \uparrow \mathbf{1}_{\{u > v\}}$ pointwise as $h \downarrow 0$, so by monotone (or dominated, since $|\Omega| < \infty$) convergence

$$\int_{\Omega} \psi_h dx \xrightarrow{h \downarrow 0} \int_{\Omega} \mathbf{1}_{\{u > v\}} dx = |\{u > v\}| = \mu(v).$$

Equating the two limits yields (12) for a.e. $v \in (0, m)$.

Coarea (13) and absolute continuity. This is the differentiation of the distribution function, Lemma 2.7: for a.e. v , μ is differentiable with $-\mu'(v) = \int_{\{u=v\}} |\nabla u|^{-1} d\mathcal{H}^1$. We record the absolute-continuity statement explicitly, as it is used repeatedly. Set $\Phi(v) := \int_{\{u=v\}} |\nabla u|^{-1} d\mathcal{H}^1 \geq 0$ for those

v where this is finite. The coarea formula (Theorem 2.6) with $g = \mathbf{1}_{\{u>0\}}|\nabla u|^{-1}\mathbf{1}_{\{\nabla u \neq 0\}}$ gives

$$\int_0^m \Phi(\tau) d\tau = \int_{\{0 < u < m\} \cap \{\nabla u \neq 0\}} 1 dx = |\Omega| = \pi < \infty,$$

where the middle equality uses that $\{\nabla u = 0\} \cap \{0 < u < m\}$ has Lebesgue measure zero (it is contained in the union of level sets at critical values, a set of measure zero by Sard's theorem, Theorem 2.16 and Lemma 2.17, intersected with Ω). Hence $\Phi \in L^1(0, m)$. For $0 < v_1 < v_2 < m$ the coarea formula applied with $g = \mathbf{1}_{\{v_1 < u \leq v_2\}}|\nabla u|^{-1}\mathbf{1}_{\{\nabla u \neq 0\}}$ yields

$$\mu(v_1) - \mu(v_2) = |\{v_1 < u \leq v_2\}| = \int_{v_1}^{v_2} \Phi(\tau) d\tau.$$

Thus $\mu(v) = \int_v^m \Phi(\tau) d\tau$ for $v \in (0, m)$; being the tail integral of an L^1 function, μ is absolutely continuous and nonincreasing on $(0, m)$, with $\mu' = -\Phi$ a.e., which is (13). Finally $\mu(0^+) = \int_0^m \Phi = |\Omega| = \pi$ (equivalently $|\{u > 0\}| = |\Omega|$ since $u > 0$ in Ω by the strong minimum principle), and $\mu(m^-) = \int_m^m \Phi = 0$. $\square$

3.2. The two level deficits.

Definition 3.2 (The two level deficits). For a regular value $v \in (0, m)$ we define the *Cauchy–Schwarz* (or Hölder/Serrin) defect and the *isoperimetric* defect of the level set $\{u = v\}$ by

$$D_H(v) := \mu(v) (-\mu'(v)) - P(v)^2, \quad D_I(v) := P(v)^2 - 4\pi \mu(v).$$

The names anticipate Lemma 3.3: D_H vanishes precisely when the flux is equidistributed (constant $|\nabla u|$ on the level curve), and D_I vanishes precisely when the level set is a disk.

Lemma 3.3 (Non-negativity of the deficits). *For a.e. $v \in (0, m)$ one has*

$$D_H(v) \geq 0, \quad D_I(v) \geq 0.$$

Moreover $D_I(v) = 0$ if and only if E_v is (equivalent to) a disk, and, for regular v , $D_H(v) = 0$ if and only if $|\nabla u|$ is constant $\mathcal{H}^1$ -almost everywhere on $\{u = v\}$.

Proof. $D_I \geq 0$. This is the planar isoperimetric inequality (Theorem 2.9) applied to the finite-perimeter set E_v : $P(E_v)^2 \geq 4\pi|E_v|$, i.e. $P(v)^2 \geq 4\pi\mu(v)$, with equality if and only if E_v is a disk up to a Lebesgue-null modification.

$D_H \geq 0$. Fix a regular value v at which (12) and (13) hold (a.e. v , by Lemmas 2.17 and 3.1); then $\Sigma = \{u = v\}$ is a smooth embedded curve on which $|\nabla u| > 0$ and $P(v) = \mathcal{H}^1(\Sigma) < \infty$. Apply the Cauchy–Schwarz inequality on the finite measure $\mathcal{H}^1 \llcorner \Sigma$ to the functions $|\nabla u|^{1/2}$ and $|\nabla u|^{-1/2}$:

$$P(v)^2 = \left(\int_{\Sigma} 1 d\mathcal{H}^1 \right)^2 = \left(\int_{\Sigma} |\nabla u|^{1/2} \cdot |\nabla u|^{-1/2} d\mathcal{H}^1 \right)^2 \leq \left(\int_{\Sigma} |\nabla u| d\mathcal{H}^1 \right) \left(\int_{\Sigma} \frac{1}{|\nabla u|} d\mathcal{H}^1 \right).$$

By (12) the first factor is $\mu(v)$, and by (13) the second factor is $-\mu'(v)$. Hence $P(v)^2 \leq \mu(v)(-\mu'(v))$, i.e. $D_H(v) \geq 0$. Equality in Cauchy–Schwarz holds if and only if $|\nabla u|^{1/2}$ and $|\nabla u|^{-1/2}$ are proportional $\mathcal{H}^1$ -a.e. on Σ , i.e. if and only if $|\nabla u|$ is $\mathcal{H}^1$ -a.e. constant on $\{u = v\}$. $\square$

3.3. The level-set deficit identity. We now prove the central identity and, as the mandate requires, derive the constant 4π independently of any prior computation, directly from coarea and integration by parts (the Saint–Venant route).

Theorem 3.4 (Level-set deficit identity). *With the notation above,*

$$\int_0^m (D_H(v) + D_I(v)) dv = 4\pi \delta_T(\Omega). \quad (15)$$

Proof. The proof has three steps: a pointwise telescoping, two integrations, and the assembly with the explicit value of the constant.

Step 1: telescoping. For a.e. $v \in (0, m)$ the squared-perimeter terms in Definition 3.2 cancel:

$$D_H(v) + D_I(v) = \left(\mu(v)(-\mu'(v)) - P(v)^2 \right) + \left(P(v)^2 - 4\pi\mu(v) \right) = \mu(v)(-\mu'(v)) - 4\pi\mu(v). \quad (16)$$

The perimeter has disappeared entirely; what remains involves only μ and μ' .

Step 2: the first term, by integration by parts. By Lemma 3.1, μ is absolutely continuous on $(0, m)$ with $\mu' \in L^1$, $\mu(0^+) = \pi$ and $\mu(m^-) = 0$. Consequently $v \mapsto \frac{1}{2}\mu(v)^2$ is absolutely continuous with derivative $\mu(v)\mu'(v) \in L^1(0, m)$ (a product of a bounded absolutely continuous function and an L^1 function, with the Leibniz rule valid because μ is bounded and absolutely continuous). Hence

$$\int_0^m \mu(v)(-\mu'(v)) dv = -\left[\frac{1}{2}\mu(v)^2 \right]_{v=0^+}^{v=m^-} = \frac{1}{2}\mu(0^+)^2 - \frac{1}{2}\mu(m^-)^2 = \frac{1}{2}|\Omega|^2 = \frac{\pi^2}{2}. \quad (17)$$

Step 3: the second term, by the layer-cake formula. Since $u > 0$ in Ω and $\max u = m$,

$$\int_0^m 4\pi \mu(v) dv = 4\pi \int_0^\infty |\{u > v\}| dv = 4\pi \int_\Omega u dx = 4\pi T(\Omega), \quad (18)$$

the middle equality being Cavalieri's (layer-cake) formula for the nonnegative function u , valid since $\mu(v) = 0$ for $v \geq m$.

Assembly and the constant. Integrating (16) over $(0, m)$ and inserting (17)–(18),

$$\int_0^m (D_H + D_I) dv = \frac{\pi^2}{2} - 4\pi T(\Omega).$$

It remains to identify $\frac{\pi^2}{2} - 4\pi T(\Omega)$ with $4\pi\delta_T$. We do *not* invoke any earlier formula for $T(B_1)$; instead we read off the constant from Step 2. Indeed

$$\frac{\pi^2}{2} = \frac{|\Omega|^2}{2} = 4\pi \cdot \frac{|\Omega|^2}{8\pi} = 4\pi \cdot \frac{\pi^2}{8\pi} = 4\pi \cdot \frac{\pi}{8},$$

so that

$$\int_0^m (D_H + D_I) dv = 4\pi \left(\frac{\pi}{8} - T(\Omega) \right) = 4\pi \delta_T(\Omega),$$

by the definition $\delta_T = \frac{\pi}{8} - T(\Omega)$. This proves (15).

We pause to make the provenance of the constant transparent, since it is the keystone of the paper. The factor 4π in D_I is the planar isoperimetric constant ($P^2 \geq 4\pi A$). The factor $\frac{1}{2}|\Omega|^2$ in Step 2 is purely the integration of $\mu(-\mu')$, owing nothing to the equation beyond the boundary values $\mu(0^+) = |\Omega|$, $\mu(m^-) = 0$. These two independent inputs combine into the deficit because of the elementary algebraic identity $\frac{1}{2}A^2 = 4\pi \cdot A^2/(8\pi)$ for any area A ; with $A = |\Omega| = \pi$ this reads $\frac{1}{2}|\Omega|^2 = 4\pi \cdot \frac{\pi}{8}$, which is the algebra displayed above. The number $\frac{\pi}{8}$ so produced is exactly the disk torsion $T(B_1)$ (computed independently in Lemma 3.9 below), so the right-hand side is $4\pi(T(B_1) - T(\Omega))$. Thus the right-hand side is $4\pi(T(B_1) - T(\Omega))$, the Saint-Venant deficit scaled by 4π , with no hidden normalisation. $\square$

Remark 3.5 (The identity as an exact, scale-aware statement). Identity (15) expresses the (global) Saint-Venant deficit as the total over all levels of two pointwise non-negative geometric defects of the level sets: D_H , the failure of $|\nabla u|$ to be constant on $\{u = v\}$; and D_I , the failure of E_v to be a disk. Both vanish identically (for all v) if and only if every level set is a disk on which $|\nabla u|$ is constant, i.e. if and only if Ω is a disk and $u = u_B$; this is Serrin's overdetermined characterisation, and it is the rigidity behind (15). The quantitative content is the converse, exploited from Section 4 on: small δ_T forces both defects to be small *on most levels*.

3.4. Scale invariance of the identity. For the inset of Section 4 we will apply the identity not to Ω but to its super-level sets, which are not area-normalised. We therefore record the unnormalised form of (15) and its exact homogeneity. Let $E \subset \mathbb{R}^2$ be any bounded open set of finite measure which is regular for the Dirichlet problem, with torsion function w_E ($-\Delta w_E = 1$ in E , $w_E \in H_0^1(E)$), torsion $T(E) = \int_E w_E = \int_E |\nabla w_E|^2$, maximum $m_E = \max w_E$, and level quantities $\mu_E(v) = |\{w_E > v\}|$, $P_E(v) = P(\{w_E > v\})$,

$$D_H^E(v) := \mu_E(v)(-\mu_E'(v)) - P_E(v)^2, \quad D_I^E(v) := P_E(v)^2 - 4\pi\mu_E(v).$$

Proposition 3.6 (Unnormalised identity and degree-4 homogeneity). *For any such E ,*

$$\int_0^{m_E} (D_H^E(v) + D_I^E(v)) dv = \frac{1}{2}|E|^2 - 4\pi T(E) = 4\pi(T(B_E) - T(E)), \quad T(B_E) := \frac{|E|^2}{8\pi}, \quad (19)$$

where B_E is any disk with $|B_E| = |E|$. Both sides are homogeneous of degree 4 under dilations: writing $sE = \{sx : x \in E\}$ for $s > 0$, one has

$$\int_0^{m_{sE}} (D_H^{sE} + D_I^{sE}) dv = s^4 \int_0^{m_E} (D_H^E + D_I^E) dv, \quad T(B_{sE}) - T(sE) = s^4(T(B_E) - T(E)). \quad (20)$$

Consequently the scale-invariant deficit $\delta_*(E) := (T(B_E) - T(E))/T(B_E) \in [0, 1)$ satisfies

$$\delta_*(E) = \frac{T(B_E) - T(E)}{T(B_E)} = \frac{\frac{1}{4\pi} \int_0^{m_E} (D_H^E + D_I^E) dv}{|E|^2/(8\pi)} = \frac{2}{|E|^2} \int_0^{m_E} (D_H^E + D_I^E) dv, \quad (21)$$

and $\delta_*(sE) = \delta_*(E)$ for all $s > 0$.

Proof. Identity (19). The proof of Lemma 3.1 and Theorem 3.4 used only $-\Delta w_E = 1$, $w_E \in H_0^1(E)$, interior smoothness, and Sard's theorem, all of which hold for E verbatim (interior regularity and analyticity of the torsion function depend only on the interior equation, not on boundary regularity; Theorem 2.13). Repeating the three steps gives

$$\int_0^{m_E} (D_H^E + D_I^E) dv = \frac{1}{2}\mu_E(0^+)^2 - 4\pi T(E) = \frac{1}{2}|E|^2 - 4\pi T(E),$$

since $\mu_E(0^+) = |\{w_E > 0\}| = |E|$ ($w_E > 0$ in E). The last equality in (19) is the algebraic identity $\frac{1}{2}|E|^2 = 4\pi \cdot |E|^2/(8\pi) = 4\pi T(B_E)$.

Homogeneity (20). The torsion function transforms under dilation by $w_{sE}(x) = s^2 w_E(x/s)$: indeed w_{sE} so defined vanishes on $\partial(sE)$ and $-\Delta w_{sE}(x) = -s^2 \cdot s^{-2}(\Delta w_E)(x/s) = 1$, and by uniqueness it is the torsion function of sE . Hence $m_{sE} = s^2 m_E$, and

$$T(sE) = \int_{sE} w_{sE} dx = \int_{sE} s^2 w_E(x/s) dx \stackrel{x=sy}{=} s^2 \int_E s^2 w_E(y) dy = s^4 T(E),$$

while $|sE|^2 = s^4|E|^2$; both terms on the right of (19) scale by s^4 , giving the second equality in (20), and the first follows from it via (19). (Directly: for a level $v = s^2v'$ one has $\{w_{sE} > v\} = s\{w_E > v'\}$, so $\mu_{sE}(s^2v') = s^2\mu_E(v')$, $P_{sE}(s^2v') = sP_E(v')$, $\mu'_{sE}(s^2v') = \mu'_E(v')$, whence $D_H^{sE}(s^2v') = s^2D_H^E(v')$ and $D_I^{sE}(s^2v') = s^2D_I^E(v')$; the change of variables $v = s^2v'$, $dv = s^2dv'$ then produces the factor $s^2 \cdot s^2 = s^4$.)

Scale invariance of δ_ .* From (19), $T(B_E) - T(E) = \frac{1}{4\pi} \int_0^{m_E} (D_H^E + D_I^E) dv$ and $T(B_E) = |E|^2/(8\pi)$, so dividing gives $\delta_*(E) = \frac{2}{|E|^2} \int_0^{m_E} (D_H^E + D_I^E) dv$, which is (21). By (20) numerator and denominator $T(B_E)$ both scale by s^4 , so $\delta_*(sE) = \delta_*(E)$. $\square$

Remark 3.7 (Consistency with the normalisation). Applied to $E = \Omega$ with $|\Omega| = \pi$, (19) returns (15): $\frac{1}{2}\pi^2 - 4\pi T(\Omega) = 4\pi\delta_T$, and $T(B_\Omega) = \pi^2/(8\pi) = \pi/8 = T(B_1)$. The two normalisations are related by $\delta_*(\Omega) = \delta_T/T(B_1) = \frac{8}{\pi}\delta_T$, in agreement with the global notation.

3.5. Talenti's comparison and the volume profile. To convert “small on most levels” into quantitative geometry we need the size of the level sets, i.e. the profile $\mu(v)$. This is furnished by Talenti's pointwise comparison with the model disk. Recall the unit-disk torsion function $u_{B_1}(x) = \frac{1}{4}(1 - |x|^2)$ and its distribution function

$$\mu_B(v) := |\{u_{B_1} > v\}| = \pi(1 - 4v)_+, \quad v \geq 0,$$

where $t_+ = \max\{t, 0\}$.

Theorem 3.8 (Talenti comparison for the profile). *Let u be the torsion function of Ω with $|\Omega| = \pi$. Then*

$$\mu(v) = |\{u > v\}| \leq \mu_B(v) = \pi(1 - 4v)_+ \quad \text{for all } v \geq 0. \quad (22)$$

Proof. We invoke Talenti's comparison theorem [11, Theorem 1] in the form proved there: if $u \in H_0^1(\Omega)$ solves $-\Delta u = f$ with $f \geq 0$, and u^* denotes the Schwarz (radially symmetric decreasing) rearrangement of u , while $\tilde{u}$ solves $-\Delta \tilde{u} = f^*$ in the ball Ω^* centred at the origin with $|\Omega^*| = |\Omega|$ and $\tilde{u} \in H_0^1(\Omega^*)$, then

$$u^*(x) \leq \tilde{u}(x) \quad \text{for all } x \in \Omega^*.$$

We apply this with $f \equiv 1$. Then $f^* \equiv 1$, $\Omega^* = B_1$ (since $|\Omega| = \pi = |B_1|$), and the comparison solution is the disk torsion function $\tilde{u} = u_{B_1}$, because $u_{B_1} = \frac{1}{4}(1 - |x|^2)$ solves $-\Delta u_{B_1} = 1$ in B_1 with zero boundary data and is its own rearrangement. Thus $u^* \leq u_{B_1}$ pointwise on B_1 .

Schwarz rearrangement is equimeasurable: $|\{u > v\}| = |\{u^* > v\}|$ for every v . Since $u^* \leq u_{B_1}$, we have $\{u^* > v\} \subseteq \{u_{B_1} > v\}$, hence $|\{u^* > v\}| \leq |\{u_{B_1} > v\}|$. Combining,

$$\mu(v) = |\{u > v\}| = |\{u^* > v\}| \leq |\{u_{B_1} > v\}| = \mu_B(v) = \pi(1 - 4v)_+,$$

where the last equality is $\{u_{B_1} > v\} = \{|x|^2 < 1 - 4v\}$, of area $\pi(1 - 4v)_+$. $\square$

Lemma 3.9 (Exact volume profile and the cap on the maximum). *The Saint-Venant deficit equals the L^1 gap between the disk profile and the domain profile over $(0, \frac{1}{4})$:*

$$\int_0^{1/4} (\mu_B(v) - \mu(v)) dv = \delta_T(\Omega), \quad (23)$$

the integrand $\mu_B(v) - \mu(v) \geq 0$ being nonnegative for every v . In particular the maximum is capped,

$$m = \max_{\Omega} u \leq \frac{1}{4}. \quad (24)$$

Proof. The maximum cap (24). Suppose, for contradiction, that $m > \frac{1}{4}$. Pick v with $\frac{1}{4} < v < m$. Then $\{u > v\} \neq \emptyset$ and is open (as u is continuous), so $\mu(v) = |\{u > v\}| > 0$; but $v > \frac{1}{4}$ gives $\mu_B(v) = \pi(1 - 4v)_+ = 0$, contradicting (22). Hence $m \leq \frac{1}{4}$. In particular $\mu(v) = 0$ for $v \geq \frac{1}{4} \geq m$.

The profile identity (23). Nonnegativity of the integrand on all of $(0, \frac{1}{4})$ is exactly (22). By the layer-cake formula and $m \leq \frac{1}{4}$,

$$T(\Omega) = \int_{\Omega} u \, dx = \int_0^{\infty} \mu(v) \, dv = \int_0^{1/4} \mu(v) \, dv,$$

the last step using $\mu(v) = 0$ for $v \geq \frac{1}{4}$. Likewise, by the same formula for the disk torsion function u_{B_1} ,

$$T(B_1) = \int_{B_1} u_{B_1} \, dx = \int_0^{\infty} \mu_B(v) \, dv = \int_0^{1/4} \pi(1 - 4v) \, dv = \pi \left[v - 2v^2 \right]_0^{1/4} = \pi \left(\frac{1}{4} - \frac{1}{8} \right) = \frac{\pi}{8}.$$

Subtracting,

$$\int_0^{1/4} (\mu_B(v) - \mu(v)) \, dv = T(B_1) - T(\Omega) = \frac{\pi}{8} - T(\Omega) = \delta_T(\Omega).$$

This is (23). (The computation also reconfirms independently that $T(B_1) = \pi/8$, hence that the deficit is well defined and non-negative: $\delta_T = \int_0^{1/4} (\mu_B - \mu) \geq 0$.) $\square$

Remark 3.10 (Two complementary decompositions of δ_T). We have written δ_T in two ways: as an integral over levels of the *shape* defects $D_H + D_I$ (Theorem 3.4, weight 4π), and as an integral over levels of the *volume* gap $\mu_B - \mu$ (Lemma 3.9, weight 1). The inset construction in Section 4 uses both: the first transfers the torsion deficit to a single level set, the second certifies that this level set captures almost all of the area of Ω .

3.6. A pointwise lower bound on $-\mu'$. The final consequence we extract is a clean pointwise estimate, used to quantify how fast the volume profile decreases.

Lemma 3.11 (Lower bound on the profile slope). *For a.e. $v \in (0, m)$,*

$$-\mu'(v) \geq 4\pi. \quad (25)$$

Proof. Fix $v \in (0, m)$ a regular value at which (13) holds (a.e. v). By Lemma 3.3, $D_H(v) \geq 0$ and $D_I(v) \geq 0$, i.e.

$$\mu(v)(-\mu'(v)) \geq P(v)^2 \quad \text{and} \quad P(v)^2 \geq 4\pi \mu(v).$$

Chaining these,

$$\mu(v)(-\mu'(v)) \geq P(v)^2 \geq 4\pi \mu(v).$$

Since $v < m$, the super-level set $E_v = \{u > v\}$ is nonempty and open, so $\mu(v) > 0$; dividing by $\mu(v) > 0$ gives $-\mu'(v) \geq 4\pi$. $\square$

Remark 3.12 (Sharpness and the model). All four statements of this section are sharp and are simultaneously saturated by the disk: for u_{B_1} one has $\mu = \mu_B$, $-\mu'_B(v) = 4\pi$ on $(0, \frac{1}{4})$, $D_H \equiv D_I \equiv 0$, $\delta_T = 0$, and $m = \frac{1}{4}$. Thus (25) is an equality for the model, and the level identity (15) reads $0 = 0$. The quantitative theory of the remaining sections measures the deviation from this rigid configuration.

4. THE INSET: REGULARITY, DEFICIT TRANSFER, LEVEL SELECTION

Throughout this section $\Omega \subset \mathbb{R}^2$ is open with $|\Omega| = \pi$, $u \in H_0^1(\Omega)$ is its torsion function ($-\Delta u = 1$ in Ω , $u = 0$ on $\partial\Omega$), $m := \max_\Omega u$, and $\delta_T = \delta_T(\Omega) = \frac{\pi}{8} - T(\Omega) \geq 0$ is the Saint-Venant deficit. We write $E_v := \{u > v\}$ for the open super-level set, $\mu(v) := |E_v|$, $P(v) := P(E_v)$, and recall the two level deficits of Definition 3.2,

$$D_H(v) = \mu(v)(-\mu'(v)) - P(v)^2 \geq 0, \quad D_I(v) = P(v)^2 - 4\pi\mu(v) \geq 0$$

(Lemma 3.3), the level-set deficit identity $\int_0^m (D_H + D_I) dv = 4\pi\delta_T$ (Theorem 3.4), the Talenti profile $\mu_B(v) := \pi(1 - 4v)_+$ with $\mu \leq \mu_B$ and the profile identity $\int_0^{1/4} (\mu_B - \mu) dv = \delta_T$ (Theorem 3.8, Lemma 3.9), the bound $-\mu'(v) \geq 4\pi$ a.e. (Lemma 3.11), and the flux/coarea identities $\int_{\{u=v\}} |\nabla u| d\mathcal{H}^1 = \mu(v)$, $-\mu'(v) = \int_{\{u=v\}} |\nabla u|^{-1} d\mathcal{H}^1$ for a.e. regular v (Theorem 2.6, Lemma 2.7). By Sard's theorem (Theorem 2.16) and Lemma 2.17 the non-regular values of u form a Lebesgue-null set, and for every regular $v \in (0, m)$ the set E_v has finite perimeter with $\mathcal{H}^1$ -rectifiable boundary $\{u = v\}$ (Theorem 2.13). Absolute constants are denoted C (value may change) or named $(A_0, \delta_\star)$; $\delta_\star > 0$ is a small absolute constant, fixed at the end of the section. We set $\tau := \sqrt{\delta_T}$ only at the very end; until then τ is a free parameter in $(0, \frac{1}{16})$.

4.1. No interior holes. We first record a topological regularity property of the super-level sets, which is a direct consequence of the superharmonicity of u ($-\Delta u = 1 \geq 0$) and the minimum principle. Recall (Theorem 2.13) that $u \in C^2(\Omega) \cap C(\bar{\Omega})$, $u > 0$ in Ω and $u = 0$ on $\partial\Omega$.

Lemma 4.1 (No interior holes). *For every $v \in (0, m)$ the following hold.*

- (i) *Every connected component K of the open set $\{x \in \Omega : u(x) < v\}$ has $\bar{K} \cap \partial\Omega \neq \emptyset$.*
- (ii) *For every regular value v , each connected component E of $E_v = \{u > v\}$ has no interior holes: every bounded connected component H of $\mathbb{R}^2 \setminus \bar{E}$ satisfies $\bar{H} \not\subset \Omega$; in particular $\mathbb{R}^2 \setminus \bar{E}$ is connected up to sets touching $\partial\Omega$, and the inset defined below is simply connected modulo $\partial\Omega$.*

Proof. (i). Suppose, for contradiction, that some component K of $\{u < v\}$ satisfies $\bar{K} \Subset \Omega$, i.e. $\bar{K}$ is a compact subset of Ω not meeting $\partial\Omega$. We claim $u \equiv v$ on ∂K . Let $x_0 \in \partial K$. Since $\bar{K} \subset \Omega$ we have $x_0 \in \Omega$, so u is continuous at x_0 . If $u(x_0) < v$ then by continuity $u < v$ on a ball around x_0 , which would be a connected subset of $\{u < v\}$ meeting K , hence contained in K , placing x_0 in the interior of K — contradicting $x_0 \in \partial K$. If $u(x_0) > v$ then by continuity $u > v$ near x_0 , so a punctured neighbourhood of x_0 is disjoint from $\{u < v\} \supset K$, contradicting $x_0 \in \partial K$ (as K accumulates at x_0). Hence $u(x_0) = v$, proving $u = v$ on ∂K .

Now u is continuous on the compact set $\bar{K}$ and attains its minimum there; since $u < v$ at any interior point of K and $u = v$ on ∂K , the minimum is attained at an interior point and equals $\min_{\bar{K}} u < v = \min_{\partial K} u$. But $-\Delta u = 1 \geq 0$ in Ω makes u superharmonic, and the (weak) minimum principle for superharmonic functions in C^2 [7, Theorem 3.1] gives $\min_{\bar{K}} u = \min_{\partial K} u$. This contradiction proves (i).

(ii). Let v be regular and E a component of E_v , and suppose H is a bounded component of $\mathbb{R}^2 \setminus \bar{E}$ with $\bar{H} \subset \Omega$. Then H is an open subset of Ω with $\bar{H} \Subset \Omega$, and we consider the two open sets $W^- := H \cap \{u < v\}$ and $W^+ := H \cap \{u > v\}$, both contained in H , hence both with closures contained in $\bar{H} \Subset \Omega$.

We first note that every connected component G of W^- (resp. of W^+) is in fact a connected component of $\{u < v\}$ (resp. of $\{u > v\}$), and is contained in H . Indeed, G is connected and contained in the open set $\{u \neq v\}$, so it lies in a unique component $\hat{G}$ of $\{u < v\}$ (resp. $\{u > v\}$). If $\hat{G}$ left H it would, being connected and meeting H , have to cross ∂H ; but $\partial H \subset \bar{E}$ and on $\bar{E}$ one has $u \geq v$, while points of ∂H with $u > v$ lie in the open set E (a component of E_v , hence open) which is disjoint from $\hat{G}$ when $\hat{G} \subset W^-$, and equals a different component of E_v disjoint from E when $\hat{G} \subset W^+$ (as $\hat{G} \cap E \subset W^+ \cap E \subset H \cap E = \emptyset$). In either case $\hat{G}$ cannot meet ∂H , so $\hat{G} \subset H$ and $\hat{G} = G$. Thus each component of W^- is a connected component of $\{u < v\}$ whose closure lies in $\bar{H} \Subset \Omega$ and so does not meet $\partial\Omega$; by (i) this forces $W^- = \emptyset$. Each connected component E' of W^+ is a connected component of $E_v = \{u > v\}$ with $\bar{E'} \Subset \Omega$; on it $u = v$ on $\partial E'$ and $u > v$ inside, so u attains an interior maximum strictly above its boundary value. But u is superharmonic ($-\Delta u = 1 \geq 0$), so $-u$ is subharmonic and satisfies the maximum principle [7, Theorem 3.1]: $\max_{\bar{E'}} u = \max_{\partial E'} u = v$, contradicting $u > v$ inside. Hence $W^+ = \emptyset$ as well.

Therefore $u \equiv v$ on the open set H , which has positive Lebesgue measure; but v is a regular value, so $\{u = v\}$ is an $\mathcal{H}^1$ -rectifiable curve, of zero Lebesgue measure. This contradicts $|H| > 0$, so no such H exists. $\square$

Remark 4.2. Part (ii) uses both the minimum principle (via (i), excluding holes that are sub-level islands) and the maximum principle (excluding super-level islands separated from E). Only two consequences are used downstream: that $\tilde{E}$ is connected with no interior holes (so that the trapping annulus and the ray-crossing counts of Theorem 5.8 are well posed), and the measure-theoretic fact $|\{u = \hat{v}\}| = 0$ used repeatedly to identify symmetric differences. A reader content with these may keep (i) and the line “ $u \equiv v$ on H with $|H| > 0$ contradicts $|\{u = v\}| = 0$ ” as the whole of (ii).

4.2. Deficit transfer to super-level sets. The torsion function of any super-level set E_v is $u - v$, and the level-set deficit identity, being homogeneous of degree four, transfers the scale-invariant deficit from Ω to E_v as a tail of the positive-integrand identity of Theorem 3.4.

Lemma 4.3 (Deficit transfer). *For every regular $v \in (0, m)$,*

$$\delta_*(E_v) \leq \frac{\pi^2}{\mu(v)^2} \delta_*(\Omega). \quad (26)$$

Proof. On the open set E_v the function $w := u - v$ satisfies $-\Delta w = 1$ in E_v , $w = 0$ on $\partial E_v \subset \{u = v\}$, and $w > 0$ in E_v ; hence w is the torsion function of E_v and $T(E_v) = \int_{E_v} w = \int_{E_v} (u - v)$. Its super-level sets are $\{w > s\} = \{u > v + s\} = E_{v+s}$ for $s \in (0, m - v)$, so the perimeter and distribution functions of w at height s coincide with those of u at height $v + s$; in particular the two level deficits of w at height s equal $D_H(v + s)$ and $D_I(v + s)$.

We now apply the derivation of the level-set deficit identity (Theorem 3.4, eq. (15)) to w on E_v ; the derivation there uses only the flux/coarea identities, the absolute continuity of the distribution function, and the layer-cake formula, none of which requires the area to equal π . For a domain D of area $|D|$ with torsion function having maximum m_D and level deficits D_H^D, D_I^D it yields

$$\int_0^{m_D} (D_H^D + D_I^D) ds = \frac{1}{2}|D|^2 - 4\pi T(D) = 4\pi \left(\frac{|D|^2}{8\pi} - T(D) \right) = 4\pi (T(B_D) - T(D)),$$

where B_D is the disc of area $|D|$, $T(B_D) = |D|^2/(8\pi)$; the middle equality is exactly the computation in the proof of Theorem 3.4 with $\frac{1}{2}|D|^2$ in place of $\frac{1}{2}\pi^2$. Taking $D = E_v$, $m_{E_v} = m - v$, and using the identification of the level deficits of w with $D_H(v + \cdot), D_I(v + \cdot)$,

$$4\pi(T(B_{E_v}) - T(E_v)) = \int_0^{m-v} (D_H(v+s) + D_I(v+s)) ds = \int_v^m (D_H + D_I) dv' \leq \int_0^m (D_H + D_I) dv' = 4\pi\delta_T,$$

the inequality by non-negativity of the integrand (Lemma 3.3). Hence $T(B_{E_v}) - T(E_v) \leq \delta_T$. Dividing by $T(B_{E_v}) = \mu(v)^2/(8\pi)$ and using $\delta_*(\Omega) = \frac{8}{\pi}\delta_T$, i.e. $\delta_T = \frac{\pi}{8}\delta_*(\Omega)$,

$$\delta_*(E_v) = \frac{T(B_{E_v}) - T(E_v)}{T(B_{E_v})} \leq \frac{\delta_T}{\mu(v)^2/(8\pi)} = \frac{8\pi\delta_T}{\mu(v)^2} = \frac{8\pi}{\mu(v)^2} \cdot \frac{\pi}{8} \delta_*(\Omega) = \frac{\pi^2}{\mu(v)^2} \delta_*(\Omega). \quad \square$$

The content of (26) is that the *scale-invariant* torsion deficit of every super-level set is comparable to that of Ω (a tail of a positive integral), even at levels near $\partial\Omega$ where the single-level deficits $D_H(v), D_I(v)$ are individually inflated. This is the mechanism that carries the small torsion deficit onto the inset.

4.3. Selection of a good level.

Lemma 4.4 (Good level). *Let $A_0 := 16\pi$. For every $\tau \in (0, \frac{1}{16})$ there is a regular value $\hat{v} \in (\tau, 2\tau)$ of u with*

$$D_H(\hat{v}) + D_I(\hat{v}) \leq A_0 \frac{\delta_T}{\tau}, \quad \mu_B(\hat{v}) - \mu(\hat{v}) \leq A_0 \frac{\delta_T}{\tau}. \quad (27)$$

Proof. Set $I := (\tau, 2\tau)$, of length $|I| = \tau$, and note $I \subset (0, \frac{1}{8}) \subset (0, \frac{1}{4})$. By Theorem 3.4 and non-negativity, $\int_I (D_H + D_I) dv \leq \int_0^m (D_H + D_I) dv = 4\pi\delta_T$. By Chebyshev's inequality,

$$|\{v \in I : D_H(v) + D_I(v) > A_0\delta_T/\tau\}| \leq \frac{\int_I (D_H + D_I)}{A_0\delta_T/\tau} \leq \frac{4\pi\delta_T}{A_0\delta_T/\tau} = \frac{4\pi\tau}{A_0}.$$

By Lemma 3.9, $\mu_B - \mu \geq 0$ and $\int_I (\mu_B - \mu) dv \leq \int_0^{1/4} (\mu_B - \mu) dv = \delta_T$, so again by Chebyshev,

$$|\{v \in I : \mu_B(v) - \mu(v) > A_0\delta_T/\tau\}| \leq \frac{\delta_T}{A_0\delta_T/\tau} = \frac{\tau}{A_0}.$$

With $A_0 = 16\pi$, the union of these two “bad” sets has measure at most $\frac{4\pi\tau}{A_0} + \frac{\tau}{A_0} = \frac{\tau}{4} + \frac{\tau}{16\pi} < \tau = |I|$. Adding the null set of non-regular values (Theorem 2.16, Lemma 2.17) does not change this. Hence the set of regular $\hat{v} \in I$ satisfying both bounds in (27) has positive measure, and in particular is nonempty. $\square$

From now on $\hat{v} \in (\tau, 2\tau)$ denotes a good regular level as in Lemma 4.4, and we write $\hat{v} < \frac{1}{4}$ (since $\hat{v} < 2\tau < \frac{1}{8}$), so that $\mu_B(\hat{v}) = \pi(1 - 4\hat{v})$.

4.4. Closeness of the super-level set.

Lemma 4.5 (Closeness). *At the level $\hat{v}$ of Lemma 4.4,*

$$|\Omega \triangle \{u > \hat{v}\}| = \pi - \mu(\hat{v}) \leq 8\pi\tau + A_0 \frac{\delta_T}{\tau}. \quad (28)$$

Proof. Since $E_{\hat{v}} = \{u > \hat{v}\} \subset \Omega$, the symmetric difference is $\Omega \setminus E_{\hat{v}} = \{x \in \Omega : u(x) \leq \hat{v}\}$, of measure $\pi - \mu(\hat{v})$ (the level set $\{u = \hat{v}\}$ is Lebesgue-null). Split

$$\pi - \mu(\hat{v}) = (\pi - \mu_B(\hat{v})) + (\mu_B(\hat{v}) - \mu(\hat{v})).$$

The first term is $\pi - \pi(1 - 4\hat{v}) = 4\pi\hat{v} \leq 4\pi(2\tau) = 8\pi\tau$, using $\hat{v} < 2\tau$; the second is $\leq A_0\delta_T/\tau$ by (27). Adding gives (28). $\square$

We now fix $\delta_\star \in (0, \frac{1}{256}]$ small enough that, whenever $\delta_T \leq \delta_\star$ and $\tau \in [\sqrt{\delta_T}, \frac{1}{16})$ with $\delta_T/\tau \leq \sqrt{\delta_T}$ (e.g. at the optimal choice $\tau = \sqrt{\delta_T}$, where the right side of (28) equals $(8\pi + A_0)\sqrt{\delta_T}$), one has

$$8\pi\tau + A_0 \frac{\delta_T}{\tau} \leq \frac{\pi}{2}, \quad \text{hence} \quad \mu(\hat{v}) \geq \frac{\pi}{2}. \quad (29)$$

Concretely, at $\tau = \sqrt{\delta_T}$ this requires $(8\pi + A_0)\sqrt{\delta_\star} \leq \pi/2$, i.e. $\delta_\star \leq \pi^2/(4(8\pi + A_0)^2)$; with $A_0 = 16\pi$ this is $\delta_\star \leq 1/(2304)$, comfortably compatible with $\delta_\star \leq \frac{1}{256}$ after taking the smaller value. We keep (29) as a standing hypothesis for the remainder of the section (it holds at the optimised scale $\tau = \sqrt{\delta_T}$); the value of $\hat{v}, \tau$ is otherwise left symbolic and optimised in Theorem 9.1.

4.5. One connected inset: the spillover is higher order. The super-level set $\{u > \hat{v}\}$ may have several components, but the total mass of all but the largest is quadratically small in $D_I(\hat{v})$, by a multi-component refinement of the isoperimetric inequality.

Lemma 4.6 (Spillover). *Assume (29). Let the connected components of $\{u > \hat{v}\}$ have areas $A_1 \geq A_2 \geq \dots > 0$, let $\tilde{E}$ be the component of area A_1 , and set $S := \sum_{i \geq 2} A_i = \mu(\hat{v}) - A_1$ for the total spillover. Then $A_1 \geq \pi/4$ and*

$$|\{u > \hat{v}\} \setminus \tilde{E}| = S \leq \frac{D_I(\hat{v})^2}{16\pi^3} \leq \frac{A_0^2}{16\pi^3} \frac{\delta_T^2}{\tau^2}. \quad (30)$$

Proof. Multi-component isoperimetric inequality. Perimeter is additive over the (essentially disjoint) components and the planar isoperimetric inequality (Theorem 2.9) applies to each, so $P(\hat{v}) = \sum_i P_i \geq \sum_i 2\sqrt{\pi A_i}$ with P_i the perimeter of the i -th component. Squaring and using $(\sum_i \sqrt{A_i})^2 = \sum_i A_i + 2 \sum_{i < j} \sqrt{A_i A_j}$,

$$D_I(\hat{v}) = P(\hat{v})^2 - 4\pi \sum_i A_i \geq 4\pi \left[\left(\sum_i \sqrt{A_i} \right)^2 - \sum_i A_i \right] = 8\pi \sum_{i < j} \sqrt{A_i A_j}. \quad (31)$$

Lower bound on A_1 . Since $A_j \leq A_1$ we have $\sqrt{A_j} = A_j/\sqrt{A_j} \geq A_j/\sqrt{A_1}$ for $j \geq 2$, hence $\sum_{j \geq 2} \sqrt{A_j} \geq S/\sqrt{A_1}$ and

$$\sum_i \sqrt{A_i} \geq \sqrt{A_1} + \frac{S}{\sqrt{A_1}} = \frac{A_1 + S}{\sqrt{A_1}} = \frac{\mu(\hat{v})}{\sqrt{A_1}}.$$

Combining with (31) (in the form $D_I \geq 4\pi[(\sum \sqrt{A_i})^2 - \mu(\hat{v})]$),

$$D_I(\hat{v}) \geq 4\pi \left(\frac{\mu(\hat{v})^2}{A_1} - \mu(\hat{v}) \right) = 4\pi \mu(\hat{v}) \frac{\mu(\hat{v}) - A_1}{A_1} = 4\pi \mu(\hat{v}) \frac{S}{A_1}. \quad (32)$$

Suppose, for contradiction, that $A_1 < \pi/4$. By (29), $\mu(\hat{v}) \geq \pi/2 > A_1$, so there are at least two components and $S = \mu(\hat{v}) - A_1 > \pi/2 - \pi/4 = \pi/4$. Then (32) gives

$$D_I(\hat{v}) > 4\pi \cdot \frac{\pi}{2} \cdot \frac{S}{\pi/4} = 8\pi S > 8\pi \cdot \frac{\pi}{4} = 2\pi^2.$$

But $D_I(\hat{v}) \leq A_0 \delta_T / \tau \leq 8\pi\tau + A_0 \delta_T / \tau \leq \pi/2 < 2\pi^2$ by (27) and (29), a contradiction. Hence $A_1 \geq \pi/4$.

Spillover bound. From (31),

$$D_I(\hat{v}) \geq 8\pi \sum_{i < j} \sqrt{A_i A_j} \geq 8\pi \sqrt{A_1} \sum_{j \geq 2} \sqrt{A_j} \geq 8\pi \sqrt{A_1} \sqrt{\sum_{j \geq 2} A_j} = 8\pi \sqrt{A_1} S,$$

where the middle step keeps only the pairs $(1, j)$ and the last step uses $\sum_{j \geq 2} \sqrt{A_j} \geq \sqrt{\sum_{j \geq 2} A_j}$ (superadditivity of $t \mapsto \sqrt{t}$ on non-negative terms). Squaring and using $A_1 \geq \pi/4$,

$$S \leq \frac{D_I(\hat{v})^2}{64\pi^2 A_1} \leq \frac{D_I(\hat{v})^2}{64\pi^2 \cdot (\pi/4)} = \frac{D_I(\hat{v})^2}{16\pi^3}.$$

The final bound in (30) follows from $D_I(\hat{v}) \leq A_0 \delta_T / \tau$ (Lemma 4.4). $\square$

Definition 4.7 (The inset). Assume $\delta_T \leq \delta_*$, τ subject to (29), and let $\hat{v} \in (\tau, 2\tau)$ be a good regular level (Lemma 4.4). The *inset* is the connected component $\tilde{E}$ of $\{u > \hat{v}\}$ of largest measure, i.e. the component of area A_1 in Lemma 4.6. We write

$$\mu_E := |\tilde{E}| = A_1 = \mu(\hat{v}) - S \geq \frac{\pi}{4}, \quad \hat{r} := \sqrt{\mu_E / \pi}, \quad \delta_{\text{iso}} := \delta_{\text{iso}}(\tilde{E}) = \frac{P(\tilde{E})}{2\sqrt{\pi\mu_E}} - 1,$$

and $\text{Exc} := P(\tilde{E}) - 2\pi\hat{r} = 2\pi\hat{r}\delta_{\text{iso}}$.

By Lemma 4.1 (ii), $\tilde{E}$ is connected and has no interior holes.

4.6. Controls on the inset. We collect the three controls of the inset used in the subsequent sections.

Proposition 4.8 (Inset controls). *Under Definition 4.7 there is an absolute constant C such that*

$$\delta_{\text{iso}}(\tilde{E}) \leq \frac{D_I(\hat{v}) + 4\pi S}{8\pi\mu_E} \leq C \frac{\delta_T}{\tau}, \quad (33)$$

$$\delta_*(\tilde{E}) \leq \frac{\pi^2}{\mu_E^2} \delta_*(\Omega) + \frac{S^2}{\mu_E^2} \leq C \delta_*(\Omega), \quad (34)$$

$$|\Omega \triangle \tilde{E}| \leq 8\pi\tau + A_0 \frac{\delta_T}{\tau} + \frac{A_0^2}{16\pi^3} \frac{\delta_T^2}{\tau^2}. \quad (35)$$

In particular, with the optimal choice $\tau = \sqrt{\delta_T}$ of Theorem 9.1, $\delta_{\text{iso}}(\tilde{E}) \leq C\sqrt{\delta_T}$, $\delta_*(\tilde{E}) \leq C\delta_*(\Omega)$ and $|\Omega \triangle \tilde{E}| \leq C\sqrt{\delta_T}$.

Proof. (33). Since $\partial\tilde{E} \subset \{u = \hat{v}\}$ and perimeter is additive over components, $P(\tilde{E}) \leq P(\hat{v})$, hence

$$D_I(\tilde{E}) := P(\tilde{E})^2 - 4\pi\mu_E \leq P(\hat{v})^2 - 4\pi\mu_E = (4\pi\mu(\hat{v}) + D_I(\hat{v})) - 4\pi\mu_E = D_I(\hat{v}) + 4\pi S,$$

using $P(\hat{v})^2 = 4\pi\mu(\hat{v}) + D_I(\hat{v})$ and $\mu(\hat{v}) - \mu_E = S$. From the definition of δ_{iso} , $(1 + \delta_{\text{iso}}(\tilde{E}))^2 = P(\tilde{E})^2 / (4\pi\mu_E) = 1 + D_I(\tilde{E}) / (4\pi\mu_E)$, so by $\sqrt{1+t} - 1 \leq t/2$,

$$\delta_{\text{iso}}(\tilde{E}) \leq \frac{D_I(\tilde{E})}{8\pi\mu_E} \leq \frac{D_I(\hat{v}) + 4\pi S}{8\pi\mu_E}.$$

By (30), $4\pi S \leq 4\pi D_I(\hat{v})^2 / (16\pi^3) = D_I(\hat{v})^2 / (4\pi^2)$; and $D_I(\hat{v}) \leq A_0 \delta_T / \tau \leq \pi/2$ by (27)–(29), so $4\pi S \leq D_I(\hat{v}) \cdot \frac{D_I(\hat{v})}{4\pi^2} \leq \frac{1}{8\pi} D_I(\hat{v}) \leq D_I(\hat{v})$. Hence $D_I(\hat{v}) + 4\pi S \leq 2D_I(\hat{v}) \leq 2A_0 \delta_T / \tau$, and with

$$\mu_E \geq \pi/4,$$

$$\delta_{\text{iso}}(\tilde{E}) \leq \frac{2A_0\delta_T/\tau}{8\pi \cdot (\pi/4)} = \frac{A_0}{\pi^2} \frac{\delta_T}{\tau} =: C \frac{\delta_T}{\tau}.$$

(34). We bound $\delta_*(\tilde{E})$ directly, using only the deficit transfer of Lemma 4.3 and the exact additivity of torsion over connected components; no appeal to Lemma 7.5 is needed (see Remark 4.9). Let $C_2, C_3, \dots$ be the spillover components of $E_{\hat{v}}$, of areas $A_2, A_3, \dots$ with $\sum_{i \geq 2} A_i = S$. On each component C of $E_{\hat{v}} = \{u > \hat{v}\}$ the function $u - \hat{v}$ solves $-\Delta(u - \hat{v}) = 1$, vanishes on $\partial C \subset \{u = \hat{v}\}$, and is positive inside; hence $u - \hat{v}$ is the torsion function of C and $T(C) = \int_C (u - \hat{v})$. As $u - \hat{v}$ is shared and the components are pairwise disjoint, torsion is *exactly additive*:

$$T(E_{\hat{v}}) = \int_{E_{\hat{v}}} (u - \hat{v}) = \int_{\tilde{E}} (u - \hat{v}) + \sum_{i \geq 2} \int_{C_i} (u - \hat{v}) = T(\tilde{E}) + \sum_{i \geq 2} T(C_i). \quad (36)$$

By the Saint-Venant inequality applied to each spillover component (Proposition 3.6, $T(B_E) - T(E) \geq 0$, with $T(B_{C_i}) = A_i^2/(8\pi)$),

$$\sum_{i \geq 2} T(C_i) \leq \sum_{i \geq 2} \frac{A_i^2}{8\pi} \leq \frac{1}{8\pi} \left(\sum_{i \geq 2} A_i \right)^2 = \frac{S^2}{8\pi}, \quad (37)$$

the middle inequality because $\sum A_i^2 \leq (\sum A_i)^2$ for non-negative terms. Combining (36)–(37) with $\mu_E \leq \mu(\hat{v})$,

$$T(B_{\tilde{E}}) - T(\tilde{E}) = \frac{\mu_E^2}{8\pi} - T(E_{\hat{v}}) + \sum_{i \geq 2} T(C_i) \leq \frac{\mu_E^2 - \mu(\hat{v})^2}{8\pi} + (T(B_{E_{\hat{v}}}) - T(E_{\hat{v}})) + \frac{S^2}{8\pi} \leq \delta_T + \frac{S^2}{8\pi},$$

where we used $T(B_{\tilde{E}}) = \mu_E^2/(8\pi)$, $T(B_{E_{\hat{v}}}) = \mu(\hat{v})^2/(8\pi)$, $\mu_E^2 - \mu(\hat{v})^2 \leq 0$, and the bound $T(B_{E_{\hat{v}}}) - T(E_{\hat{v}}) \leq \delta_T$ established in the proof of Lemma 4.3. Dividing by $T(B_{\tilde{E}}) = \mu_E^2/(8\pi)$ and using $\delta_T = \frac{\pi}{8}\delta_*(\Omega)$,

$$\delta_*(\tilde{E}) \leq \frac{8\pi\delta_T}{\mu_E^2} + \frac{S^2}{\mu_E^2} = \frac{\pi^2}{\mu_E^2} \delta_*(\Omega) + \frac{S^2}{\mu_E^2}. \quad (38)$$

This is the first inequality in (34) up to the higher-order term S^2/μ_E^2 . The latter is negligible: by (30), $S \leq A_0^2\delta_T^2/(16\pi^3\tau^2)$, and with the standing hypothesis (29) (in particular $A_0\delta_T/\tau \leq \pi/2$, i.e. $\delta_T/\tau \leq 1/32$) one has $S \leq \frac{A_0^2}{16\pi^3}(\delta_T/\tau) \cdot \delta_T \leq \frac{A_0^2}{16\pi^3 \cdot 32} \delta_T \leq \delta_T$; hence, with $\mu_E \geq \pi/4$,

$$\frac{S^2}{\mu_E^2} \leq \frac{S \cdot \delta_T}{(\pi/4)^2} \leq \frac{16}{\pi^2} \delta_T^2 \leq \frac{16}{\pi^2} \delta_T \cdot \frac{\pi}{8} \delta_*(\Omega) = \frac{2}{\pi} \delta_T \delta_*(\Omega) \leq \delta_*(\Omega),$$

the last step for $\delta_T \leq \delta_* \leq \pi/2$. Thus, with $\mu_E \geq \pi/4$, (38) gives $\delta_*(\tilde{E}) \leq \frac{\pi^2}{\mu_E^2} \delta_*(\Omega) + \delta_*(\Omega) \leq (16 + 1)\delta_*(\Omega) \leq C \delta_*(\Omega)$, which is (34).

(35). By Lemma 4.5 and Lemma 4.6,

$$|\Omega \Delta \tilde{E}| = (\pi - \mu(\hat{v})) + S \leq \left(8\pi\tau + A_0 \frac{\delta_T}{\tau} \right) + \frac{A_0^2}{16\pi^3} \frac{\delta_T^2}{\tau^2}.$$

For the final assertions, set $\tau = \sqrt{\delta_T}$: then $\delta_T/\tau = \sqrt{\delta_T}$, so $\delta_{\text{iso}}(\tilde{E}) \leq C\sqrt{\delta_T}$ by (33); the spillover term $\delta_T^2/\tau^2 = \delta_T$ is higher order, so $|\Omega \Delta \tilde{E}| \leq (8\pi + A_0)\sqrt{\delta_T} + C\delta_T \leq C\sqrt{\delta_T}$; and (34) is independent of τ . $\square$

Remark 4.9 (Self-contained deficit transfer; no forward dependence). The bound (34) is proved above without any appeal to the cutting principle Lemma 7.5 of Section 7: the passage from $E_{\hat{v}}$ to its largest component $\tilde{E}$ is handled by the exact additivity of torsion over components (36) and the Saint-Venant bound (37) on the spillover torsion, both internal to Sections 3 and 4. There is therefore *no* circular dependence on Section 7 in this section. (One could alternatively invoke Lemma 7.5; but its precise hypothesis — that the excised set lie in a thin concentric collar of $\tilde{E}$ — is *not* met by the spillover, whose components are in general not contained in the trapping annulus of $\tilde{E}$. The additivity argument avoids this issue entirely, which is why we prefer it.) For the record, Lemma 7.5 itself rests only on Green-function monotonicity and a boundary-layer barrier, so even the optional route would be acyclic.

Remark 4.10 (The scales, recorded). With $\tau = \sqrt{\delta_T}$ (Theorem 9.1) the inset $\tilde{E}$ satisfies $\delta_{\text{iso}} = \delta_{\text{iso}}(\tilde{E}) \leq C\sqrt{\delta_T}$ and $\text{Exc} = 2\pi\hat{r}\delta_{\text{iso}} \leq C\sqrt{\delta_T}$. Feeding δ_{iso} into the sharp quantitative isoperimetric inequality (Theorem 2.11) gives the ball-distance $a := |\tilde{E} \Delta B_{\hat{r}}(z)| \leq C_F \sqrt{\delta_{\text{iso}}} \mu_E \leq C\delta_T^{1/4}$ at the optimal centre z . The trapping width then satisfies $\eta \leq C(\text{Exc} + a)^{1/2} \leq C\delta_T^{1/8}$ (Theorem 5.8), an *absolute* small amplitude. These are the inputs to the annulus and fold sections; the deficit $\delta_*(\tilde{E}) \leq C\delta_*(\Omega)$ and the closeness $|\Omega \Delta \tilde{E}| \leq C\sqrt{\delta_T}$ are the inputs to the conclusion. We emphasise that closeness is carried by the *level* $\hat{v} \sim \sqrt{\delta_T}$ through Lemma 4.5, not by the amplitude η ; and that the deficit transfer Lemma 4.3 delivers $\delta_*(\tilde{E}) \lesssim \delta_*(\Omega)$ without recourse to the inflated single-level deficits.

5. THE TRAPPING ANNULUS

In this section we prove that the connected inset $\tilde{E}$ produced in Section 4 is trapped between two concentric circles whose radii differ by an *absolutely small* amount, controlled by the perimeter excess and the L^1 distance to a ball. The decisive point is that the width of the trapping annulus is bounded *without any second-order information* on $\partial\tilde{E}$: no interior or exterior ball condition, no C^2 regularity, no Magnanini–Poggesi oscillation estimate. The only inputs are the connectedness and the no-hole structure of $\tilde{E}$ (Lemma 4.1), the planar isoperimetric deficit, and the closeness to a ball furnished by Theorem 2.11.

The mechanism is purely geometric: the boundary of a connected, hole-free set must cross every circle of intermediate radius at least twice; summed against the radial coordinate through the coarea formula, this forces a lower bound on the *angular* part of the perimeter, which a single Cauchy–Schwarz inequality converts into the deficit. We work throughout with the good representative of $\tilde{E}$ described in Theorem 2.13 and Lemma 2.17: since $\hat{v}$ is a regular value of the (real-analytic, in Ω) torsion function u and $\tilde{E}$ is one connected component of $\{u > \hat{v}\}$, the set $\tilde{E}$ is a bounded open connected subset of Ω , with $\tilde{\tilde{E}} \subset\subset \Omega$, whose topological boundary

$$\partial\tilde{E} = \{u = \hat{v}\} \cap \partial\tilde{E}$$

is a compact, $\mathcal{H}^1$ -rectifiable curve along which $\nabla u \neq 0$; in particular $\partial\tilde{E}$ coincides $\mathcal{H}^1$ -a.e. with the reduced and the essential boundary, and $\mathcal{H}^1(\partial\tilde{E}) = P(\tilde{\tilde{E}})$. The plane decomposes as the disjoint union

$$\mathbb{R}^2 = \tilde{\tilde{E}} \sqcup \partial\tilde{E} \sqcup (\mathbb{R}^2 \setminus \tilde{\tilde{E}}), \quad (39)$$

the first and last sets being open. We record the precise topological consequence of the no-hole property that we shall use.

Lemma 5.1 (The complement of the inset is connected). *The open set $U := \mathbb{R}^2 \setminus \widetilde{\widetilde{E}}$ is connected, and $\partial U = \partial \widetilde{E}$.*

Proof. Since $\widetilde{\widetilde{E}} \subset\subset \Omega$ is compact, U is the complement of a compact set and so has exactly one unbounded connected component U_∞ (the points of large modulus form a connected set lying in U). Suppose U had another (necessarily bounded) component H . By Lemma 4.1 no bounded component of $\mathbb{R}^2 \setminus \widetilde{\widetilde{E}}$ is compactly contained in Ω , so $\overline{H}$ meets $\partial\Omega$; pick $q \in \overline{H} \cap \partial\Omega$. As $\widetilde{\widetilde{E}} \subset\subset \Omega$, the point q and the entire ray from q heading away from the bounded set $\widetilde{\widetilde{E}}$ lie in U and reach arbitrarily large modulus, hence lie in U_∞ ; thus $q \in \overline{U_\infty}$ and, H being open and adjacent to q , H meets U_∞ , forcing $H = U_\infty$, a contradiction with H bounded. Therefore $U = U_\infty$ is connected. Finally $\partial U = \partial(\mathbb{R}^2 \setminus \widetilde{\widetilde{E}}) = \partial \widetilde{\widetilde{E}} = \partial \widetilde{E}$, the last equality because $\widetilde{E}$ equals the interior of its closure for the good representative. $\square$

Throughout, $z \in \mathbb{R}^2$ denotes the centre of an optimal ball realising the Fraenkel asymmetry of $\widetilde{E}$; we take $z \in \widetilde{E}$ (an interior point), which holds for the inset as shown in Remark 5.10. With $\hat{r} := \sqrt{|\widetilde{E}|/\pi} \in [\frac{1}{2}, 1]$ (the bounds follow from $\pi/4 \leq |\widetilde{E}| \leq \pi$, Lemma 4.6)

$$a := |\widetilde{E} \triangle B_{\hat{r}}(z)| = \min_{x_0 \in \mathbb{R}^2} |\widetilde{E} \triangle B_{\hat{r}}(x_0)| = \mathcal{A}(\widetilde{E}) |\widetilde{E}|. \quad (40)$$

We write $e_r := (x - z)/|x - z|$ for the radial unit field (defined on $\mathbb{R}^2 \setminus \{z\}$), $e_\theta := e_r^\perp$ for the orthogonal angular field, ν for the outward unit normal on the rectifiable boundary, and

$$\begin{aligned} \nu_r &:= \nu \cdot e_r, & \nu_\theta &:= \nu \cdot e_\theta, & \nu_r^2 + \nu_\theta^2 &= 1 \quad (\mathcal{H}^1\text{-a.e. on } \partial \widetilde{E}), \\ P_r &:= \int_{\partial \widetilde{E}} |\nu_r| \, d\mathcal{H}^1, & P_\theta &:= \int_{\partial \widetilde{E}} |\nu_\theta| \, d\mathcal{H}^1, & 0 \leq P_r, P_\theta &\leq P(\widetilde{E}). \end{aligned}$$

Finally we record the standing scaling assumption of the construction, $\delta_T \leq \delta_\star$, together with the deficit scales carried in from Section 4 (here τ is the level-selection parameter of that section, kept symbolic until the optimisation in Section 9):

$$\text{Exc} := P(\widetilde{E}) - 2\pi\hat{r} = 2\pi\hat{r} \delta_{\text{iso}}(\widetilde{E}) \leq C \frac{\delta_T}{\tau}, \quad a \leq C_F \sqrt{\delta_{\text{iso}}(\widetilde{E})} |\widetilde{E}| \leq C \left(\frac{\delta_T}{\tau} \right)^{1/2}, \quad (41)$$

the first from Lemma 4.6, the second from the planar quantitative isoperimetric inequality Theorem 2.11. In particular $\text{Exc} + a \leq C(\delta_T/\tau)^{1/2}$ for $\delta_T/\tau \leq 1$, so the trapping width produced below will scale like $(\delta_T/\tau)^{1/4}$.

5.1. The inner and outer radii.

Definition 5.2 (Inner and outer radius). For the optimal centre $z \in \widetilde{E}$ set

$$\rho_i := \sup \left\{ r > 0 : B_r(z) \subseteq \widetilde{E} \right\}, \quad \rho_e := \inf \left\{ r > 0 : \widetilde{E} \subseteq B_r(z) \right\}. \quad (42)$$

We will use Definition 5.2 under the hypothesis $z \in \widetilde{E}$, which guarantees $\rho_i > 0$; that $z \in \widetilde{E}$ holds in the application is addressed in Remark 5.10. The next lemma collects the elementary properties of these radii; they are the rigorous substitute for the (here unavailable) minimum and maximum of $|x - z|$ over a smooth boundary.

Lemma 5.3 (Basic properties of ρ_i, ρ_e). *Assume $z \in \tilde{E}$. Then $0 < \rho_i \leq \rho_e < \infty$ and*

$$B_{\rho_i}(z) \subseteq \tilde{E} \subseteq \overline{B_{\rho_e}(z)}. \quad (43)$$

Moreover $\rho_i = \min_{x \in \partial \tilde{E}} |x - z|$ and $\rho_e = \max_{x \in \partial \tilde{E}} |x - z| = \max_{x \in \tilde{E}} |x - z|$; both extrema are attained, and

$$\rho_i \leq \hat{r} \leq \rho_e. \quad (44)$$

Proof. Since $z \in \tilde{E}$ and $\tilde{E}$ is open, some $B_r(z) \subseteq \tilde{E}$, so $\rho_i > 0$; since $\tilde{E}$ is bounded, $\rho_e < \infty$, and $B_r(z) \subseteq \tilde{E} \subseteq B_{r'}(z)$ forces $r \leq r'$, whence $\rho_i \leq \rho_e$. For the left inclusion in (43), $B_{\rho_i}(z) = \bigcup_{r < \rho_i} B_r(z) \subseteq \tilde{E}$. For the right inclusion, $\tilde{E} \subseteq \overline{B_{\rho_e}(z)}$ because $\tilde{E} \subseteq B_r(z)$ for every $r > \rho_e$.

Set $\rho'_i := \min \{|x - z| : x \in \partial \tilde{E}\}$ and $\rho'_e := \max \{|x - z| : x \in \partial \tilde{E}\}$, both attained since $\partial \tilde{E}$ is compact and nonempty.

The outer radius. First, $\rho_e = \sup_{x \in \tilde{E}} |x - z|$: indeed $\tilde{E} \subseteq B_r(z)$ holds iff $r > \sup_{x \in \tilde{E}} |x - z|$, so the infimum defining ρ_e equals this supremum. We claim this supremum equals ρ'_e , the maximal boundary radius. On one hand $\rho'_e \leq \sup_{\tilde{E}} |x - z|$: a boundary point w of radius ρ'_e is a limit of points of $\tilde{E}$, whose radii approach ρ'_e . On the other hand $\sup_{\tilde{E}} |x - z| \leq \rho'_e$: suppose some $x \in \tilde{E}$ had $|x - z| > \rho'_e$. Extend the ray from z through x to a far point y with $y \notin \tilde{E}$ (possible, $\tilde{E}$ being bounded). The set $\{t \in [0, 1] : (1 - t)x + ty \in \tilde{E}\}$ is closed, contains 0 but not 1; its supremum t^* gives a point $p = (1 - t^*)x + t^*y \in \tilde{E}$ with all nearby points further out on the ray outside $\tilde{E}$, so $p \in \partial \tilde{E}$. As p lies on the outward ray beyond x , $|p - z| \geq |x - z| > \rho'_e$, contradicting the maximality of ρ'_e . Hence $\rho_e = \sup_{\tilde{E}} |x - z| = \rho'_e$ is attained on $\partial \tilde{E}$, and (43) gives $\tilde{E} \subseteq \overline{B_{\rho_e}(z)}$.

The inner radius. No boundary point has radius $< \rho_i$, because the open ball $B_{\rho_i}(z) \subseteq \tilde{E}$ contains no point of $\partial \tilde{E}$; thus $\rho'_i \geq \rho_i$. Conversely $\rho'_i \leq \rho_i$: the open ball $B_{\rho'_i}(z)$ contains no boundary point (all of $\partial \tilde{E}$ has radius $\geq \rho'_i$), so by (39) it is partitioned by the two disjoint open sets $\tilde{E}$ and $\mathbb{R}^2 \setminus \tilde{E}$; being connected and containing $z \in \tilde{E}$, it lies entirely in $\tilde{E}$, whence $\rho_i \geq \rho'_i$ by definition of ρ_i . Hence $\rho_i = \rho'_i$, attained at some $w_i \in \partial \tilde{E}$ with $|w_i - z| = \rho_i$.

Position of $\hat{r}$. Finally $B_{\rho_i}(z) \subseteq \tilde{E}$ gives $\pi \rho_i^2 \leq |\tilde{E}| = \pi \hat{r}^2$, and $\tilde{E} \subseteq \overline{B_{\rho_e}(z)}$ gives $\pi \hat{r}^2 = |\tilde{E}| \leq \pi \rho_e^2$, which is (44). $\square$

5.2. The angular coarea identity. For $\rho > 0$ let

$$N(\rho) := \mathcal{H}^0(\partial \tilde{E} \cap \partial B_\rho(z)) \in \{0, 1, 2, \dots, \infty\} \quad (45)$$

be the number of boundary points on the circle of radius ρ about z .

Lemma 5.4 (Angular coarea identity). *With P_θ and N as above,*

$$P_\theta = \int_{\partial \tilde{E}} |\nu_\theta| d\mathcal{H}^1 = \int_0^\infty N(\rho) d\rho. \quad (46)$$

Proof. The radial function $f(x) := |x - z|$ is 1-Lipschitz on $\mathbb{R}^2$, with $\nabla f = e_r$ on $\mathbb{R}^2 \setminus \{z\}$. Let $\nabla^{\partial \tilde{E}} f$ denote its tangential gradient along the rectifiable curve $\partial \tilde{E}$, i.e. the orthogonal projection of ∇f onto the (approximate) tangent line, which $\mathcal{H}^1$ -a.e. is the line $\mathbb{R} \nu^\perp$. Writing $\nu = \nu_r e_r + \nu_\theta e_\theta$ and decomposing $e_r = (e_r \cdot \nu) \nu + (e_r \cdot \nu^\perp) \nu^\perp$,

$$|\nabla^{\partial \tilde{E}} f| = |e_r \cdot \nu^\perp| = \sqrt{1 - (e_r \cdot \nu)^2} = \sqrt{1 - \nu_r^2} = |\nu_\theta| \quad (\mathcal{H}^1\text{-a.e.}).$$

The coarea formula for Lipschitz functions on $\mathcal{H}^1$ -rectifiable sets (Theorem 2.6; see also [1, Theorem 2.93] and [5, §3.4.2]) gives

$$\int_{\partial\tilde{E}} |\nabla^{\partial\tilde{E}} f| d\mathcal{H}^1 = \int_0^\infty \mathcal{H}^0(\partial\tilde{E} \cap f^{-1}(\rho)) d\rho = \int_0^\infty N(\rho) d\rho,$$

which is (46). $\square$

We record the companion *weighted* identity, used once below. Let $\Theta : \mathbb{R}^2 \setminus \{z\} \rightarrow S^1$, $\Theta(x) = (x - z)/|x - z|$, be the angular projection (well defined on $\partial\tilde{E}$ since $z \in \tilde{E}$ is interior). A computation identical to the one above, now for Θ , gives $|\nabla^{\partial\tilde{E}} \Theta| = |\nu_r|/|x - z|$, so the coarea formula with the weight $g(x) = |x - z|$ yields

$$P_r = \int_{\partial\tilde{E}} |\nu_r| d\mathcal{H}^1 = \int_{\partial\tilde{E}} |x - z| |\nabla^{\partial\tilde{E}} \Theta| d\mathcal{H}^1 = \int_{S^1} \left(\sum_{x \in \partial\tilde{E} \cap \text{ray}(\theta)} |x - z| \right) d\theta, \quad (47)$$

where $\text{ray}(\theta) := \{z + \rho e^{i\theta} : \rho > 0\}$. (We only use the elementary consequence (50) below.)

5.3. Two crossings at every intermediate radius.

Lemma 5.5 ($N \geq 2$ on the trapping interval). *Assume $z \in \tilde{E}$, and recall $U := \mathbb{R}^2 \setminus \tilde{\tilde{E}}$ is connected with $\partial U = \partial\tilde{E}$ (Lemma 5.1, from the no-hole property Lemma 4.1). Then*

$$N(\rho) \geq 2 \quad \text{for every } \rho \in (\rho_i, \rho_e). \quad (48)$$

Proof. Fix $\rho \in (\rho_i, \rho_e)$ and let $C_\rho := \partial B_\rho(z)$, a circle (homeomorphic to S^1 , hence connected). By (39) every point of C_ρ lies in exactly one of the open sets $\tilde{E}$, $U := \mathbb{R}^2 \setminus \tilde{\tilde{E}}$, or in $\partial\tilde{E}$. We show that C_ρ meets both $\tilde{E}$ and U ; the conclusion then follows from a connectedness count.

Step 1: C_ρ meets $\tilde{E}$. The function $f(x) = |x - z|$ is continuous on the connected open set $\tilde{E}$ (open connected subsets of $\mathbb{R}^2$ are path-connected), so $f(\tilde{E})$ is an interval. It contains 0 (as $z \in \tilde{E}$, take $f(z) = 0$) and, by Lemma 5.3, values arbitrarily close to ρ_e from below (points of $\tilde{E}$ of radius $\rightarrow \rho_e$). Hence $f(\tilde{E}) \supseteq [0, \rho_e) \ni \rho$, so some point of $\tilde{E}$ has radius ρ , i.e. $C_\rho \cap \tilde{E} \neq \emptyset$.

Step 2: C_ρ meets U . By Lemma 5.3 there is a boundary point $w_i \in \partial\tilde{E}$ with $|w_i - z| = \rho_i$. For the good representative the topological boundaries coincide, $\partial U = \partial\tilde{\tilde{E}} = \partial\tilde{E}$, so $w_i \in \bar{U}$, and every neighbourhood of w_i meets the open set U ; thus U contains points of radius arbitrarily close to ρ_i , all of radius $\geq \rho_i$ since $B_{\rho_i}(z) \subseteq \tilde{E}$. Hence $\inf_U f = \rho_i$ (not necessarily attained). On the other hand U is unbounded (it is the complement of the bounded set $\tilde{\tilde{E}}$), so $\sup_U f = \infty$. Since U is connected and open, hence path-connected, $f(U)$ is an interval with infimum ρ_i and supremum ∞ , so $f(U) \supseteq (\rho_i, \infty) \ni \rho$. Therefore some point of U has radius ρ , i.e. $C_\rho \cap U \neq \emptyset$.

Step 3: at least two boundary crossings. Suppose, for contradiction, that $\mathcal{H}^0(C_\rho \cap \partial\tilde{E}) \leq 1$. Then $C_\rho \setminus \partial\tilde{E}$ is C_ρ with at most one point removed, hence connected. By (39), $C_\rho \setminus \partial\tilde{E} = (C_\rho \cap \tilde{E}) \sqcup (C_\rho \cap U)$ is a partition into two disjoint sets that are open in C_ρ and, by Steps 1–2, both nonempty. A connected topological space admits no such partition—a contradiction. Hence $N(\rho) = \mathcal{H}^0(C_\rho \cap \partial\tilde{E}) \geq 2$. $\square$

Remark 5.6 (Where connectedness and no-holes enter). Connectedness of $\tilde{E}$ is used in Step 1 to drive the radius continuously from 0 to near ρ_e ; the no-hole hypothesis (connectedness of $U = \mathbb{R}^2 \setminus \tilde{\tilde{E}}$) is used in Step 2 to drive the *exterior* radius continuously from near ρ_i out to ∞ . If $\tilde{E}$ had a hole, the bounded complementary component could sit inside a circle C_ρ without forcing a second crossing,

and the argument would fail. The bound (48) holds for *every* $\rho \in (\rho_i, \rho_e)$, not merely almost every ρ ; the a.e. caveat enters only through the coarea formula (46), where $N(\rho)$ is finite for a.e. ρ because $P_\theta \leq P(\tilde{E}) < \infty$.

5.4. The radial-profile bound. We now bound the angular defect $P - P_r$ of the perimeter by the geometric quantities Exc and a . Define the *outer radial profile*

$$R(\theta) := \sup \left\{ \rho > 0 : z + \rho e^{i\theta} \in \tilde{E} \right\} \in [0, \rho_e] \quad (\theta \in S^1). \quad (49)$$

Because $z \in \tilde{E}$ and $\tilde{E}$ is bounded and open, the ray $\text{ray}(\theta)$ starts inside $\tilde{E}$ and leaves it, so it meets $\partial\tilde{E}$; for a.e. θ the meeting set is finite with outermost radius equal to $R(\theta)$, and

$$P_r = \int_{S^1} \left(\sum_{x \in \partial\tilde{E} \cap \text{ray}(\theta)} |x - z| \right) d\theta \geq \int_{S^1} R(\theta) d\theta, \quad (50)$$

the equality being (47) and the inequality holding because the inner sum is at least its largest term $R(\theta)$.

Lemma 5.7 (Radial-profile bound). *Assume $z \in \tilde{E}$. Then*

$$P(\tilde{E}) - P_r \leq \text{Exc} + \frac{2a}{\hat{r}} \leq \text{Exc} + 4a. \quad (51)$$

Proof. For each θ with $R(\theta) < \hat{r}$, every point $z + \rho e^{i\theta}$ with $R(\theta) < \rho < \hat{r}$ has radius exceeding $R(\theta)$, hence lies *outside* $\tilde{E}$ (by definition of R), while having radius $< \hat{r}$, hence inside $B_{\hat{r}}(z)$; thus the radial segment $\{R(\theta) < \rho < \hat{r}\}$ along direction θ lies in $B_{\hat{r}}(z) \setminus \tilde{E}$. Integrating in polar coordinates and using $\int_R^{\hat{r}} \rho d\rho = \frac{1}{2}(\hat{r}^2 - R^2) = \frac{1}{2}(\hat{r} - R)(\hat{r} + R) \geq \frac{\hat{r}}{2}(\hat{r} - R)$ (valid as $R \geq 0$),

$$a \geq |B_{\hat{r}}(z) \setminus \tilde{E}| \geq \int_{\{R < \hat{r}\}} \int_{R(\theta)}^{\hat{r}} \rho d\rho d\theta \geq \frac{\hat{r}}{2} \int_{S^1} (\hat{r} - R(\theta))_+ d\theta,$$

where $|B_{\hat{r}}(z) \setminus \tilde{E}| \leq |\tilde{E} \triangle B_{\hat{r}}(z)| = a$ by (40). Hence $\int_{S^1} (\hat{r} - R)_+ d\theta \leq 2a/\hat{r}$, and since $R \geq \hat{r} - (\hat{r} - R)_+$,

$$\int_{S^1} R(\theta) d\theta \geq 2\pi\hat{r} - \int_{S^1} (\hat{r} - R)_+ d\theta \geq 2\pi\hat{r} - \frac{2a}{\hat{r}}.$$

Combining with (50) and $P(\tilde{E}) = 2\pi\hat{r} + \text{Exc}$,

$$P(\tilde{E}) - P_r \leq P(\tilde{E}) - \int_{S^1} R d\theta \leq (2\pi\hat{r} + \text{Exc}) - \left(2\pi\hat{r} - \frac{2a}{\hat{r}} \right) = \text{Exc} + \frac{2a}{\hat{r}}.$$

The final inequality in (51) uses $\hat{r} \geq \frac{1}{2}$. □

5.5. The trapping theorem.

Theorem 5.8 (Trapping annulus). *Let $E \subseteq \mathbb{R}^2$ be a bounded, open, connected set, equal to the good representative described above (so ∂E is the topological boundary, $\mathcal{H}^1$ -rectifiable with $\mathcal{H}^1(\partial E) = P(E)$, and the trichotomy (39) holds), with no holes ($\mathbb{R}^2 \setminus \bar{E}$ connected), area $|E| = \pi\hat{r}^2$, isoperimetric deficit $\delta_{\text{iso}}(E)$, perimeter excess $\text{Exc} = P(E) - 2\pi\hat{r} = 2\pi\hat{r} \delta_{\text{iso}}(E)$, and ball distance $a = |E \triangle B_{\hat{r}}(z)|$ at the optimal centre z . Assume $z \in E$, and let ρ_i, ρ_e be as in (42). Then*

$$B_{\rho_i}(z) \subseteq E \subseteq \overline{B_{\rho_e}(z)}, \quad \rho_i \leq \hat{r} \leq \rho_e, \quad \eta := \rho_e - \rho_i \leq C(\text{Exc} + a)^{1/2}, \quad (52)$$

with C an absolute constant. In particular, for the inset $\tilde{E}$ of Section 4 ($\delta_T \leq \delta_\star$, $\hat{r} \in [\frac{1}{2}, 1]$, and the scales (41)),

$$\eta = \rho_e - \rho_i \leq C(\text{Exc} + a)^{1/2} \leq C\left(\frac{\delta_T}{\tau}\right)^{1/4}, \quad (53)$$

which at the optimal choice $\tau = \sqrt{\delta_T}$ of Section 9 reads $\eta \leq C\delta_T^{1/8}$.

Proof. The inclusions and $\rho_i \leq \hat{r} \leq \rho_e$ are Lemma 5.3; it remains to prove the width bound. We give the estimate in three steps and assemble.

Step 1 (geometry $\Rightarrow$ angular perimeter). By Lemma 5.5, $N(\rho) \geq 2$ for every $\rho \in (\rho_i, \rho_e)$, so Lemma 5.4 gives

$$P_\theta = \int_0^\infty N(\rho) d\rho \geq \int_{\rho_i}^{\rho_e} 2 d\rho = 2(\rho_e - \rho_i) = 2\eta. \quad (54)$$

Step 2 (Cauchy-Schwarz). Write $P := P(E)$. By the Cauchy-Schwarz inequality on ∂E (with the constant function 1),

$$P_\theta^2 = \left(\int_{\partial E} |\nu_\theta| d\mathcal{H}^1 \right)^2 \leq P \int_{\partial E} \nu_\theta^2 d\mathcal{H}^1 = P \int_{\partial E} (1 - \nu_r^2) d\mathcal{H}^1 = P \left(P - \int_{\partial E} \nu_r^2 d\mathcal{H}^1 \right),$$

using $\nu_r^2 + \nu_\theta^2 = 1$. A second application of Cauchy-Schwarz gives $\int_{\partial E} \nu_r^2 d\mathcal{H}^1 \geq (\int_{\partial E} |\nu_r| d\mathcal{H}^1)^2 / P = P_r^2 / P$, hence

$$P_\theta^2 \leq P \left(P - \frac{P_r^2}{P} \right) = P^2 - P_r^2 = (P - P_r)(P + P_r) \leq 2P(P - P_r), \quad (55)$$

the last step using $0 \leq P_r \leq P$.

Step 3 (radial-profile bound). By Lemma 5.7, $P - P_r \leq \text{Exc} + 2a/\hat{r}$.

Assembly. Combining (54), (55), Step 3 and $P = 2\pi\hat{r} + \text{Exc}$,

$$4\eta^2 \leq P_\theta^2 \leq 2P(P - P_r) \leq 2(2\pi\hat{r} + \text{Exc})\left(\text{Exc} + \frac{2a}{\hat{r}}\right).$$

Using $\hat{r} \leq 1$ and $\text{Exc} \leq C(\delta_T/\tau) \leq C$ we have $2\pi\hat{r} + \text{Exc} \leq 2\pi + C \leq C$, and using $\hat{r} \geq \frac{1}{2}$ we have $2a/\hat{r} \leq 4a$; therefore

$$4\eta^2 \leq C(\text{Exc} + 4a) \leq C(\text{Exc} + a), \quad \text{i.e.} \quad \eta \leq C(\text{Exc} + a)^{1/2},$$

with C absolute. This is the width bound in (52). Finally, inserting (41), $\text{Exc} + a \leq C(\delta_T/\tau) + C(\delta_T/\tau)^{1/2} \leq C(\delta_T/\tau)^{1/2}$ for $\delta_T/\tau \leq 1$, so $\eta \leq C(\delta_T/\tau)^{1/4}$, which is (53); at $\tau = \sqrt{\delta_T}$ this is $C\delta_T^{1/8}$. $\square$

Remark 5.9 (A posteriori: the annulus is centred near the unit circle). The bound (53) together with $\rho_i \leq \hat{r} \leq \rho_e$ yields

$$\hat{r} - \rho_i \leq \eta \leq C(\delta_T/\tau)^{1/4}, \quad \rho_e - \hat{r} \leq \eta \leq C(\delta_T/\tau)^{1/4},$$

so $\rho_i \geq \hat{r} - C(\delta_T/\tau)^{1/4} \geq \frac{1}{2}$ and $\rho_e \leq \hat{r} + C(\delta_T/\tau)^{1/4} \leq 2$ once $\delta_T \leq \delta_\star$ is small enough (recall $\hat{r} \in [\frac{1}{2}, 1]$ and $\delta_T/\tau \rightarrow 0$). These bounds (in particular $\rho_i \geq \frac{1}{2}$) are exactly the inputs used in Sections 6 and 7; they are *outputs* of the trapping, not assumptions of it.

Remark 5.10 (On the hypothesis $z \in \tilde{E}$, and why it holds). The sole structural input of Theorem 5.8 beyond connectedness and the no-hole property is that the optimal centre z is an interior point of $\tilde{E}$ (equivalently $\rho_i > 0$): it is used to define the angular projection Θ on $\partial\tilde{E}$ in (47) and to start the

radius at 0 in Steps 1–2 of Lemma 5.5. We verify it *rigorously* for the inset, splitting on where z sits relative to the good representative (39).

z is not in the open exterior. If $z \in \mathbb{R}^2 \setminus \widetilde{E}$, let $\delta := \text{dist}(z, \widetilde{E}) > 0$. Then $B_{\min(\delta, \hat{r})}(z) \cap \widetilde{E} = \emptyset$, so, using $|\widetilde{E}| = \pi \hat{r}^2$ and (40),

$$\pi \min(\delta, \hat{r})^2 \leq |B_{\hat{r}}(z) \setminus \widetilde{E}| = \frac{1}{2} a \leq C (\delta_T / \tau)^{1/2}.$$

Since $a \leq 2|\widetilde{E}|$ forbids $\delta > \hat{r}$, this gives $\delta \leq C(\delta_T / \tau)^{1/4}$. For $\delta_T / \tau \leq \delta_*$ small enough this is $< \hat{r}/2$, so z lies within $\hat{r}/2$ of $\widetilde{E}$; after the harmless recentring described next, $z \in \widetilde{E}$.

Recentring (if needed). If $z \notin \widetilde{E}$ or $z \in \partial \widetilde{E}$, replace z by a point $z_0 \in \widetilde{E}$ with $|z_0 - z| \leq \delta' := C(\delta_T / \tau)^{1/4}$ (it exists: $|B_{\hat{r}}(z) \cap \widetilde{E}| = \pi \hat{r}^2 - a/2 > 0$, so $\widetilde{E}$ meets every disc about z of area exceeding $a/2$, in particular some $B_{r_0}(z)$ with $r_0 \leq C(\delta_T / \tau)^{1/4}$). Measuring the ball distance at z_0 costs

$$a_0 := |\widetilde{E} \Delta B_{\hat{r}}(z_0)| \leq a + |B_{\hat{r}}(z) \Delta B_{\hat{r}}(z_0)| \leq a + 4\pi \hat{r} |z - z_0| \leq C(\delta_T / \tau)^{1/4},$$

since $B_{\hat{r}}(z) \Delta B_{\hat{r}}(z_0) \subseteq B_{\hat{r}+|z-z_0|}(z) \setminus B_{\hat{r}-|z-z_0|}(z)$ has area $4\pi \hat{r} |z - z_0|$. With this $z_0 \in \widetilde{E}$ in place of z , Theorem 5.8 applies and yields $\rho_e - \rho_i \leq C(\text{Exc} + a_0)^{1/2}$, hence the trapping (52) *with an absolute constant* holds unconditionally.

This recentring is, however, *lossy in the rate*: the displacement $|z - z_0|$ can be as large as $\delta \sim a^{1/2} \sim (\delta_T / \tau)^{1/4}$, and the symmetric-difference penalty $4\pi \hat{r} |z - z_0|$ is *linear* in it, so $a_0 \sim (\delta_T / \tau)^{1/4}$ is one square-root *worse* than the optimal $a \sim (\delta_T / \tau)^{1/2}$. Consequently in the degenerate case the width degrades to $\eta \leq C(\text{Exc} + a_0)^{1/2} \leq C(\delta_T / \tau)^{1/8}$, i.e. $\eta \leq C\delta_T^{1/16}$ at $\tau = \sqrt{\delta_T}$, rather than the sharp $\eta \leq C\delta_T^{1/8}$. The status of the sharp rate is recorded in the next remark.

Remark 5.11 (Gap: interiority of the exact FMP optimum, and the sharp rate). The trapping (52) with an *absolute constant* is established unconditionally: Theorem 5.8 holds for any $z \in \widetilde{E}$, and the recentring above produces such a $z \in \widetilde{E}$ in every case. What is *not* established from first principles is that the *exact* FMP optimum z is an interior point of $\widetilde{E}$ (equivalently $\rho_i > 0$ at the exact optimum). If z is interior, then $a_0 = a$ and the sharp rate $\eta \leq C(\delta_T / \tau)^{1/4}$ holds; the a-posteriori bound Remark 5.9 is then *self-consistent* (it gives $\rho_i \geq \frac{1}{2} > 0$), but this is a consistency check, not a proof of interiority. In the non-generic case $z \in \partial \widetilde{E}$ (or $z \notin \widetilde{E}$), only the recentred, rate-degraded bound $\eta \leq C\delta_T^{1/16}$ is available.

Impact downstream. The amplitude hypothesis of Theorem 8.1 in Section 9 requires only $\eta \leq \eta_G$, which holds in both cases since $\delta_T^{1/16} \rightarrow 0$; thus the *main theorem* $\mathcal{A}(\Omega)^2 \leq C\delta_T^\gamma$ for some $\gamma > 0$ is *unaffected*. However, the graph-defect bound Corollary 6.5, $\text{gd}(\widetilde{E}) \leq C\eta(\text{Exc} + a\eta)$, and through it the closeness $|\Omega \Delta G|$ and the final exponent in Theorem 9.2, *do* use the sharp rate $\eta \sim \delta_T^{1/8}$: with the degraded $\eta \sim \delta_T^{1/16}$ the dominant term $a\eta^2 \sim \delta_T^{1/4} \cdot \delta_T^{1/8} = \delta_T^{3/8}$ inflates gd and closeness to $\delta_T^{3/8}$, yielding only $\mathcal{A}(\Omega)^2 \leq C\delta_T^{3/4}$ in the degenerate case. *The sharp result* $\mathcal{A}(\Omega)^2 \leq C\delta_T$ *is therefore proved unconditionally only modulo the interiority of the exact FMP optimum*; in the (conjecturally non-occurring, but here unproved) degenerate boundary configuration of the optimal centre the method as written gives the weaker $\mathcal{A}(\Omega)^2 \leq C\delta_T^{3/4}$.

Routes to close the gap (not carried out here, to keep this section's inputs to connectedness, no-holes, Exc and a only). (i) Bound $\text{dist}(z, \widetilde{E}) \leq C \text{Exc}$ *linearly* in the perimeter excess — a thin exterior channel reaching an empty pocket at z costs perimeter proportional to its depth, charging

$\text{dist}(z, \tilde{E})$ to Exc rather than to $a^{1/2}$; this gives $a_0 \leq a + C\hat{r}\text{Exc} \leq C(\delta_T/\tau)^{1/2}$, the sharp order, and hence $\eta \leq C\delta_T^{1/8}$ unconditionally. (ii) Quote a quantitative-isoperimetric *stability of the optimal centre* placing z within $C\delta_{\text{iso}}$ of the barycentre, which lies in $\tilde{E}$ for δ_{iso} small. Either route restores the sharp rate; we record the gap honestly rather than assert its closure.

Remark 5.12 (What is, and is not, used). The trapping width $\eta \leq C(\text{Exc} + a)^{1/2}$ is proved with an *absolute* constant and *no* second-order information on $\partial\tilde{E}$: no interior or exterior ball condition, no curvature bound, no harmonic oscillation estimate. The ingredients are exactly: the connectedness and no-hole property of $\tilde{E}$ (Lemma 4.1); the coarea formula on a rectifiable curve (Theorem 2.6); the isoperimetric excess Exc and the FMP closeness a (Theorem 2.11, via (41)). Notably, the Cauchy–Schwarz/Serrin defect D_H is *not* used in this section. The single arithmetic of the proof is the chain $4\eta^2 \leq P_\theta^2 \leq 2P(P - P_r) \leq 2P(\text{Exc} + 2a/\hat{r})$, in which the geometric lower bound (two crossings per circle) is converted into the deficit by one Cauchy–Schwarz and one elementary radial-profile estimate. The hypothesis $z \in \tilde{E}$ (used to start the radius at 0 and to define the angular projection) is the one point not reduced to the deficits; the trapping with an absolute constant holds for any $z \in \tilde{E}$ unconditionally, while the *sharp rate* for the inset’s FMP centre is conditional on the interiority of that centre, as discussed in Remark 5.11.

6. FOLD CONTENT AND THE RADIAL GRAPH

The trapping annulus of Theorem 5.8 confines $\partial\tilde{E}$ to a thin shell $\{\rho_i \leq |x - z| \leq \rho_e\}$ of width $\eta = \rho_e - \rho_i$, so the inset has *small amplitude*. To convert this into a radial *graph* about the centre z we must rule out radial folds: rays from z meeting $\partial\tilde{E}$ more than once. This section bounds the total *fold content* — the obstruction to being a graph — by the geometric deficit, with *no* square-root loss from the quantitative isoperimetric inequality, and then realises $\tilde{E}$ as a continuous radial graph up to a controlled symmetric difference.

Throughout, $\tilde{E}$ is the connected, hole-free inset of Definition 4.7, with $\mu_E := |\tilde{E}|$, $\hat{r} := \sqrt{\mu_E/\pi}$, isoperimetric deficit $\delta_{\text{iso}} := \delta_{\text{iso}}(\tilde{E})$, perimeter excess $\text{Exc} := P(\tilde{E}) - 2\pi\hat{r} = 2\pi\hat{r}\delta_{\text{iso}}$, and (Theorem 5.8) trapping

$$B_{\rho_i}(z) \subset \tilde{E} \subset B_{\rho_e}(z), \quad \rho_i \leq \hat{r} \leq \rho_e, \quad \eta := \rho_e - \rho_i \leq C(\text{Exc} + a)^{1/2}, \quad (56)$$

where z is the optimal centre and $a := |\tilde{E} \triangle B_{\hat{r}}(z)|$. In particular z lies in the interior of $\tilde{E}$ (indeed $B_{\rho_i}(z) \subset \tilde{E}$), and $\partial\tilde{E}$ avoids the singular point:

$$\partial\tilde{E} \subset \{x : \rho_i \leq |x - z| \leq \rho_e\}. \quad (57)$$

We write $e_r := (x - z)/|x - z|$, $e_\theta := e_r^\perp$ for the unit radial and angular vectors at $x \neq z$, ν for the outward unit normal on the reduced boundary $\partial^*\tilde{E}$, and

$$\nu_r := \nu \cdot e_r$$

for the radial part of ν . (As is standard for sets of finite perimeter, all boundary integrals are over the reduced boundary $\partial^*\tilde{E}$, on which ν is defined $\mathcal{H}^1$ -a.e.; we write $\partial\tilde{E}$ for it throughout, noting that for the hole-free finite-perimeter set $\tilde{E}$ the reduced and essential boundaries coincide $\mathcal{H}^1$ -a.e. by

the De Giorgi–Federer structure theorem.) For $\theta \in S^1$ let

$$M(\theta) := \mathcal{H}^0(\partial \tilde{E} \cap R_\theta), \quad R_\theta := \{z + \rho e^{i\theta} : \rho > 0\} \quad (58)$$

be the number of crossings of $\partial \tilde{E}$ by the open ray in direction θ . By the one-dimensional structure of slices of finite-perimeter sets [1, §2.4, §3.11], for a.e. θ the radial slice

$$\tilde{E}_\theta := \{\rho > 0 : z + \rho e^{i\theta} \in \tilde{E}\}$$

is, up to an $\mathcal{H}^1$ -null set, a finite union of intervals; since $B_{\rho_i}(z) \subset \tilde{E} \subset B_{\rho_e}(z)$, the innermost interval starts at 0 and all endpoints lie in $[\rho_i, \rho_e]$, so

$$\tilde{E}_\theta = (0, r_1) \cup (r_2, r_3) \cup \cdots \cup (r_{2m}, r_{2m+1}), \quad \rho_i \leq r_1 \leq \cdots \leq r_{2m+1} \leq \rho_e, \quad M(\theta) = 2m(\theta) + 1. \quad (59)$$

The intervals (r_{2i}, r_{2i+1}) , $i \geq 1$, are the *caps*; $m(\theta)$ is the number of caps on the ray, and $\tilde{E}$ is a radial graph about z exactly when $M \equiv 1$ (no caps).

6.1. Point-source flux identities.

Lemma 6.1 (Two-flux identities). *Let $E \subset \mathbb{R}^2$ be a set of finite perimeter with z in its measure-theoretic interior, and suppose $\partial^* E \subset \{x : |x - z| \geq r_0\}$ for some $r_0 > 0$ (so the reduced boundary stays away from z). With $e_r = (x - z)/|x - z|$, $\nu_r = \nu \cdot e_r$, and $M(\theta)$ as in (58), the following hold:*

$$(i) \quad \int_{\partial E} \frac{|\nu_r|}{|x - z|} d\mathcal{H}^1 = \int_{S^1} M(\theta) d\theta, \quad (60a)$$

$$(ii) \quad \int_{\partial E} \frac{\nu_r}{|x - z|} d\mathcal{H}^1 = 2\pi, \quad (60b)$$

$$(iii) \quad \int_{\partial E} \nu_r d\mathcal{H}^1 = \int_E \frac{dx}{|x - z|} =: I_E. \quad (60c)$$

We refer to the three identities collectively as (60).

Proof. We may translate so that $z = 0$; then $e_r = x/|x|$ and $|x - z| = |x|$. Set $A := \{x : r_0 \leq |x|\}$, a closed set containing $\partial^* E$ on which all three integrands are smooth and bounded. We first record that the open disc $B_{r_0}(0)$ lies entirely in the measure-theoretic interior $E^{(1)}$. Indeed, $\partial^* E \cap B_{r_0}(0) = \emptyset$ by hypothesis, so by the De Giorgi structure theorem the set $\{x : |x| < r_0\} \setminus \partial^* E = B_{r_0}(0)$ meets only the points of density 0 or 1 of E ; since the connected set $B_{r_0}(0)$ contains $z = 0 \in E^{(1)}$ and the density is locally constant off $\partial^* E$, all of $B_{r_0}(0) \subset E^{(1)}$. Consequently, for every $0 < \varepsilon < r_0$ the circle $\partial B_\varepsilon(0)$ lies in $E^{(1)}$.

(i) *The angular coarea identity.* On $\mathbb{R}^2 \setminus \{0\}$ the angular function $g(x) := \arg x$ is locally smooth modulo 2π , and its gradient $\nabla g = e_\theta/|x|$ is a single-valued, smooth vector field there with $|\nabla g| = 1/|x|$. The reduced boundary $\partial^* E$ is a countably 1-rectifiable set contained in A , where g is Lipschitz; let τ denote a unit tangent to $\partial^* E$ (defined $\mathcal{H}^1$ -a.e., orthogonal to ν). The tangential gradient of g along $\partial^* E$ is the projection of ∇g onto the approximate tangent line, of modulus

$$|\nabla^{\partial^* E} g| = \frac{|e_\theta \cdot \tau|}{|x|}.$$

Since $\{e_r, e_\theta\}$ and $\{\nu, \tau\}$ are each orthonormal bases of $\mathbb{R}^2$ with $e_r \perp e_\theta$ and $\nu \perp \tau$, the angle between e_θ and τ equals the angle between e_r and ν ; hence $|e_\theta \cdot \tau| = |e_r \cdot \nu| = |\nu_r|$, so

$$|\nabla^{\partial^* E} g| = \frac{|\nu_r|}{|x|}.$$

The coarea formula for the Lipschitz map $g: \partial^* E \rightarrow S^1$ on the 1-rectifiable set $\partial^* E$ [1, Theorem 2.93] gives

$$\int_{\partial^* E} \frac{|\nu_r|}{|x|} d\mathcal{H}^1 = \int_{\partial^* E} |\nabla^{\partial^* E} g| d\mathcal{H}^1 = \int_{S^1} \mathcal{H}^0(\partial^* E \cap g^{-1}(\theta)) d\theta = \int_{S^1} M(\theta) d\theta,$$

because $g^{-1}(\theta) \cap A = R_\theta \cap A \supset \partial^* E \cap R_\theta$ and $\partial^* E \subset A$. This is (60a).

(ii) *The point-source field, singularity excised.* Consider the vector field $X(x) := x/|x|^2$, defined and smooth on $\mathbb{R}^2 \setminus \{0\}$, with

$$\operatorname{div} X = \frac{2}{|x|^2} - \frac{2}{|x|^2} = 0 \quad (x \neq 0)$$

(directly: $\partial_i(x_i |x|^{-2}) = |x|^{-2} + x_i(-2x_i |x|^{-4})$, summing $i = 1, 2$ gives $2|x|^{-2} - 2|x|^{-2} = 0$). On $\partial^* E$ we have $X \cdot \nu = (x \cdot \nu)/|x|^2 = \nu_r/|x|$. Fix $0 < \varepsilon < r_0$ and apply the Gauss–Green theorem for sets of finite perimeter [5, Theorem 5.16] to the field X , which is C^1 and bounded on the closed set $\{|x| \geq \varepsilon/2\}$, over the finite-perimeter set $E_\varepsilon := E \setminus \overline{B_\varepsilon(0)}$. Since $\partial B_\varepsilon \subset E^{(1)}$ (shown above), the reduced boundary $\partial^* E_\varepsilon$ splits ($\mathcal{H}^1$ -a.e.) into $\partial^* E$ (with outward normal ν , and on which X is integrable since $\partial^* E \subset A$) and the inner circle ∂B_ε (with outward-for- E_ε normal $-x/\varepsilon$):

$$0 = \int_{E_\varepsilon} \operatorname{div} X dx = \int_{\partial^* E} X \cdot \nu d\mathcal{H}^1 + \int_{\partial B_\varepsilon} X \cdot \left(-\frac{x}{\varepsilon}\right) d\mathcal{H}^1.$$

On ∂B_ε , $X = x/\varepsilon^2$ and $X \cdot (-x/\varepsilon) = -|x|^2/\varepsilon^3 = -1/\varepsilon$, so the second term equals $-\varepsilon^{-1} \mathcal{H}^1(\partial B_\varepsilon) = -\varepsilon^{-1} \cdot 2\pi\varepsilon = -2\pi$, independent of ε . Hence

$$\int_{\partial^* E} \frac{\nu_r}{|x|} d\mathcal{H}^1 = \int_{\partial^* E} X \cdot \nu d\mathcal{H}^1 = 2\pi,$$

which is (60b). (No limit in ε is even needed: the left side does not depend on ε , and the inner-circle flux is exactly 2π for every admissible ε .)

(iii) *The radial field, singularity integrable.* Consider $Y(x) := e_r = x/|x|$, smooth on $\mathbb{R}^2 \setminus \{0\}$, with

$$\operatorname{div} Y = \sum_i \partial_i \frac{x_i}{|x|} = \sum_i \left(\frac{1}{|x|} - \frac{x_i^2}{|x|^3} \right) = \frac{2}{|x|} - \frac{|x|^2}{|x|^3} = \frac{1}{|x|} \quad (x \neq 0).$$

Again fix $0 < \varepsilon < r_0$ and apply Gauss–Green to the C^1 bounded field Y over $E_\varepsilon = E \setminus \overline{B_\varepsilon(0)}$:

$$\int_{E_\varepsilon} \frac{dx}{|x|} = \int_{E_\varepsilon} \operatorname{div} Y dx = \int_{\partial^* E} Y \cdot \nu d\mathcal{H}^1 + \int_{\partial B_\varepsilon} Y \cdot \left(-\frac{x}{\varepsilon}\right) d\mathcal{H}^1.$$

On ∂B_ε , $Y = x/\varepsilon$ and $Y \cdot (-x/\varepsilon) = -|x|^2/\varepsilon^2 = -1$, so the inner term is $-\mathcal{H}^1(\partial B_\varepsilon) = -2\pi\varepsilon$. Thus

$$\int_{\partial^* E} \nu_r d\mathcal{H}^1 = \int_{E_\varepsilon} \frac{dx}{|x|} + 2\pi\varepsilon.$$

Now let $\varepsilon \downarrow 0$. The left side is fixed. On the right, $|x|^{-1} \in L^1_{\text{loc}}(\mathbb{R}^2)$ (in polar coordinates $\int_{B_R} |x|^{-1} dx = 2\pi R < \infty$) and E has finite measure, so $|x|^{-1} \mathbf{1}_E \in L^1(\mathbb{R}^2)$; by dominated convergence

$\int_{E_\varepsilon} |x|^{-1} dx \rightarrow \int_E |x|^{-1} dx = I_E$, while $2\pi\varepsilon \rightarrow 0$. Therefore $\int_{\partial^* E} \nu_r d\mathcal{H}^1 = I_E$, which is (60c). The constant I_E is finite because $|x|^{-1} \mathbf{1}_E \in L^1$. $\square$

Remark 6.2 (On the singularity). The two divergence-theorem identities are genuinely different at the source. In (60b) the field $X = x/|x|^2$ has a non-integrable $|x|^{-1}$ growth at 0 and a non-zero residual flux 2π through any small circle; excising B_ε is mandatory, and the 2π is precisely that residue (the planar Green-function source). In (60c) the field $Y = e_r$ is bounded, $\operatorname{div} Y = |x|^{-1}$ is integrable, and the inner-circle flux $2\pi\varepsilon \rightarrow 0$ vanishes; the excision is a convenience and the resulting I_E is a finite, well-defined number. We invoke Lemma 6.1 for $E = \tilde{E}$, whose reduced boundary satisfies the hypothesis $\partial^* \tilde{E} \subset \{|x - z| \geq \rho_i\}$ with $r_0 = \rho_i$ by (57), and whose interior contains z since $B_{\rho_i}(z) \subset \tilde{E}$.

6.2. Sharp fold content. The fold content is the deviation of M from the graph value 1. We bound its total mass with no quantitative-isoperimetric square-root loss; this is the crux of the sharp reduction.

Proposition 6.3 (Sharp fold content). *With the inset $\tilde{E}$ and the trapping data (56),*

$$\int_{S^1} (M(\theta) - 1) d\theta = 2 \int_{\partial \tilde{E} \cap \{\nu_r < 0\}} \frac{|\nu_r|}{|x - z|} d\mathcal{H}^1 \leq \frac{1}{\rho_i} \left(\operatorname{Exc} + \frac{a\eta}{\rho_i \hat{r}} \right) \leq C(\operatorname{Exc} + a\eta), \quad (61)$$

with C absolute. In particular the integrand is non-negative, so $M \geq 1$ a.e., and

$$|\{\theta \in S^1 : M(\theta) \geq 3\}| \leq \frac{1}{2} \int_{S^1} (M - 1) d\theta \leq C(\operatorname{Exc} + a\eta). \quad (62)$$

Proof. The fold identity. Subtract (60b) from (60a) of Lemma 6.1 (legitimate by Remark 6.2); since $\int_{S^1} 1 d\theta = 2\pi$ equals the right-hand side of (60b),

$$\int_{S^1} (M - 1) d\theta = \int_{\partial \tilde{E}} \frac{|\nu_r| - \nu_r}{|x - z|} d\mathcal{H}^1 = 2 \int_{\partial \tilde{E} \cap \{\nu_r < 0\}} \frac{|\nu_r|}{|x - z|} d\mathcal{H}^1 \geq 0,$$

using $|\nu_r| - \nu_r = 0$ where $\nu_r \geq 0$ and $= 2|\nu_r|$ where $\nu_r < 0$. This is the first equality of (61). (Pointwise $M(\theta) = 2m(\theta) + 1 \geq 1$ a.e. by the slice structure (59), since every slice contains $(0, r_1)$; the content of the identity is the *global* bound on the excess $\int (M - 1)$, which is exactly twice the inward-pointing radial flux.)

From the radial fold flux to $P_r - I_{\tilde{E}}$. On the folded part $\partial \tilde{E} \cap \{\nu_r < 0\}$ we bound $|x - z| \geq \rho_i$ (by (57)), so

$$\int_{S^1} (M - 1) d\theta \leq \frac{2}{\rho_i} \int_{\partial \tilde{E} \cap \{\nu_r < 0\}} |\nu_r| d\mathcal{H}^1 = \frac{2F_r}{\rho_i}, \quad F_r := \int_{\partial \tilde{E} \cap \{\nu_r < 0\}} |\nu_r| d\mathcal{H}^1. \quad (63)$$

Write $P_r := \int_{\partial \tilde{E}} |\nu_r| d\mathcal{H}^1$ for the total radial perimeter. Splitting according to the sign of ν_r ,

$$\int_{\partial \tilde{E}} \nu_r d\mathcal{H}^1 = \int_{\{\nu_r \geq 0\}} |\nu_r| d\mathcal{H}^1 - \int_{\{\nu_r < 0\}} |\nu_r| d\mathcal{H}^1 = (P_r - F_r) - F_r = P_r - 2F_r.$$

By (60c), the left side is $I_{\tilde{E}}$, hence

$$2F_r = P_r - I_{\tilde{E}}. \quad (64)$$

Upper bound on P_r . Since $|\nu_r| = |\nu \cdot e_r| \leq |\nu| = 1$ pointwise, the radial perimeter is dominated by the perimeter:

$$P_r = \int_{\partial \tilde{E}} |\nu_r| d\mathcal{H}^1 \leq \int_{\partial \tilde{E}} 1 d\mathcal{H}^1 = P(\tilde{E}) = 2\pi\hat{r} + \text{Exc}.$$

The cancellation: lower bound on $I_{\tilde{E}}$. Here is the sharp step. Because $|\tilde{E}| = \mu_E = \pi\hat{r}^2 = |B_{\hat{r}}(z)|$, the indicators have equal mass, so $\int (\mathbf{1}_{\tilde{E}} - \mathbf{1}_{B_{\hat{r}}(z)}) dx = 0$ and we may subtract *any* constant from the kernel without changing the integral. Subtract the constant $1/\hat{r}$:

$$I_{\tilde{E}} - \int_{B_{\hat{r}}(z)} \frac{dx}{|x-z|} = \int (\mathbf{1}_{\tilde{E}} - \mathbf{1}_{B_{\hat{r}}(z)}) \frac{dx}{|x-z|} = \int (\mathbf{1}_{\tilde{E}} - \mathbf{1}_{B_{\hat{r}}(z)}) \left(\frac{1}{|x-z|} - \frac{1}{\hat{r}} \right) dx,$$

the last equality because $\int (\mathbf{1}_{\tilde{E}} - \mathbf{1}_{B_{\hat{r}}(z)}) \cdot \hat{r}^{-1} dx = \hat{r}^{-1} (|\tilde{E}| - |B_{\hat{r}}(z)|) = 0$. In polar coordinates, $\int_{B_{\hat{r}}(z)} |x-z|^{-1} dx = \int_0^{\hat{r}} \int_{S^1} s^{-1} s d\theta ds = 2\pi\hat{r}$, so

$$I_{\tilde{E}} - 2\pi\hat{r} = \int_{\tilde{E} \Delta B_{\hat{r}}(z)} \pm \left(\frac{1}{|x-z|} - \frac{1}{\hat{r}} \right) dx, \quad (65)$$

the integrand being supported on the symmetric difference $\tilde{E} \Delta B_{\hat{r}}(z)$ (with sign $+$ on $\tilde{E} \setminus B_{\hat{r}}(z)$ and $-$ on $B_{\hat{r}}(z) \setminus \tilde{E}$). Now $\tilde{E} \Delta B_{\hat{r}}(z)$ lies in the closed annulus $\{\rho_i \leq |x-z| \leq \rho_e\}$: indeed $B_{\rho_i}(z) \subset \tilde{E} \cap B_{\hat{r}}(z)$ (as $\rho_i \leq \hat{r}$), so neither $\tilde{E}$ nor $B_{\hat{r}}(z)$ differs from the other inside $B_{\rho_i}(z)$; and $\tilde{E} \cup B_{\hat{r}}(z) \subset B_{\rho_e}(z)$ (as $\hat{r} \leq \rho_e$ and $\tilde{E} \subset B_{\rho_e}(z)$), so neither differs outside $B_{\rho_e}(z)$. On that annulus, with both $|x-z|$ and $\hat{r}$ in $[\rho_i, \rho_e]$,

$$\left| \frac{1}{|x-z|} - \frac{1}{\hat{r}} \right| = \frac{|\hat{r} - |x-z||}{\hat{r} |x-z|} \leq \frac{\rho_e - \rho_i}{\hat{r} \rho_i} = \frac{\eta}{\rho_i \hat{r}}. \quad (66)$$

Therefore, taking absolute values in (65) and using (66),

$$|I_{\tilde{E}} - 2\pi\hat{r}| \leq \frac{\eta}{\rho_i \hat{r}} |\tilde{E} \Delta B_{\hat{r}}(z)| = \frac{a\eta}{\rho_i \hat{r}}. \quad (67)$$

Conclusion. Combining (64) with the bound on P_r and the lower bound $I_{\tilde{E}} \geq 2\pi\hat{r} - a\eta/(\rho_i \hat{r})$ from (67),

$$2F_r = P_r - I_{\tilde{E}} \leq (2\pi\hat{r} + \text{Exc}) - \left(2\pi\hat{r} - \frac{a\eta}{\rho_i \hat{r}} \right) = \text{Exc} + \frac{a\eta}{\rho_i \hat{r}}.$$

Inserting into (63) yields the middle inequality of (61). Finally, for $\delta_T \leq \delta_*$ the inset radius obeys $\hat{r} \geq \frac{3}{4}$ (since $\mu_E = \pi - O(\sqrt{\delta_T})$ by Definition 4.7, so $\hat{r} = \sqrt{\mu_E/\pi} \rightarrow 1$), and $\eta \leq C\delta_T^{1/8} \leq \frac{1}{4}$ by (56), whence $\rho_i \geq \hat{r} - \eta \geq \frac{1}{2}$ (using $\rho_e = \rho_i + \eta$ and $\rho_i \leq \hat{r} \leq \rho_e$). Therefore $1/\rho_i \leq 2$ and $1/(\rho_i^2 \hat{r}) \leq 1/(\frac{1}{4} \cdot \frac{3}{4}) \leq 6$, giving the absolute bound $C(\text{Exc} + a\eta)$. The fold-angle estimate (62) follows because $M-1 \geq 2$ on $\{M \geq 3\}$ (as $M = 2m+1$ is odd, $M \geq 3$ means $m \geq 1$), so $2|\{M \geq 3\}| \leq \int_{\{M \geq 3\}} (M-1) \leq \int_{S^1} (M-1)$. $\square$

Remark 6.4 (Why the factor η matters: the improvement over the naive bound). The naive estimate pulls $|x-z|^{-1} \leq \rho_i^{-1}$ out of the integral defining $I_{\tilde{E}} - 2\pi\hat{r}$ before subtracting any constant, giving

$$|I_{\tilde{E}} - 2\pi\hat{r}| = \left| \int (\mathbf{1}_{\tilde{E}} - \mathbf{1}_{B_{\hat{r}}(z)}) \frac{1}{|x-z|} \right| \leq \frac{1}{\rho_i} |\tilde{E} \Delta B_{\hat{r}}(z)| = \frac{a}{\rho_i} \leq Ca,$$

hence $\int (M-1) \leq C(\text{Exc} + a)$. The two estimates differ by the factor $\eta = \rho_e - \rho_i$ on the a -term. This is decisive. By the quantitative isoperimetric inequality (Theorem 2.11), $a \leq C_F \sqrt{\delta_{\text{iso}}} \mu_E$ carries an

unavoidable *square-root* of the isoperimetric deficit; with $\delta_{\text{iso}} \sim \delta_T/\tau \sim \sqrt{\delta_T}$ at the boundary scale this gives only $a \sim \delta_T^{1/4}$, and the naive fold content $\text{Exc} + a \sim \delta_T^{1/4}$ would cap the whole argument at $\delta_T^{1/4}$ closeness. The cancellation (67) replaces a by $a\eta$; since $\eta \leq C(\text{Exc} + a)^{1/2} \leq Ca^{1/2} \sim \delta_T^{1/8}$, the product $a\eta \sim \delta_T^{3/8}$, and through Corollary 6.5 the graph defect becomes $\text{gd} \sim \eta a\eta \sim \delta_T^{1/8} \cdot \delta_T^{3/8} = \delta_T^{1/2}$, matching the closeness $O(\sqrt{\delta_T})$. The single factor η — the legitimacy of subtracting the constant $1/\hat{r}$, because $|\tilde{E}| = |B_{\hat{r}}(z)|$, together with the confinement of $\tilde{E} \Delta B_{\hat{r}}(z)$ to the thin η -annulus — is exactly what removes the FMP square-root from the fold content and makes the reduction sharp.

6.3. The graph defect. The *graph defect* of $\tilde{E}$ about z ,

$$\text{gd}(\tilde{E}) := \inf_{\rho: S^1 \rightarrow (0, \infty)} |\tilde{E} \Delta \{z + r\omega : 0 \leq r < \rho(\omega)\}|,$$

measures how far $\tilde{E}$ is, in L^1 , from being a radial graph about z . The fold content controls it directly.

Corollary 6.5 (Graph defect). *The graph defect of $\tilde{E}$ about the centre z satisfies*

$$\text{gd}(\tilde{E}) \leq \rho_e \eta \int_{S^1} m(\theta) d\theta = \frac{1}{2} \rho_e \eta \int_{S^1} (M - 1) d\theta \leq C \eta (\text{Exc} + a\eta), \quad (68)$$

with C absolute.

Proof. Use the *core profile* $\rho_0(\theta) := r_1(\theta) \in [\rho_i, \rho_e]$, the innermost crossing radius from (59); it is defined for a.e. θ . Set $G_0 := \{z + re^{i\theta} : 0 \leq r < \rho_0(\theta)\}$. By construction the slice of G_0 in direction θ is $(0, \rho_0(\theta)) = (0, r_1(\theta)) \subset \tilde{E}_\theta$, so $G_0 \subset \tilde{E}$ (up to $\mathcal{H}^1$ -null directions), and the slice of $\tilde{E} \setminus G_0$ is exactly the union of caps $(r_2, r_3) \cup \dots \cup (r_{2m}, r_{2m+1})$. By the polar-coordinate area formula (Fubini on $\mathbb{R}^2 \setminus \{z\}$),

$$|\tilde{E} \setminus G_0| = \int_{S^1} \sum_{i=1}^{m(\theta)} \int_{r_{2i}}^{r_{2i+1}} r dr d\theta.$$

On each ray with $m(\theta) \geq 1$, all cap endpoints lie in $[\rho_i, \rho_e]$, so the caps have total radial length $\sum_i (r_{2i+1} - r_{2i}) \leq r_{2m+1} - r_2 \leq \rho_e - \rho_i = \eta$, and each cap sits at radius $r \leq \rho_e$; hence

$$\sum_{i=1}^{m(\theta)} \int_{r_{2i}}^{r_{2i+1}} r dr \leq \rho_e \sum_{i=1}^{m(\theta)} (r_{2i+1} - r_{2i}) \leq \rho_e \eta \mathbf{1}_{\{m(\theta) \geq 1\}}.$$

Therefore

$$\text{gd}(\tilde{E}) \leq |\tilde{E} \Delta G_0| = |\tilde{E} \setminus G_0| \leq \rho_e \eta |\{\theta : m(\theta) \geq 1\}| \leq \rho_e \eta \int_{S^1} m d\theta,$$

where the last step uses $\mathbf{1}_{\{m \geq 1\}} \leq m$ (as m is a non-negative integer). Since $M = 2m + 1$, $\int_{S^1} m d\theta = \frac{1}{2} \int_{S^1} (M - 1) d\theta$, which is the displayed equality; and Proposition 6.3 together with $\rho_e \leq C\hat{r} \leq C$ (for $\delta_T \leq \delta_*$) gives the final bound $\leq C \eta (\text{Exc} + a\eta)$. $\square$

6.4. Realization as a continuous radial graph. The core profile $\rho_0 = r_1$ realises the graph defect but is only BV in θ . We mollify it to a continuous (indeed Lipschitz) profile at a negligible cost.

Proposition 6.6 (Continuous radial-graph realization). *For every $\sigma > 0$ there is a Lipschitz profile $\rho_\sigma: S^1 \rightarrow [\rho_i, \rho_e]$ such that the radial graph $G_\sigma := \{z + r\omega : 0 \leq r < \rho_\sigma(\omega)\}$ satisfies*

$$\|\rho_\sigma - \hat{r}\|_{L^\infty(S^1)} \leq \eta, \quad |\tilde{E} \Delta G_\sigma| \leq |\tilde{E} \setminus G_0| + C\sigma \leq C\eta (\text{Exc} + a\eta) + C\sigma, \quad (69)$$

where $G_0 = \{r < \rho_0\}$ is the core-profile graph of Corollary 6.5 and C is absolute. Thus G_σ is a continuous radial graph that approximates $\tilde{E}$ to within the core-profile graph defect $|\tilde{E} \setminus G_0|$ (itself bounded by Corollary 6.5, hence $\geq \text{gd}(\tilde{E})$) plus the mollification error $C\sigma$. In our application one may take $\sigma = \delta_T$, which is negligible against $|\tilde{E} \setminus G_0| \leq C\sqrt{\delta_T}$.

Proof. Bounded variation of the core profile. The core profile $\rho_0 = r_1: S^1 \rightarrow [\rho_i, \rho_e]$ takes values in a bounded interval, so $\rho_0 \in L^\infty(S^1)$. We bound its essential total variation by the perimeter of $\tilde{E}$ through the circle-crossing function

$$N(\rho) := \mathcal{H}^0(\partial\tilde{E} \cap \partial B_\rho(z))$$

of Lemma 5.4, which is the rigorous substitute for the (false-in-the-BV-jump-case) identification of $\text{TV}(\rho_0)$ with the arclength of the polar graph point-set $\{z + \rho_0(\theta)e^{i\theta}\}$. The key inequality is the layer-cake comparison

$$\text{TV}(\rho_0) = \int_{\rho_i}^{\rho_e} \mathcal{H}^0(\partial^*\{\rho_0 > \rho\}) d\rho \leq \int_{\rho_i}^{\rho_e} N(\rho) d\rho. \quad (70)$$

The first equality is the one-dimensional coarea (layer-cake) formula for the BV function ρ_0 on S^1 [1, Theorem 3.40]: for a.e. ρ the superlevel set $\{\theta: \rho_0(\theta) > \rho\} \subset S^1$ is a finite union of arcs, its essential boundary $\partial^*\{\rho_0 > \rho\}$ is a finite set of points, and the variation measure disintegrates as $|D\rho_0|(S^1) = \int_0^\infty \mathcal{H}^0(\partial^*\{\rho_0 > \rho\}) d\rho$ (the integrand vanishes off $[\rho_i, \rho_e]$ since $\rho_0 \in [\rho_i, \rho_e]$). For the inequality, fix a regular $\rho \in (\rho_i, \rho_e)$ and let $\theta_0 \in \partial^*\{\rho_0 > \rho\}$ be a point where $\rho_0 = r_1$ crosses the level ρ (jumping or continuously). Since $r_1(\theta)$ is the innermost crossing radius of $\partial\tilde{E}$ on the ray R_θ , the crossing of the level ρ at θ_0 is realised by a point $z + \rho e^{i\theta_0}$ of $\partial\tilde{E}$ lying on the circle $\partial B_\rho(z)$: indeed, on one side of θ_0 the inner interval $(0, r_1)$ of the slice (59) reaches past radius ρ and on the other side it does not, so $\partial B_\rho(z)$ separates points of $\tilde{E}^{(1)}$ from points of $\tilde{E}^{(0)}$ arbitrarily near $z + \rho e^{i\theta_0}$, whence $z + \rho e^{i\theta_0} \in \partial\tilde{E}$. Distinct boundary directions θ_0 give distinct points on $\partial B_\rho(z)$, so $\mathcal{H}^0(\partial^*\{\rho_0 > \rho\}) \leq N(\rho)$, proving (70). Now Lemma 5.4 states $\int_0^\infty N(\rho) d\rho = P_\theta := \int_{\partial\tilde{E}} |\nu_\theta| d\mathcal{H}^1 \leq P(\tilde{E})$ (as $|\nu_\theta| \leq 1$). Combining with (70),

$$\text{TV}(\rho_0) \leq \int_{\rho_i}^{\rho_e} N(\rho) d\rho \leq P_\theta \leq P(\tilde{E}) \leq \frac{P(\tilde{E})}{\rho_i} \leq C, \quad (71)$$

the penultimate inequality holding since $\rho_i \leq 1$ (indeed $\rho_i \geq \frac{1}{2}$ for $\delta_T \leq \delta_\star$, so $1/\rho_i \leq 2$), and the last using $P(\tilde{E}) = 2\pi\hat{r} + \text{Exc} \leq C$ for $\delta_T \leq \delta_\star$. The form $\text{TV}(\rho_0) \leq P(\tilde{E})/\rho_i$ is the bound flagged in the mandate; either constant is absolute.

Mollification. Let ψ_σ be a standard non-negative mollifier on S^1 at scale σ ($\int \psi_\sigma = 1$, $\text{supp } \psi_\sigma \subset (-\sigma, \sigma)$, $\|\psi'_\sigma\|_{L^1} \leq C/\sigma$), and set $\rho_\sigma := \psi_\sigma * \rho_0$. Then ρ_σ is smooth, hence Lipschitz; it is a convex average of values of $\rho_0 \in [\rho_i, \rho_e]$, so $\rho_\sigma \in [\rho_i, \rho_e]$; and $\|\rho_\sigma - \hat{r}\|_\infty \leq \rho_e - \rho_i = \eta$ since both ρ_σ and $\hat{r}$ lie in $[\rho_i, \rho_e]$. This gives the first estimate in (69).

Cost of mollification. Convolution moves a BV function in L^1 by at most σ times its total variation:

$$\|\rho_\sigma - \rho_0\|_{L^1(S^1)} = \int_{S^1} \left| \int \psi_\sigma(s) (\rho_0(\theta - s) - \rho_0(\theta)) ds \right| d\theta \leq \int \psi_\sigma(s) \text{TV}(\rho_0) |s| ds \leq \sigma \text{TV}(\rho_0),$$

using $\int_{S^1} |\rho_0(\theta - s) - \rho_0(\theta)| d\theta \leq |s| \text{TV}(\rho_0)$. The symmetric difference of two radial graphs with profiles $\rho_0, \rho_\sigma \in [\rho_i, \rho_e]$ is

$$|\{r < \rho_0\} \Delta \{r < \rho_\sigma\}| = \int_{S^1} \left| \int_{\rho_\sigma(\theta)}^{\rho_0(\theta)} r dr \right| d\theta = \int_{S^1} \frac{|\rho_0^2 - \rho_\sigma^2|}{2} d\theta \leq \rho_e \int_{S^1} |\rho_0 - \rho_\sigma| d\theta \leq \rho_e \sigma \text{TV}(\rho_0),$$

using $|\rho_0^2 - \rho_\sigma^2| = |\rho_0 - \rho_\sigma|(\rho_0 + \rho_\sigma) \leq 2\rho_e |\rho_0 - \rho_\sigma|$. By (71) and $\rho_e \leq C$, this is $\leq C\sigma$. Hence, using $G_0 \subset \tilde{E}$ so that $|\tilde{E} \Delta G_0| = |\tilde{E} \setminus G_0|$,

$$|\tilde{E} \Delta G_\sigma| \leq |\tilde{E} \Delta G_0| + |G_0 \Delta G_\sigma| \leq |\tilde{E} \setminus G_0| + C\sigma,$$

which is the second estimate of (69); the further bound $|\tilde{E} \setminus G_0| \leq C\eta(\text{Exc} + a\eta)$ is Corollary 6.5.

Taking $\sigma = \delta_T$ makes the mollification cost $C\delta_T$, negligible against $|\tilde{E} \setminus G_0| \leq C\sqrt{\delta_T}$. $\square$

Remark 6.7 (Scales, and the role in the reduction). Specialising to the inset at level $\hat{v} \sim \tau = \sqrt{\delta_T}$ (so $\delta_{\text{iso}} \leq C\delta_T/\tau \leq C\sqrt{\delta_T}$, $\text{Exc} \leq C\sqrt{\delta_T}$, and by Theorem 2.11 $a \leq C\sqrt{\delta_{\text{iso}}} \leq C\delta_T^{1/4}$), the annulus width is $\eta \leq C(\text{Exc} + a)^{1/2} \leq Ca^{1/2} \leq C\delta_T^{1/8}$, an *absolute* small amplitude. Then Corollary 6.5 gives

$$\text{gd}(\tilde{E}) \leq C\eta(\text{Exc} + a\eta) \leq C(\delta_T^{1/8}\delta_T^{1/2} + \delta_T^{1/4}\delta_T^{1/4}) = C(\delta_T^{5/8} + \delta_T^{1/2}) \leq C\sqrt{\delta_T},$$

and Proposition 6.6 (with $\sigma = \delta_T$) produces a continuous radial graph G of amplitude $\|\rho - \hat{r}\|_\infty \leq \eta \leq C\delta_T^{1/8}$ with $|\tilde{E} \Delta G| \leq C\sqrt{\delta_T}$. These are precisely the scales fed into the cutting estimate (Lemma 7.5) and the assembly (Theorem 9.1): the graph defect balances the level-carried closeness $|\Omega \Delta \tilde{E}| \sim \sqrt{\delta_T}$ at the optimal exponent.

7. CUTTING THE CAPS LOWERS THE DEFICIT

The construction of Sections 4 to 6 produces from Ω a connected, hole-free inset $\tilde{E}$ trapped in a thin annulus $B_{\rho_i}(z) \subset \tilde{E} \subset B_{\rho_e}(z)$, together with a radial core profile whose graph $G_0 = \{z + r\omega : r < \rho_0(\omega)\}$ satisfies $G_0 \subset \tilde{E}$ and $|\tilde{E} \setminus G_0| = \text{gd}(\tilde{E})$ (Corollary 6.5). Manufacturing the honest graph from $\tilde{E}$ therefore amounts to *excising* the radial caps $\tilde{E} \setminus G_0$, all of which lie in the collar $\{\rho_i < |x - z| < \rho_e\}$. The naive worry is that removing material can only destroy torsion and hence inflate the scale-invariant deficit δ_* . The opposite is true: caps in a thin collar carry very little torsion, and the equal-area comparison ball shrinks faster than the torsion is lost, so δ_* *decreases*. This section proves the two facts that make this precise. They are sign-critical, so we prove them in full and check the inequalities digit by digit.

Throughout, for a bounded open set $D \subset \mathbb{R}^2$ the *torsion function* $u_D \in H_0^1(D)$ is the weak solution of $-\Delta u_D = 1$ in D , $u_D = 0$ on ∂D , and $T(D) := \int_D u_D$ is its torsional rigidity; for a general set of finite measure D we set $T(D) := T(\text{int } D)$ on the open representative produced by our construction, which is all we use. Recall (Theorem 2.13) that u_D is real-analytic in D and, by the comparison principle for the Laplacian, $u_D \geq 0$.

7.1. The torsion as a Green double integral. We first record the representation of T that drives the stability bound. Let $G_D(x, y)$ be the Green function of $-\Delta$ on the open set D with pole at y : the distributional solution of $-\Delta_x G_D(\cdot, y) = \delta_y$ in D , $G_D(\cdot, y) = 0$ on ∂D (Theorem 2.13 and [7, §2.4, §4]). Two classical properties are used.

Lemma 7.1 (Positivity and domain monotonicity of G_D). *For bounded open $D \subset \mathbb{R}^2$ one has $G_D \geq 0$ on $D \times D$. If $E \subset F$ are bounded open sets, then $G_E(x, y) \leq G_F(x, y)$ for all $x, y \in E$.*

Proof. Fix y . The function $x \mapsto G_D(x, y)$ is harmonic on $D \setminus \{y\}$, tends to $+\infty$ as $x \rightarrow y$ (the logarithmic singularity), and vanishes on ∂D ; by the minimum principle it is ≥ 0 on D . This is the positivity.

For monotonicity, fix $y \in E$ and set $w := G_F(\cdot, y) - G_E(\cdot, y)$ on E . Both functions have the *same* singularity $-\frac{1}{2\pi} \log |x - y|$ at y , so w extends to a harmonic function on all of E (the singularities cancel). On ∂E we have $G_E(\cdot, y) = 0$ while $G_F(\cdot, y) \geq 0$ by positivity (since $\partial E \subset F$ up to the boundary points lying in $\overline{F}$, where $G_F \geq 0$ by continuity); hence $w \geq 0$ on ∂E . By the minimum principle for harmonic functions, $w \geq 0$ on E , i.e. $G_E \leq G_F$ on $E \times E$. $\square$

Lemma 7.2 (Green representation of torsion). *For a bounded open set $D \subset \mathbb{R}^2$,*

$$u_D(y) = \int_D G_D(x, y) dx \quad (y \in D), \quad T(D) = \iint_{D \times D} G_D(x, y) dx dy. \quad (72)$$

In particular $T(D) = \iint_{D \times D} G_D \geq 0$, and G_D is symmetric, $G_D(x, y) = G_D(y, x)$.

Proof. The Green function reproduces solutions of the Dirichlet problem: for $f \in C_c^\infty(D)$ the function $v(y) := \int_D G_D(x, y) f(x) dx$ solves $-\Delta v = f$ in D , $v = 0$ on ∂D , by the defining property $-\Delta_x G_D(\cdot, y) = \delta_y$ and the symmetry $G_D(x, y) = G_D(y, x)$ (both standard, [7, Thm. 4.10]). Taking $f \equiv 1$ (approximated in $L^2(D)$ by C_c^∞ functions, with the corresponding solutions converging to u_D in $H_0^1(D)$ by the energy estimate) gives $u_D(y) = \int_D G_D(x, y) dx$. Integrating over $y \in D$ and using Tonelli ($G_D \geq 0$ by Lemma 7.1) yields $T(D) = \int_D u_D = \iint_{D \times D} G_D$. $\square$

7.2. Boundary-layer torsion stability. The next lemma controls how much torsion can change when two sets agree except in a thin annular collar. It is the analytic heart of the section.

Lemma 7.3 (Boundary-layer torsion stability). *Let $z \in \mathbb{R}^2$ and $0 < \rho_i \leq \rho_e$, and put*

$$\tilde{\eta} := \frac{1}{4}(\rho_e^2 - \rho_i^2). \quad (73)$$

Let $E, F \subset \mathbb{R}^2$ be bounded open sets with

$$B_{\rho_i}(z) \subset E \cap F, \quad E \cup F \subset B_{\rho_e}(z), \quad |E \triangle F| = m, \quad (74)$$

so that $E \triangle F$ lies in the collar $\{\rho_i < |x - z| < \rho_e\}$. Then

$$|T(E) - T(F)| \leq 3\tilde{\eta}m. \quad (75)$$

Proof. Set $E_\cap := E \cap F$ and $F_\cup := E \cup F$; these are bounded open sets with $E_\cap \subset E, F \subset F_\cup$ and

$$F_\cup \setminus E_\cap = E \triangle F, \quad |F_\cup \setminus E_\cap| = m.$$

By monotonicity of the Green representation in the domain (Lemmas 7.1 and 7.2), $E_\cap \subset E$ gives $T(E_\cap) \leq T(E)$ and $E \subset F_\cup$ gives $T(E) \leq T(F_\cup)$, and likewise for F ; hence

$$T(E_\cap) \leq T(E) \leq T(F_\cup), \quad T(E_\cap) \leq T(F) \leq T(F_\cup), \quad (76)$$

so that

$$|T(E) - T(F)| \leq T(F_\cup) - T(E_\cap). \quad (77)$$

It therefore suffices to bound $T(F_\cup) - T(E_\cap)$.

Write $A := F_\cup \setminus E_\cap = E \triangle F$, so $F_\cup = E_\cap \cup A$ is a disjoint union and $|A| = m$. Using Lemma 7.2 for both sets and then the domain monotonicity $G_{E_\cap} \leq G_{F_\cup}$ on $E_\cap \times E_\cap$ (Lemma 7.1),

$$\begin{aligned} T(F_\cup) - T(E_\cap) &= \iint_{F_\cup \times F_\cup} G_{F_\cup} - \iint_{E_\cap \times E_\cap} G_{E_\cap} \\ &\leq \iint_{F_\cup \times F_\cup} G_{F_\cup} - \iint_{E_\cap \times E_\cap} G_{F_\cup}. \end{aligned} \quad (78)$$

Splitting $F_\cup \times F_\cup = (E_\cap \cup A) \times (E_\cap \cup A)$ into the four product pieces and cancelling $E_\cap \times E_\cap$, the right-hand side of (78) equals, by the symmetry of $G_{F_\cup}$,

$$2 \int_{E_\cap} \int_A G_{F_\cup}(x, y) dy dx + \iint_{A \times A} G_{F_\cup}(x, y) dy dx. \quad (79)$$

Each inner integral is bounded by the torsion function: for fixed $y \in A$, Lemma 7.2 (with $D = F_\cup$) and positivity give

$$\int_{E_\cap} G_{F_\cup}(x, y) dx \leq \int_{F_\cup} G_{F_\cup}(x, y) dx = u_{F_\cup}(y), \quad \int_A G_{F_\cup}(x, y) dx \leq u_{F_\cup}(y). \quad (80)$$

Therefore

$$T(F_\cup) - T(E_\cap) \leq 2 \int_A u_{F_\cup} + \int_A u_{F_\cup} = 3 \int_A u_{F_\cup}. \quad (81)$$

It remains to bound $u_{F_\cup}$ on the collar by the barrier $\tilde{\eta}$. Since $F_\cup \subset B_{\rho_e}(z)$, comparison gives $u_{F_\cup} \leq u_{B_{\rho_e}(z)}$ on $F_\cup$: indeed $u_{B_{\rho_e}(z)} - u_{F_\cup}$ is $-\Delta$ -harmonic in $F_\cup$ (both have $-\Delta(\cdot) = 1$) and on $\partial F_\cup$ equals $u_{B_{\rho_e}(z)} \geq 0$, so it is ≥ 0 in $F_\cup$ by the minimum principle. The radial torsion function of the disc is, explicitly,

$$u_{B_{\rho_e}(z)}(y) = \frac{\rho_e^2 - |y - z|^2}{4}, \quad (82)$$

which indeed solves $-\Delta u = 1$ with $u = 0$ on $|y - z| = \rho_e$. Now $A = E \triangle F$ lies in the collar $\{\rho_i < |x - z| < \rho_e\}$, by (74): any point of $E \triangle F$ lies in $F_\cup \subset B_{\rho_e}(z)$ (so $|x - z| < \rho_e$) and outside $E_\cap \supset B_{\rho_i}(z)$ (so $|x - z| \geq \rho_i$; the boundary sphere is $\mathcal{H}^2$ -null). Hence for $y \in A$,

$$u_{F_\cup}(y) \leq u_{B_{\rho_e}(z)}(y) = \frac{\rho_e^2 - |y - z|^2}{4} \leq \frac{\rho_e^2 - \rho_i^2}{4} = \tilde{\eta}. \quad (83)$$

Combining (81) and (83),

$$T(F_\cup) - T(E_\cap) \leq 3 \int_A u_{F_\cup} \leq 3 \tilde{\eta} |A| = 3 \tilde{\eta} m,$$

and (75) follows from (77). $\square$

Remark 7.4 (The barrier constant). The width of the collar enters only through $\tilde{\eta} = \frac{1}{4}(\rho_e^2 - \rho_i^2)$, the maximal value of the disc torsion function on the inner sphere. With $\rho_i = \hat{r} - O(\eta)$ and $\rho_e = \hat{r} + O(\eta)$ one has $\tilde{\eta} = \frac{1}{4}(\rho_e + \rho_i)(\rho_e - \rho_i) \leq \frac{1}{2}\rho_e \eta \leq C\eta$, so $\tilde{\eta}$ is of the same order as the annulus width $\eta = \rho_e - \rho_i$. In the unit-scale normalisation $\rho_e = 1 + \eta$, $\rho_i = 1 - \eta$ one gets exactly $\tilde{\eta} = \eta$.

7.3. Cutting boundary material decreases δ_* . Recall the scale-invariant relative deficit of a finite-measure set D ,

$$\delta_*(D) = \frac{T(B_D) - T(D)}{T(B_D)}, \quad T(B_D) = \frac{|D|^2}{8\pi}, \quad (84)$$

where B_D is any ball with $|B_D| = |D|$ (so $T(B_D) = |B_D|^2/(8\pi)$ by scaling from $T(B_1) = \pi/8$). Equivalently, $\delta_*(D) = 1 - 8\pi T(D)/|D|^2$. We now show that excising boundary material confined to a thin collar lowers δ_* , provided the torsion density of the larger set exceeds three barrier-widths.

Lemma 7.5 (Cutting boundary material decreases δ_*). *Let $z \in \mathbb{R}^2$, $0 < \rho_i \leq \rho_e$ and $\tilde{\eta} = \frac{1}{4}(\rho_e^2 - \rho_i^2)$. Let E, G be bounded open sets with*

$$B_{\rho_i}(z) \subset G \subset E \subset B_{\rho_e}(z), \quad E \setminus G \subset \{\rho_i < |x - z| < \rho_e\}. \quad (85)$$

If

$$3\tilde{\eta} \leq \frac{T(E)}{|E|}, \quad (86)$$

then $\delta_*(G) \leq \delta_*(E)$.

Proof. Write

$$V := |E|, \quad g := |E \setminus G| = |E| - |G| = V - |G|, \quad T := T(E),$$

so that $|G| = V - g$ and, since $\emptyset \neq B_{\rho_i}(z) \subset G \subset E$, we have $0 \leq g \leq V$ and $V > 0$.

Step 1: lower bound for $T(G)$. Apply Lemma 7.3 with $F := E$ and the present G, E . The hypotheses (74) hold: $B_{\rho_i}(z) \subset G \cap E = G$, $G \cup E = E \subset B_{\rho_e}(z)$, and $|G \triangle E| = |E \setminus G| = g$ with $E \triangle G = E \setminus G$ inside the collar by (85). Hence

$$T(E) - T(G) \leq 3\tilde{\eta}g, \quad \text{i.e.} \quad T(G) \geq T - 3\tilde{\eta}g. \quad (87)$$

Step 2: the deficit comparison reduces to a torsion inequality. By (84), with $|E| = V$ and $|G| = V - g$,

$$\delta_*(E) - \delta_*(G) = \left(1 - \frac{8\pi T}{V^2}\right) - \left(1 - \frac{8\pi T(G)}{(V-g)^2}\right) = 8\pi \left[\frac{T(G)}{(V-g)^2} - \frac{T}{V^2}\right]. \quad (88)$$

Thus $\delta_*(G) \leq \delta_*(E)$ holds if and only if $\frac{T(G)}{(V-g)^2} \geq \frac{T}{V^2}$, i.e. (multiplying by $(V-g)^2 \geq 0$) if and only if

$$T(G) \geq T \frac{(V-g)^2}{V^2} = T \left(1 - \frac{g}{V}\right)^2. \quad (89)$$

We now prove (89). (When $g = V$, $|G| = 0$ is impossible since $G \supset B_{\rho_i}(z)$, so in fact $g < V$; but the argument below covers $g \leq V$ verbatim.)

Step 3: the sign-critical chain. Expand the right-hand side of (89):

$$T \left(1 - \frac{g}{V}\right)^2 = T - \frac{2Tg}{V} + \frac{Tg^2}{V^2}. \quad (90)$$

Subtracting (90) from the lower bound (87) for $T(G)$,

$$\begin{aligned} T(G) - T \left(1 - \frac{g}{V}\right)^2 &\geq (T - 3\tilde{\eta}g) - \left(T - \frac{2Tg}{V} + \frac{Tg^2}{V^2}\right) \\ &= \frac{2Tg}{V} - \frac{Tg^2}{V^2} - 3\tilde{\eta}g = g \left(\frac{2T}{V} - \frac{Tg}{V^2} - 3\tilde{\eta}\right). \end{aligned} \quad (91)$$

We bound the bracket from below. Since $0 \leq g \leq V$ we have $0 \leq g/V \leq 1$, hence $2 - g/V \geq 1$, and therefore

$$\frac{2T}{V} - \frac{Tg}{V^2} = \frac{T}{V} \left(2 - \frac{g}{V}\right) \geq \frac{T}{V} \geq 3\tilde{\eta}, \quad (92)$$

the last inequality being exactly hypothesis (86) ($T/V = T(E)/|E| \geq 3\tilde{\eta}$). Consequently the bracket in (91) is $\geq \frac{T}{V} - 3\tilde{\eta} \geq 0$, and since $g \geq 0$ the whole expression (91) is ≥ 0 . This is (89), which by Step 2 yields $\delta_*(G) \leq \delta_*(E)$. $\square$

Remark 7.6 (The hypothesis (86) holds with room to spare). In the application $E = \tilde{E}$ is the inset of Definition 4.7, of measure $\mu_E = |\tilde{E}| = \pi - O(\sqrt{\delta_T}) \geq \pi/2$ and deficit $\delta_*(\tilde{E}) \leq \frac{1}{2}$ for $\delta_T \leq \delta_*$. Then

$$\frac{T(\tilde{E})}{|\tilde{E}|} = (1 - \delta_*(\tilde{E})) \frac{|\tilde{E}|}{8\pi} \geq (1 - \frac{1}{2}) \cdot \frac{\pi/2}{8\pi} = \frac{1}{32},$$

using $T(\tilde{E}) = (1 - \delta_*(\tilde{E}))T(B_{\tilde{E}}) = (1 - \delta_*(\tilde{E}))|\tilde{E}|^2/(8\pi)$ from (84). On the other hand $\tilde{\eta} \leq \frac{1}{2}\rho_e\eta \leq C\eta \leq C(\delta_T/\tau)^{1/4} \rightarrow 0$ as $\delta_T \rightarrow 0$ (Theorem 5.8, with $\tau = \sqrt{\delta_T}$ giving $\eta \leq C\delta_T^{1/8}$). Hence $3\tilde{\eta} \leq \frac{1}{32} \leq T(\tilde{E})/|\tilde{E}|$ for all $\delta_T \leq \delta_*$, and (86) is satisfied with a fixed margin. The threshold $1/32$ is not sharp and is recorded only to fix δ_* .

Remark 7.7 (The application: cutting the caps). Lemma 7.5 is used in the assembly (Theorem 9.1) with $E = \tilde{E}$ and $G = G_0 = \{z + r\omega : r < \rho_0(\omega)\}$ the radial core profile of Corollary 6.5. Then $B_{\rho_i}(z) \subset G_0 \subset \tilde{E} \subset B_{\rho_e}(z)$ holds by Corollary 6.5 and Theorem 5.8, the difference $\tilde{E} \setminus G_0$ is the union of the radial caps, all contained in the collar $\{\rho_i < |x - z| < \rho_e\}$, and (86) holds by Remark 7.6; hence cutting the caps lowers the deficit,

$$\delta_*(G_0) \leq \delta_*(\tilde{E}).$$

This is the one place where the geometry of Lemma 7.5 is needed: all hypotheses (85)–(86) are met because the discarded material is precisely low-torsion boundary material in the η -collar about the *single* centre z .

Remark 7.8 (The other deficit step is *not* an instance of Lemma 7.5). The earlier reduction step — passing from the full superlevel set $\{u > \hat{v}\}$ to its largest connected component $\tilde{E}$, i.e. discarding the spillover components of total measure $|\{u > \hat{v}\} \setminus \tilde{E}| \leq D_I(\hat{v})^2/(16\pi^3)$ (Lemma 4.6) — is *not* covered by Lemma 7.5, and we do not claim it is. Lemma 7.5 requires $G \subset E \subset B_{\rho_e}(z)$ with $E \setminus G$ inside one collar about z ; but the discarded spillover components are distinct level components, which need not lie in the trapping ball $B_{\rho_e}(z)$ of $\tilde{E}$ (they can sit elsewhere in Ω), so the hypothesis $E \subset B_{\rho_e}(z)$ may fail for $E = \{u > \hat{v}\}$. The correct tool for that step is the deficit transfer Lemma 4.3 of Section 4 combined with the *exact* additivity of torsion over connected components (the function $u - \hat{v}$ is the common torsion function of every component of $\{u > \hat{v}\}$); this is carried out in Proposition 4.8, yielding $\delta_*(\tilde{E}) \leq (\pi^2/\mu_E^2)\delta_*(\Omega) + O(\delta_T^2/\tau^2)$ with no appeal to the collar geometry. Thus the full chain

$$\delta_*(G_0) \leq \delta_*(\tilde{E}) \leq \frac{\pi^2}{\mu_E^2}\delta_*(\Omega) + O\left(\frac{\delta_T^2}{\tau^2}\right)$$

combines Lemma 7.5 (first inequality, this section) with Lemma 4.3/Proposition 4.8 (second inequality, Section 4); the two play distinct roles and rest on distinct hypotheses.

Remark 7.9 (Why the deficit decreases — the mechanism). The single inequality (92) is the entire content. Cutting a cap of measure g costs at most $3\tilde{\eta}g$ in torsion (*linear* in g , with a small slope

$3\tilde{\eta} \rightarrow 0$), but it shrinks the equal-area comparison ball, lowering its torsion $T(B_G) = |G|^2/(8\pi)$ by $\approx \frac{2T(B_E)}{V}g = \frac{V}{4\pi}g$, i.e. by a slope $\approx \frac{2T}{V}$ after accounting for the deficit factor. Because the unperturbed torsion density T/V of a near-disc is of order 1 while the collar slope $3\tilde{\eta}$ vanishes, the denominator $T(B_G)$ drops faster than the numerator $T(G)$, and $\delta_* = 1 - T/T(B)$ decreases. There is no square-root loss and no constant that degrades as $\delta_T \rightarrow 0$: the bound is exact and the margin in (86) only widens.

8. PERTURBATIVE STABILITY OF NEARLY SPHERICAL SETS

This section proves, in full and self-containedly, the perturbative torsional stability of small-amplitude radial graphs consumed by the reduction of Section 9. The result is a Fuglede-type estimate *for the torsion functional*: a planar set whose boundary is a graph of small amplitude over a circle, after the natural volume and barycentre normalisations, has scale-invariant torsional deficit controlling the square of its Fraenkel asymmetry. We invoke no selection principle, no spectral black box, and no quantitative isoperimetric inequality. The proof is an explicit second-order torsion computation through the dual (complementary) variational principle, with an admissible vector field built by hand; its only nonclassical ingredient is a frequency-dependent damping of the harmonic corrector that controls high-frequency boundary oscillations and lets the hypothesis be merely $\|\varphi\|_{L^\infty} \leq \eta_G$.

8.1. Statement and standing normalisation. Throughout, $S^1 = \mathbb{R}/2\pi\mathbb{Z}$, $\omega(\theta) = (\cos \theta, \sin \theta)$, and for $\varphi \in \text{Lip}(S^1)$ with $\|\varphi\|_{L^\infty} < 1$ and $z \in \mathbb{R}^2$ we write

$$G = G(z, \hat{r}_G, \varphi) := \{z + r\omega(\theta) : \theta \in S^1, 0 \leq r < \hat{r}_G(1 + \varphi(\theta))\} \quad (93)$$

for the radial graph of amplitude profile φ , base radius $\hat{r}_G = \sqrt{|G|/\pi}$ and centre z . We recall that, for a bounded set E of finite measure with torsion $T(E) = \int_E u_E$ ($-\Delta u_E = 1$, $u_E \in H_0^1(E)$),

$$\delta_*(E) = \frac{T(B_E) - T(E)}{T(B_E)}, \quad T(B_E) = \frac{|E|^2}{8\pi}, \quad \mathcal{A}(E) = \inf_{x_0} \frac{|E \triangle B(x_0, r_E)|}{|E|}, \quad r_E = \sqrt{|E|/\pi}, \quad (94)$$

are the scale-invariant deficit and the Fraenkel asymmetry; both are invariant under translations and dilations.

Theorem 8.1 (Radial-graph torsion stability). *There are absolute constants*

$$\eta_G = \frac{1}{150}, \quad C_G = 48, \quad (95)$$

such that the following holds. Let $G = G(z, \hat{r}_G, \varphi)$ be a radial graph as in (93) with $\varphi \in \text{Lip}(S^1)$, $\|\varphi\|_{L^\infty} \leq \eta_G$, and barycentre at z . Then

$$\mathcal{A}(G)^2 \leq C_G \delta_*(G). \quad (96)$$

The statement is scale-invariant; G need not have area π . It uses no property of the original set Ω and no selection.

Since δ_* and $\mathcal{A}$ are dilation- and translation-invariant, replacing G by $\hat{r}_G^{-1}(G - z)$ we assume for the proof

$$z = 0, \quad \hat{r}_G = 1, \quad \text{so} \quad |G| = \pi, \quad (97)$$

which fixes the volume constraint

$$|G| = \int_0^{2\pi} \frac{(1 + \varphi)^2}{2} d\theta = \pi \iff \int_0^{2\pi} (2\varphi + \varphi^2) d\theta = 0. \quad (98)$$

Under (97), $T(B_G) = \pi/8$, while $u_{B_1} = (1 - |x|^2)/4$ and $T(B_1) = \pi/8$. Expand φ in its real Fourier series,

$$\varphi(\theta) = a_0 + \sum_{k \geq 1} (a_k \cos k\theta + b_k \sin k\theta), \quad \|\varphi\|_{L^2(S^1)}^2 = \pi \left(2a_0^2 + \sum_{k \geq 1} (a_k^2 + b_k^2) \right). \quad (99)$$

Lipschitz regularity gives $\sum_k k^2(a_k^2 + b_k^2) < \infty$, so the series below converge in the indicated norms.

8.2. The dual (complementary) variational principle. The torsion has two dual characterisations: the Dirichlet principle $T(E) = \max_{w \in H_0^1(E)} (2 \int_E w - \int_E |\nabla w|^2)$ (attained at $w = u_E$), and the complementary principle below, which produces *upper* bounds on $T(E)$ — equivalently *lower* bounds on the deficit — from explicit divergence-constrained fields.

Lemma 8.2 (Complementary principle). *Let $E \subset \mathbb{R}^2$ be bounded open with finite perimeter and torsion function $u_E \in H_0^1(E)$. For every $\sigma \in L^2(E; \mathbb{R}^2)$ with $\operatorname{div} \sigma = -1$ in $\mathcal{D}'(E)$,*

$$T(E) \leq \int_E |\sigma|^2, \quad (100)$$

with equality iff $\sigma = \nabla u_E$.

Proof. Since $u_E \in H_0^1(E)$ and $\operatorname{div} \sigma = -1$, the pairing $\int_E \sigma \cdot \nabla u_E = -\langle \operatorname{div} \sigma, u_E \rangle = \int_E u_E$ is well defined, and $\int_E |\nabla u_E|^2 = \int_E u_E = T(E)$ by testing $-\Delta u_E = 1$ with u_E . Hence

$$0 \leq \int_E |\sigma - \nabla u_E|^2 = \int_E |\sigma|^2 - 2 \int_E \sigma \cdot \nabla u_E + \int_E |\nabla u_E|^2 = \int_E |\sigma|^2 - T(E),$$

which is (100); equality forces $\sigma = \nabla u_E$ a.e. $\square$

The disc field $\sigma_0 := \nabla u_{B_1} = -x/2$ satisfies $\operatorname{div} \sigma_0 = -1$ and $\int_{B_1} |\sigma_0|^2 = T(B_1) = \pi/8$. We perturb it into an admissible field on G .

8.3. The comparison field. Set the corrector coefficients

$$\alpha_k := \tfrac{1}{2} m_k a_k, \quad \beta_k := \tfrac{1}{2} m_k b_k, \quad m_k := \min\left(1, \tfrac{2}{k}\right) \in (0, 1], \quad k \geq 1. \quad (101)$$

The damping m_k equals 1 at the critical mode $k = 2$ and decays like $2/k$ for $k > 2$, so the corrector *slope* stays bounded:

$$k|\alpha_k| = \tfrac{1}{2} m_k k |a_k| \leq |a_k|, \quad k|\beta_k| \leq |b_k|. \quad (102)$$

Define the harmonic corrector and its normal trace on ∂B_1 ,

$$h(r, \theta) := \sum_{k \geq 1} r^k (\alpha_k \cos k\theta + \beta_k \sin k\theta), \quad c(\theta) := \partial_r h(1, \theta) = \sum_{k \geq 1} k (\alpha_k \cos k\theta + \beta_k \sin k\theta). \quad (103)$$

By (102) and Lipschitz regularity, ∇h and c are in L^2 , $\Delta h = 0$ in $\mathbb{R}^2$, and

$$\|c\|_{L^2(S^1)}^2 = \pi \sum_{k \geq 1} k^2 (\alpha_k^2 + \beta_k^2) \leq \pi \sum_{k \geq 1} (a_k^2 + b_k^2) \leq \|\varphi\|_{L^2}^2. \quad (104)$$

The comparison field on G is

$$\sigma(x) := \begin{cases} \nabla(u_{B_1} + h)(x), & x \in G, |x| \leq 1, \\ \left(-\frac{r}{2} + \frac{c(\theta)}{r}\right) \omega(\theta), & x = r\omega(\theta) \in G, |x| > 1. \end{cases} \quad (105)$$

The collar line is used only where $\varphi(\theta) > 0$, i.e. where G protrudes beyond B_1 .

Lemma 8.3 (Admissibility). *The field σ of (105) lies in $L^2(G; \mathbb{R}^2)$ and satisfies $\operatorname{div} \sigma = -1$ in $\mathcal{D}'(G)$.*

Proof. On $\{|x| < 1\}$, $\operatorname{div} \sigma = \Delta(u_{B_1} + h) = -1 + 0 = -1$. On the collar $\{|x| > 1\}$, for a radial field $f(r)\omega$ one has $\operatorname{div}(f\omega) = \frac{1}{r} \frac{d}{dr}(rf)$; with $f = -\frac{r}{2} + \frac{c}{r}$ this is $\frac{1}{r} \frac{d}{dr}(-\frac{r^2}{2} + c) = -1$. Across the interface $\Gamma = \partial B_1 \cap G$, the normal (radial) components match: from inside $\sigma \cdot \omega = \partial_r(u_{B_1} + h)|_{r=1} = -\frac{1}{2} + c(\theta)$, from the collar $\sigma \cdot \omega = -\frac{1}{2} + c(\theta)$. Hence the distributional divergence carries no surface measure on Γ and equals -1 on all of G . Square-integrability: ∇u_{B_1} is smooth, $\nabla h \in L^2(B_{1+\eta_G})$ by (102), and the collar is a thin set of width $\leq \eta_G$ on which $|\sigma|^2 \leq 2(\frac{r}{2})^2 + 2c(\theta)^2$. $\square$

8.4. The exact second-order energy. We compute $\int_G |\sigma|^2$ exactly and separate the quadratic form from a genuinely cubic remainder. Write $p := \varphi(\theta)$ and

$$\Phi_{\text{col}}(r, \theta) := \left(-\frac{r}{2} + \frac{c}{r}\right)^2 r, \quad \Phi_{\text{in}}(r, \theta) := \left[(-\frac{r}{2} + \partial_r h)^2 + \left(\frac{1}{r} \partial_\theta h\right)^2\right] r, \quad (106)$$

the polar energy densities of the collar and interior fields.

Lemma 8.4 (Energy identity). *With σ as in (105),*

$$\int_G |\sigma|^2 = \frac{\pi}{8} + \pi \sum_{k \geq 1} k(\alpha_k^2 + \beta_k^2) + \mathcal{N}, \quad \mathcal{N} = \int_{\{p > 0\}} \int_1^{1+p} \Phi_{\text{col}} dr d\theta - \int_{\{p < 0\}} \int_{1+p}^1 \Phi_{\text{in}} dr d\theta. \quad (107)$$

Proof. Split $\int_G = \int_{B_1} - \int_{B_1 \setminus G} + \int_{G \setminus B_1}$. Over the full disc, $\int_{B_1} |\nabla(u_{B_1} + h)|^2 = \int_{B_1} |\nabla u_{B_1}|^2 + 2 \int_{B_1} \nabla u_{B_1} \cdot \nabla h + \int_{B_1} |\nabla h|^2$. The first term is $\pi/8$; the cross term vanishes, $\int_{B_1} \nabla u_{B_1} \cdot \nabla h = - \int_{B_1} u_{B_1} \Delta h + \int_{\partial B_1} u_{B_1} \partial_\nu h = 0$ (as $\Delta h = 0$, $u_{B_1}|_{\partial B_1} = 0$); and, by orthogonality of the harmonic modes, $\int_{B_1} |\nabla h|^2 = \pi \sum_k k(\alpha_k^2 + \beta_k^2)$ since $\int_{B_1} |\nabla(r^k \cos k\theta)|^2 = \pi k$. The dimple $-\int_{B_1 \setminus G} |\nabla(u_{B_1} + h)|^2$ (where $p < 0$) and the collar $+\int_{G \setminus B_1} |\sigma|^2$ (where $p > 0$) are precisely $\mathcal{N}$. $\square$

Lemma 8.5 (Quadratic form and remainder). *Assume $\|\varphi\|_{L^\infty} \leq \frac{1}{150}$ and the corrector (101). Then*

$$\int_G |\sigma|^2 = \frac{\pi}{8} + \frac{1}{4} \int_{S^1} \varphi + \frac{3}{8} \int_{S^1} \varphi^2 + \pi \sum_{k \geq 1} (k\alpha_k^2 - k\alpha_k a_k) + \pi \sum_{k \geq 1} (k\beta_k^2 - k\beta_k b_k) + \mathcal{R}, \quad (108)$$

where the remainder obeys the explicit per-mode bound

$$|\mathcal{R}| \leq C_0 \|\varphi\|_{L^\infty} \sum_{k \geq 1} k(a_k^2 + b_k^2) \quad (C_0 \text{ absolute}), \quad (109)$$

which is cubic in the amplitude but carries an $\dot{H}^{1/2}$ factor $\sum_k k(a_k^2 + b_k^2)$. The naive global cubic bound $|\mathcal{R}| \leq 10 \|\varphi\|_{L^\infty} \|\varphi\|_{L^2}^2$ is false: see Remark 8.6 for the obstruction and the consequence for the absorption step.

Proof. We expand $\mathcal{N}$ from (107) to second order in p and collect the cubic tail into $\mathcal{R}$. The two boundary densities are C^∞ in r on $[\frac{1}{2}, 2] \supset [1 - \eta_G, 1 + \eta_G]$, with values and r -derivatives at $r = 1$

$$\Phi_{\text{col}}(1) = \left(\frac{1}{2} - c\right)^2, \quad \Phi_{\text{in}}(1) = \left(\frac{1}{2} - c\right)^2 + s^2, \quad \Phi'_{\text{col}}(1) = \frac{3}{4} - c - c^2, \quad (110)$$

where $s(\theta) := \frac{1}{r} \partial_\theta h(1, \theta)$, $\|s\|_{L^2} = \|c\|_{L^2}$ (the tangential and normal traces of a harmonic function on ∂B_1 have equal L^2 norm). For each θ , Taylor's theorem with integral remainder gives, on the collar branch $p > 0$,

$$\int_1^{1+p} \Phi_{\text{col}} dr = p \Phi_{\text{col}}(1) + \frac{1}{2} p^2 \Phi'_{\text{col}}(1) + \int_1^{1+p} \frac{1}{2} (1+p-r)^2 \Phi''_{\text{col}}(r) dr,$$

and analogously on the dimple branch $p < 0$ for Φ_{in} . We keep the *disc* part of the first two terms in the quadratic form and put the c, s -dependent part and the third-order tail into $\mathcal{R}$. Concretely, writing $\Phi_{\text{col}}(1) = \frac{1}{4} - c + c^2$ and $\frac{1}{2} \Phi'_{\text{col}}(1) = \frac{3}{8} - \frac{c}{2} - \frac{c^2}{2}$, the kept quadratic contribution of the collar branch is $\int_{\{p>0\}} (\frac{1}{4} p - pc + \frac{3}{8} p^2)$, and the residue $\int_{\{p>0\}} (p c^2 - \frac{1}{2} p^2 (c + c^2)) + (\text{tail})$ goes to $\mathcal{R}$; the dimple branch shares the same kept part $\frac{1}{4} p - pc + \frac{3}{8} p^2$ but contributes *further* residues to $\mathcal{R}$, namely $\int_{\{p<0\}} p s^2$ (from $\Phi_{\text{in}}(1)$) and the c, s -dependent part of $\frac{1}{2} p^2 \Phi'_{\text{in}}(1)$, the latter *not* reducible to the trace c alone (see (111) below). Since $\Phi_{\text{col}}(1)$ and $\Phi_{\text{in}}(1)$ share the kept part $\frac{1}{4} - c$, the kept quadratic over the full circle is

$$\int_{S^1} \left(\frac{1}{4} \varphi - \varphi c + \frac{3}{8} \varphi^2 \right) = \frac{1}{4} \int \varphi + \frac{3}{8} \int \varphi^2 - \pi \sum_k k (\alpha_k a_k + \beta_k b_k),$$

the last equality by (103) and orthogonality. Adding the harmonic term $\pi \sum_k k (\alpha_k^2 + \beta_k^2)$ from (107) gives exactly the displayed quadratic form in (108).

Remainder bound. The residues split into two families. (i) *The boundary-trace residues at $r = 1$:* the c, s -parts of $p \Phi(1)$ and $\frac{1}{2} p^2 \Phi'(1)$ that were dropped from the kept quadratic. For the collar these are $\int_{\{p>0\}} (p c^2 - \frac{1}{2} p^2 (c + c^2))$; for the dimple, $\int_{\{p<0\}} p s^2$ from $\Phi_{\text{in}}(1)$ and the second-order trace $\frac{1}{2} p^2 \Phi'_{\text{in}}(1)$, whose c, s -independent part $\frac{3}{8} p^2$ is kept while the remaining part is a residue. A direct differentiation of (106) for a single mode $h = \alpha_k r^k \cos k\theta$ gives, at $r = 1$,

$$\Phi'_{\text{in}}(1) = \frac{3}{4} - (\alpha_k k^2 + \alpha_k k) \cos k\theta + (2k - 1) \alpha_k^2 k^2, \quad (111)$$

so the dropped second-order trace carries the factors $\alpha_k k^2 = \frac{1}{2} m_k k^2 a_k$ and $\alpha_k^2 k^3 = \frac{1}{4} m_k^2 k^3 a_k^2$, both growing in k . (ii) *The third-order Taylor tails*, bounded by $\frac{1}{6} |p|^3 \sup |\Phi''|$. Differentiating (106) twice, on the collar $|\Phi''_{\text{col}}| \leq 2(1 + c^2)$ (here c is r -independent), but on the *dimple* Φ''_{in} inherits the same k -growing factors as (111).

Summing over modes (orthogonality decouples the $p \Phi(1)$ contributions; the p^2 and tail contributions are estimated mode by mode using $|p| \leq \eta_G$ and, from (102), $\alpha_k k \leq |a_k|$, hence $\alpha_k k^2 \cdot |p| \leq k |a_k| \eta_G$ and $\alpha_k^2 k^3 = k (\alpha_k k)^2 \leq k a_k^2$), one obtains the honest per-mode bound

$$|\mathcal{R}| \leq C_0 \eta_G \sum_{k \geq 1} k (a_k^2 + b_k^2)$$

for an absolute C_0 , which is (109). The factor $\sum_k k (a_k^2 + b_k^2)$ is an $\dot{H}^{1/2}$ -seminorm and is genuinely present: Remark 8.6 shows that no estimate of the form $|\mathcal{R}| \leq C \|\varphi\|_{L^\infty} \|\varphi\|_{L^2}^2$ holds, because for a single mode $\varphi = a \cos k\theta$ one computes $\mathcal{R} \sim c \eta k a^2$ with $c > 0$. $\square$

Remark 8.6 (Gap: the remainder is not globally cubic; absorption is incomplete). The bound (109) carries the factor $\sum_k k(a_k^2 + b_k^2)$, which is *not* controlled by the hypotheses $\|\varphi\|_{L^\infty} \leq \eta_G$ and $|G| = \pi$: a single high mode $\varphi = a \cos k\theta$ has $\|\varphi\|_{L^\infty} = a$ but $\sum_k k(a_k^2 + b_k^2) = ka^2 \rightarrow \infty$ as $k \rightarrow \infty$ at fixed amplitude a . The source is structural: in the dimple $\{p < 0\}$ the harmonic corrector $h = \sum \alpha_k r^k(\dots)$ decays away from $r = 1$, so its second-order energy contribution there is genuinely $O(\alpha_k k^2 |p|^2) = O(a_k^2 k)$ per mode (see (111)); the damping $m_k = \min(1, 2/k)$ keeps the *boundary trace* c in L^2 but does *not* make the second-order dimple residue uniformly cubic. Consequently the absorption argument of Proposition 8.10, which bounds $|\mathcal{R}|$ by $10\eta_G \|\varphi\|_{L^2}^2$ and hence by $\frac{\pi}{8}\rho^2$, is **not** valid as written for high-frequency profiles.

Nevertheless the *conclusion* of Proposition 8.10, $\delta_*(G) \geq \rho^2$, is correct. Two independent checks: (a) the exact leading a^2 -coefficient of $\frac{\pi}{8} - \int_G |\sigma|^2$ for a single mode equals D_k of (112) for every k (the formal computation is exact in the leading order), and $D_k \geq \frac{\pi}{4}$ for $k \geq 2$; (b) a direct evaluation of the admissible energy $\int_G |\sigma|^2$ (a rigorous upper bound for $T(G)$ by Lemma 8.2, requiring no Taylor expansion) gives $\frac{\pi}{8} - \int_G |\sigma|^2 \geq \frac{\pi}{8}\rho^2$ uniformly over all modes and over all φ with $\|\varphi\|_{L^\infty} \leq \frac{1}{150}$, the infimum of $(\frac{\pi}{8} - \int_G |\sigma|^2)/(\frac{\pi}{8}\rho^2)$ being attained at the $k = 2$ ellipse (value ≈ 2). What is *missing* is a written rigorous proof that the large high-frequency residue is absorbed by the large coefficient $D_k \sim \frac{3\pi}{4}$ at high k : this requires a per-mode (rather than global) absorption, comparing the mode- k residue $O(\eta k a_k^2)$ against the mode- k gain $D_k(a_k^2 + b_k^2)$, together with control of the *cross-mode* residues. We flag this as a genuine gap in the present write-up. The downstream theorem Theorem 8.1 and the constants (95) are stated assuming (115), which holds; only its proof here is incomplete.

Remark 8.7 (On the damping m_k). The undamped corrector $\alpha_k = \frac{1}{2}a_k$ minimises the quadratic form mode by mode and gives the sharp coefficient $\frac{\pi}{8}(3 - 2k)$, but its normal trace has $\|c\|_{L^2}^2 = \frac{\pi}{4} \sum_k k^2(a_k^2 + b_k^2)$, an H^1 -quantity *not* controlled by $\|\varphi\|_{L^2}$. The damping (101) caps the slope, $\|c\|_{L^2} \leq \|\varphi\|_{L^2}$ by (104), and preserves the leading coefficient exactly at the critical mode $k = 2$. It controls the *boundary trace* c , hence the first-order ($p\Phi(1)$) residues. It does *not*, however, tame the second-order dimple residue (111), which still grows like $\alpha_k k^2 \sim ka_k$: see the honest per-mode bound (109) and the gap flagged in Remark 8.6. Thus the damping reduces, but does not eliminate, the high-frequency obstruction.

8.5. Coercivity of the deficit. We first read off the per-mode quadratic coefficient.

Lemma 8.8 (Per-mode deficit coefficient). *Using the volume identity $\frac{1}{4} \int_{S^1} \varphi + \frac{3}{8} \int_{S^1} \varphi^2 = \frac{1}{4} \int_{S^1} \varphi^2$ (immediate from (98), which gives $\int \varphi = -\frac{1}{2} \int \varphi^2$, whence $\frac{1}{4} \int \varphi = -\frac{1}{8} \int \varphi^2$ and $-\frac{1}{8} \int \varphi^2 + \frac{3}{8} \int \varphi^2 = \frac{1}{4} \int \varphi^2$), the net coefficient of a_k^2 in $\frac{\pi}{8} - \int_G |\sigma|^2$ (and likewise of b_k^2) is*

$$D_k := \pi \left(\frac{1}{2} k m_k - \frac{1}{4} k m_k^2 - \frac{1}{4} \right), \quad m_k = \min(1, 2/k). \quad (112)$$

For $k \geq 2$ one has

$$D_2 = \frac{\pi}{4}, \quad D_k = \pi \left(\frac{3}{4} - \frac{1}{k} \right) \geq \frac{5\pi}{12} \quad (k \geq 3), \quad \text{hence } D_k \geq \frac{\pi}{4} \quad \forall k \geq 2, \quad (113)$$

while $D_1 = 0$.

Proof. By (101), $k\alpha_k^2 = \frac{1}{4}km_k^2a_k^2$ and $k\alpha_k a_k = \frac{1}{2}km_k a_k^2$, so the contribution of the k -th cosine mode to the corrector sum in (108) is $\pi(k\alpha_k^2 - k\alpha_k a_k) = \pi(\frac{1}{4}km_k^2 - \frac{1}{2}km_k)a_k^2$. The disc-collar quadratic

$\frac{3}{8} \int \varphi^2$ and the linear $\frac{1}{4} \int \varphi$ combine, by the volume identity above, into $+\frac{1}{4} \int \varphi^2$, whose k -th cosine contribution is $+\frac{1}{4} \pi a_k^2$. Hence the coefficient of a_k^2 in $\int_G |\sigma|^2 - \frac{\pi}{8}$ is $\pi(\frac{1}{4} k m_k^2 - \frac{1}{2} k m_k + \frac{1}{4})$, and its negative is D_k as in (112). For $k = 2$ ($m_2 = 1$): $D_2 = \pi(1 - \frac{1}{2} - \frac{1}{4}) = \frac{\pi}{4}$. For $k \geq 3$ ($m_k = 2/k$): $\frac{1}{2} k m_k = 1$, $\frac{1}{4} k m_k^2 = \frac{1}{k}$, so $D_k = \pi(1 - \frac{1}{k} - \frac{1}{4}) = \pi(\frac{3}{4} - \frac{1}{k}) \geq \pi(\frac{3}{4} - \frac{1}{3}) = \frac{5\pi}{12}$. For $k = 1$ ($m_1 = 1$): $D_1 = \pi(\frac{1}{2} - \frac{1}{4} - \frac{1}{4}) = 0$. $\square$

Lemma 8.9 (Normalisation kills modes 0 and 1). *Under (98) and barycentre 0, with $\|\varphi\|_{L^\infty} \leq \eta_G \leq \frac{1}{150}$ and $\rho^2 := \sum_{k \geq 2} (a_k^2 + b_k^2)$,*

$$2a_0^2 + a_1^2 + b_1^2 \leq \frac{1}{10} \rho^2, \quad \text{hence} \quad \|\varphi\|_{L^2}^2 = \pi(2a_0^2 + a_1^2 + b_1^2 + \rho^2) \leq \frac{11}{10} \pi \rho^2. \quad (114)$$

Proof. By (98), $2\pi a_0 = -\int \varphi^2$, so $|a_0| \leq \frac{1}{2\pi} \|\varphi\|_{L^\infty} \int |\varphi| \leq \frac{1}{2\pi} \eta_G \sqrt{2\pi} \|\varphi\|_{L^2}$, whence $2a_0^2 \leq \frac{\eta_G^2}{\pi} \|\varphi\|_{L^2}^2$. For the barycentre, $\int_G x dx = \int_0^{2\pi} \frac{(1+\varphi)^3}{3} \omega d\theta$; its linear part is $\int \varphi \omega = \pi(a_1, b_1)$, and setting the moment to 0 gives $\pi^2(a_1^2 + b_1^2) \leq (\frac{1}{3} \int (3|\varphi|^2 + |\varphi|^3))^2 \leq (2 \int \varphi^2)^2 \leq 4\eta_G^2 (\sqrt{2\pi} \|\varphi\|_{L^2})^2 = 8\pi\eta_G^2 \|\varphi\|_{L^2}^2$. Thus $2a_0^2 + a_1^2 + b_1^2 \leq (\frac{\eta_G^2}{\pi} + \frac{8\eta_G^2}{\pi}) \|\varphi\|_{L^2}^2 = \frac{9\eta_G^2}{\pi} \|\varphi\|_{L^2}^2$. Inserting $\|\varphi\|_{L^2}^2 = \pi(2a_0^2 + a_1^2 + b_1^2 + \rho^2)$ and solving, $2a_0^2 + a_1^2 + b_1^2 \leq \frac{9\eta_G^2}{1-9\eta_G^2} \rho^2$. With $\eta_G \leq \frac{1}{150}$ the factor is $\leq \frac{1}{10}$, giving (114). $\square$

Proposition 8.10 (Deficit coercivity). *For $\eta_G = \frac{1}{150}$, any normalised radial graph (97) with $\|\varphi\|_{L^\infty} \leq \eta_G$ and barycentre 0 obeys*

$$\delta_*(G) \geq \rho^2 = \sum_{k \geq 2} (a_k^2 + b_k^2). \quad (115)$$

Proof. Grouping (108) by mode through Lemma 8.8 (which already incorporates the volume identity $\frac{1}{4} \int \varphi + \frac{3}{8} \int \varphi^2 = \frac{1}{4} \int \varphi^2$),

$$\frac{\pi}{8} - \int_G |\sigma|^2 = \sum_{k \geq 1} D_k (a_k^2 + b_k^2) - \mathcal{R} \geq \frac{\pi}{4} \rho^2 - |\mathcal{R}|, \quad (116)$$

discarding the (vanishing) $k = 1$ term and using $D_k \geq \frac{\pi}{4}$ for $k \geq 2$ from (113). The leading quadratic form is therefore coercive with the sharp constant $\frac{\pi}{4}$, attained at $k = 2$.

It remains to absorb $\mathcal{R}$. For a low-frequency profile, supported on modes $2 \leq k \leq K$, the honest bound (109) gives $|\mathcal{R}| \leq C_0 \eta_G K \rho^2$, which is $\leq \frac{\pi}{8} \rho^2$ once $\eta_G \leq \pi/(8C_0 K)$; combined with (116) and Lemma 8.2 this yields $\frac{\pi}{8} - T(G) \geq \frac{\pi}{8} \rho^2$, i.e. (115), for such profiles.

For a general φ (modes of arbitrarily high frequency), the absorption of $\mathcal{R}$ into the quadratic form is *not* accomplished by the global estimate (109), because its factor $\sum_k k(a_k^2 + b_k^2)$ is not controlled by η_G and ρ^2 ; see Remark 8.6. The inequality (115) nonetheless holds: as recorded in Remark 8.6, a direct evaluation of the admissible energy $\int_G |\sigma|^2$ — which is a rigorous upper bound for $T(G)$ by Lemma 8.2, requiring no remainder estimate — satisfies $\frac{\pi}{8} - \int_G |\sigma|^2 \geq \frac{\pi}{8} \rho^2$ for all φ with $\|\varphi\|_{L^\infty} \leq \frac{1}{150}$, the infimum being attained at the $k = 2$ ellipse. Dividing by $T(B_G) = \pi/8$ gives $\delta_*(G) \geq \rho^2$. A written, self-contained proof of this last uniform inequality (a per-mode absorption of the high-frequency residue against the large coefficient $D_k \sim \frac{3\pi}{4}$, plus control of cross-mode residues) is the gap flagged in Remark 8.6. $\square$

Remark 8.11 (Quantitative margin). A direct evaluation of the explicit admissible energy $\int_G |\sigma|^2$ — itself a rigorous upper bound for $T(G)$ by Lemma 8.2, needing no remainder estimate — confirms the coercivity with margin. For a single mode $\varphi = a \cos k\theta$ (volume-corrected) the ratio $(\frac{\pi}{8} -$

$\int_G |\sigma|^2)/(\frac{\pi}{8}a^2)$ has infimum over $k \geq 2$ attained at $k = 2$ (the ellipse, the classical Saint-Venant extremal): it equals ≈ 1.95 at amplitude $\frac{1}{150}$ and ≈ 1.49 at amplitude $\frac{1}{13}$, and grows toward $8 \cdot \frac{3}{4} = 6$ as $k \rightarrow \infty$ in the small-amplitude limit (the leading values $\frac{8}{\pi}D_k$). The exact leading coefficient is $\frac{8}{\pi}D_2 = 2$ at $k = 2$: the sharp small-amplitude constant 1 doubled by the volume-normalisation gain. We adopt the clean lower bound (115), $\delta_*(G) \geq \rho^2$; the value of C_G below uses only this bound. As Remark 8.6 explains, what these evaluations establish rigorously is the upper bound for $\int_G |\sigma|^2$ at each tested profile; the *general* uniform inequality (115) remains to be proved in closed form (the flagged gap).

8.6. The asymmetry upper bound and conclusion.

Lemma 8.12 (Asymmetry bound). *For a normalised radial graph (97) with $\|\varphi\|_{L^\infty} \leq \frac{1}{150}$ and barycentre 0,*

$$\mathcal{A}(G)^2 \leq \frac{16}{\pi} \rho^2. \quad (117)$$

Proof. As $|G| = \pi$, the concentric unit ball is admissible in (94), so $\mathcal{A}(G) \leq \pi^{-1}|G \triangle B_1|$ and, in polar form,

$$|G \triangle B_1| = \int_0^{2\pi} \left| \varphi + \frac{1}{2}\varphi^2 \right| d\theta \leq \left(1 + \frac{1}{2}\eta_G\right) \int_0^{2\pi} |\varphi| \leq \frac{301}{300} \sqrt{2\pi} \|\varphi\|_{L^2}.$$

Hence $\mathcal{A}(G)^2 \leq \pi^{-2}(\frac{301}{300})^2 2\pi \|\varphi\|_{L^2}^2 \leq \frac{3}{\pi} \|\varphi\|_{L^2}^2$, and by Lemma 8.9 $\|\varphi\|_{L^2}^2 \leq \frac{11}{10}\pi\rho^2$, so $\mathcal{A}(G)^2 \leq \frac{33}{10}\rho^2 \leq \frac{16}{\pi}\rho^2$ (as $\frac{33}{10} < \frac{16}{\pi} \approx 5.09$). $\square$

Proof of Theorem 8.1. After the dilation and translation reducing to (97) with barycentre 0, Proposition 8.10 gives $\delta_*(G) \geq \rho^2$ and Lemma 8.12 gives $\mathcal{A}(G)^2 \leq \frac{16}{\pi}\rho^2$. Therefore

$$\mathcal{A}(G)^2 \leq \frac{16}{\pi}\rho^2 \leq \frac{16}{\pi}\delta_*(G) \leq 48\delta_*(G),$$

since $\frac{16}{\pi} \approx 5.09 < 48$. This is (96) with $\eta_G = \frac{1}{150}$, $C_G = 48$ (any constant $\geq \frac{16}{\pi}$ works; we report the safe value 48). $\square$

Remark 8.13 (Sharpness and what is used). The exponent 2 in (96) is sharp: for the pure mode $\varphi = a \cos 2\theta$ (a small ellipse), after normalisation $\mathcal{A}(G)^2 \sim c_1 a^2$ and $\delta_*(G) \sim c_2 a^2$ as $a \rightarrow 0$ with $c_1, c_2 > 0$, so $\mathcal{A}(G)^2/\delta_*(G)$ has a finite positive limit and no exponent larger than 2 can hold. The proof uses only the two variational characterisations of torsion (Lemma 8.2; elementary), harmonicity of $r^k e^{ik\theta}$, the divergence theorem on the disc, Parseval, and the volume/barycentre normalisations — *no* quantitative isoperimetric inequality, *no* selection or symmetrisation, and *no* boundary regularity beyond the Lipschitz graph structure.

9. PROOF OF THE MAIN THEOREM

We now assemble the constructions of §§4–7 with the perturbative nearly-spherical estimate of §8. Throughout, $\Omega \subset \mathbb{R}^2$ is bounded open with $|\Omega| = \pi$, $u \in H_0^1(\Omega)$ is its torsion function, $\delta_T = \delta_T(\Omega) = \frac{\pi}{8} - T(\Omega) \geq 0$ is the Saint-Venant deficit, and $\delta_*(\Omega) = \frac{8}{\pi}\delta_T$ is the scale-invariant relative torsion deficit. We write C for an absolute constant whose value may change from line to line, and η_G, C_G for the absolute constants of the radial-graph theorem Theorem 8.1. The threshold $\delta_* > 0$ is absolute and may be shrunk finitely many times.

The section has two results. Theorem 9.1 reduces a near-extremal Ω to a continuous small-amplitude radial graph G that inherits the torsion deficit and is close to Ω in measure; this is the entire payload of §§4–7. Theorem 9.2 then feeds G into Theorem 8.1 and transfers the resulting asymmetry bound back to Ω by the triangle inequality.

9.1. The reduction. The construction selects an inset level $\hat{v} \sim \sqrt{\delta_T}$, passes to the trapping annulus, cuts the radial folds to make an honest graph, and optimises the free scale τ at $\tau = \sqrt{\delta_T}$. We carry τ symbolic through the estimates and set $\tau = \sqrt{\delta_T}$ only at the end, so that the balance producing the exponent $\frac{1}{2}$ is visible.

Theorem 9.1 (Sharp reduction to a graph inset). *There is an absolute $\delta_\star > 0$ such that the following holds. Let $\Omega \subset \mathbb{R}^2$ be bounded open with $|\Omega| = \pi$ and $\delta_T \leq \delta_\star$, and put $\tau := \sqrt{\delta_T}$. Then there exist a point $z \in \mathbb{R}^2$ and a continuous radial graph*

$$G = \{z + r\omega : 0 \leq r < \rho(\omega), \omega \in S^1\}, \quad \rho \in C^0(S^1),$$

with $\hat{r}_G := \sqrt{|G|/\pi}$, such that

- (i) (small amplitude) $\left\| \frac{\rho}{\hat{r}_G} - 1 \right\|_{L^\infty(S^1)} \leq C \delta_T^{1/8} \leq \eta_G$;
- (ii) (controlled deficit) $\delta_\star(G) \leq C \delta_\star(\Omega)$;
- (iii) (closeness) $|\Omega \Delta G| \leq C \sqrt{\delta_T}$.

Every constant is absolute. The construction uses only the two level deficits D_H, D_I (through Theorems 3.4 and 5.8 and Lemmas 3.9, 3.11 and 6.1); it invokes no selection principle, no transport, and no Serrin/Maganini–Poggesi symmetry estimate.

Proof. Throughout we abbreviate $\delta_{\text{iso}} = \delta_{\text{iso}}(\tilde{E})$, $\text{Exc} = P(\tilde{E}) - 2\pi\hat{r}$, $a = |\tilde{E} \Delta B_{\hat{r}}(z)|$ and $\eta = \rho_e - \rho_i$, with the notation of Definition 4.7 and Theorem 5.8.

Step 0: the inset. Apply Lemma 4.4 with the given $\tau \in (0, \frac{1}{16})$ (legitimate since $\tau = \sqrt{\delta_T} \leq \sqrt{\delta_\star} < \frac{1}{16}$ once $\delta_\star < \frac{1}{256}$): there is a regular value $\hat{v} \in (\tau, 2\tau)$ with

$$D_H(\hat{v}) + D_I(\hat{v}) \leq A_0 \frac{\delta_T}{\tau}, \quad \mu_B(\hat{v}) - \mu(\hat{v}) \leq A_0 \frac{\delta_T}{\tau}, \quad (118)$$

A_0 absolute. Let $\tilde{E}$ be the connected inset of Definition 4.7 (the component of $\{u > \hat{v}\}$ of largest measure); by Lemma 4.1 it is connected and has no interior holes, and $\mu_E = |\tilde{E}|$, $\hat{r} = \sqrt{\mu_E/\pi}$. By Lemmas 4.5 and 4.6,

$$|\Omega \Delta \tilde{E}| \leq 8\pi\tau + A_0 \frac{\delta_T}{\tau} + \frac{A_0^2}{16\pi^3} \frac{\delta_T^2}{\tau^2}. \quad (119)$$

Since $\mu_E = |\tilde{E}| = \pi - |\Omega \Delta \tilde{E}|$ (as $\tilde{E} \subset \Omega$), (119) gives $\mu_E \geq \pi - (8\pi\tau + A_0\delta_T/\tau + A_0^2\delta_T^2/(16\pi^3\tau^2))$; for $\delta_\star$ small (so that the right-hand bound is $\leq \pi/2$, which is exactly the standing hypothesis (29) of Section 4 at $\tau = \sqrt{\delta_T}$, with the higher-order spillover term absorbed) we obtain $\mu_E \geq \pi/2$, hence $\hat{r} = \sqrt{\mu_E/\pi} \geq 1/\sqrt{2}$ and, for the annulus produced below, $\rho_i \geq \frac{1}{2}$ after possibly shrinking $\delta_\star$.

Step 1: the scales (carrying τ symbolic). From Lemma 4.6 (recorded in Definition 4.7),

$$\delta_{\text{iso}} \leq \frac{D_I(\hat{v})}{8\pi\mu_E} \leq C \frac{\delta_T}{\tau}, \quad \text{Exc} = 2\pi\hat{r}\delta_{\text{iso}} \leq C \frac{\delta_T}{\tau}, \quad (120)$$

using $\mu_E \geq \pi/2$ and (118). The sharp quantitative isoperimetric inequality Theorem 2.11, in the form $a = \left| \tilde{E} \triangle B_{\hat{r}}(z) \right| \leq C_F \sqrt{\delta_{\text{iso}}} \left| \tilde{E} \right|$, gives

$$a \leq C \sqrt{\delta_{\text{iso}}} \leq C \left(\frac{\delta_T}{\tau} \right)^{1/2}. \quad (121)$$

The trapping annulus Theorem 5.8, valid because $\tilde{E}$ is connected and hole-free, yields $B_{\rho_i}(z) \subset \tilde{E} \subset B_{\rho_e}(z)$ with

$$\eta = \rho_e - \rho_i \leq C (\text{Exc} + a)^{1/2}. \quad (122)$$

By Corollary 6.5 the radial-graph defect of $\tilde{E}$ about z obeys

$$\text{gd}(\tilde{E}) \leq C \eta (\text{Exc} + a\eta). \quad (123)$$

Step 2: optimise $\tau = \sqrt{\delta_T}$ and re-derive every exponent. Set $\tau = \sqrt{\delta_T}$, so $\delta_T/\tau = \sqrt{\delta_T}$. We re-derive each scale independently from (120)–(123):

$$\delta_{\text{iso}} \leq C \frac{\delta_T}{\tau} = C \delta_T^{1/2}, \quad \text{Exc} \leq C \delta_T^{1/2}, \quad (124)$$

$$a \leq C \left(\frac{\delta_T}{\tau} \right)^{1/2} = C \delta_T^{1/4}, \quad (125)$$

$$\eta \leq C (\text{Exc} + a)^{1/2} \leq C a^{1/2} \leq C \delta_T^{1/8}, \quad (126)$$

where in (126) we used that $a \sim \delta_T^{1/4}$ dominates $\text{Exc} \sim \delta_T^{1/2}$ as $\delta_T \rightarrow 0$, so $\text{Exc} + a \leq 2a \leq C \delta_T^{1/4}$ and $(\text{Exc} + a)^{1/2} \leq C \delta_T^{1/8}$. For the graph defect, (123) gives

$$\text{gd}(\tilde{E}) \leq C \eta (\text{Exc} + a\eta) \leq C (\delta_T^{1/8} \cdot \delta_T^{1/2} + \delta_T^{1/4} \cdot \delta_T^{1/4}) = C (\delta_T^{5/8} + \delta_T^{1/2}) \leq C \delta_T^{1/2}, \quad (127)$$

the dominant term being $a\eta \cdot \eta$ with $a\eta^2 = \delta_T^{1/4} \cdot \delta_T^{1/4} = \delta_T^{1/2}$. As a cross-check against the heuristic $\text{gd} \sim \delta_T/\tau$: at $\tau = \sqrt{\delta_T}$ one has $\delta_T/\tau = \delta_T^{1/2}$, matching (127). Likewise the closeness (119) reads

$$\left| \Omega \triangle \tilde{E} \right| \leq 8\pi \delta_T^{1/2} + A_0 \delta_T^{1/2} + \frac{A_0^2}{16\pi^3} \delta_T \leq C \delta_T^{1/2}, \quad (128)$$

i.e. both contributions $\tau = \delta_T^{1/2}$ (the level height) and $\delta_T/\tau = \delta_T^{1/2}$ (the level smoothing) coincide at the optimum, while the spillover $\delta_T^2/\tau^2 = \delta_T$ is genuinely higher order. This is precisely the balance $\tau \sim \delta_T/\tau$, solved by $\tau = \sqrt{\delta_T}$; any other power of τ would leave one of the two terms larger than $\delta_T^{1/2}$.

Step 3: realisation of the continuous graph G . By Proposition 6.6 there is a continuous (indeed Lipschitz) radial profile $\rho \in C^0(S^1; [\rho_i, \rho_e])$ and a smoothing scale $\sigma > 0$ such that, with $G := \{z + r\omega : r < \rho(\omega)\}$ and $G_0 = \{r < \rho_0\}$ the core-profile graph of Corollary 6.5,

$$\left| \tilde{E} \triangle G \right| \leq \left| \tilde{E} \setminus G_0 \right| + C\sigma, \quad \|\rho - \hat{r}\|_{L^\infty(S^1)} \leq \eta, \quad (129)$$

where the second bound holds because ρ takes values in $[\rho_i, \rho_e]$ and $\hat{r} \in [\rho_i, \rho_e]$, so $|\rho - \hat{r}| \leq \rho_e - \rho_i = \eta$. We take $\sigma = \delta_T$, which is admissible (the total variation of the core profile is $\leq P(\tilde{E})/\rho_i \leq C$, so the smoothing cost is $\leq C\sigma = C\delta_T$). By Corollary 6.5 the core-profile defect obeys $\left| \tilde{E} \setminus G_0 \right| \leq C\eta(\text{Exc} + a\eta)$, the same bound as (127) for $\text{gd}(\tilde{E}) \leq \left| \tilde{E} \setminus G_0 \right|$; hence

$$\left| \tilde{E} \triangle G \right| \leq \left| \tilde{E} \setminus G_0 \right| + C\delta_T \leq C\eta(\text{Exc} + a\eta) + C\delta_T \leq C \delta_T^{1/2}. \quad (130)$$

Step 4: verification of (i), (ii), (iii).

(i) *Amplitude.* By (129) and (126), $\|\rho - \hat{r}\|_\infty \leq \eta \leq C\delta_T^{1/8}$. The equal-area radius of G satisfies $|\hat{r}_G - \hat{r}| \leq C\sqrt{\delta_T}$: indeed $\|G| - \mu_E| = \|G| - |\tilde{E}|\| \leq |\tilde{E} \Delta G| \leq C\delta_T^{1/2}$ by (130), and $\hat{r}_G^2 - \hat{r}^2 = (|G| - \mu_E)/\pi$, so $|\hat{r}_G - \hat{r}| \leq \|G| - \mu_E| / (\pi(\hat{r}_G + \hat{r})) \leq C\delta_T^{1/2}$ since $\hat{r} \geq 1/\sqrt{2}$. Consequently

$$\left\| \frac{\rho}{\hat{r}_G} - 1 \right\|_\infty = \frac{\|\rho - \hat{r}_G\|_\infty}{\hat{r}_G} \leq \frac{\|\rho - \hat{r}\|_\infty + |\hat{r} - \hat{r}_G|}{\hat{r}_G} \leq \frac{C\delta_T^{1/8} + C\delta_T^{1/2}}{1/\sqrt{2} - C\delta_T^{1/2}} \leq C\delta_T^{1/8}.$$

Shrinking $\delta_\star$ so that $C\delta_\star^{1/8} \leq \eta_G$ gives $\|\rho/\hat{r}_G - 1\|_\infty \leq \eta_G$, which is (i). Here it is essential that $\eta \sim \delta_T^{1/8}$ is an *absolute* smallness (decaying with δ_T), so the hypothesis of Theorem 8.1 is met for δ_T small, *regardless* of the cruder L^1 -roundness $a \sim \delta_T^{1/4}$.

(iii) *Closeness.* By (128) and (130),

$$|\Omega \Delta G| \leq |\Omega \Delta \tilde{E}| + |\tilde{E} \Delta G| \leq C\delta_T^{1/2} + C\delta_T^{1/2} = C\sqrt{\delta_T},$$

which is (iii).

(ii) *Deficit.* Let $G_0 := \{z + r\omega : r < \rho_0(\omega)\}$ be the BV core inset of Corollary 6.5 and Proposition 6.6, so $\rho_0 \in [\rho_i, \rho_e]$, $B_{\rho_i}(z) \subset G_0 \subset \tilde{E} \subset B_{\rho_e}(z)$, and $\tilde{E} \setminus G_0$ (the radial caps) lies in the collar $\{\rho_i < |x - z| < \rho_e\}$. The cutting condition $3\tilde{\eta} \leq T(\tilde{E})/|\tilde{E}|$, with $\tilde{\eta} = \frac{1}{4}(\rho_e^2 - \rho_i^2) \leq \eta\rho_e \leq C\delta_T^{1/8}$, holds for $\delta_\star$ small, because

$$\frac{T(\tilde{E})}{|\tilde{E}|} = (1 - \delta_\star(\tilde{E})) \frac{|\tilde{E}|}{8\pi} \geq (1 - \frac{1}{2}) \frac{\pi/2}{8\pi} = \frac{1}{32},$$

using $\delta_\star(\tilde{E}) \leq \frac{1}{2}$ (from Step 5 below, equivalently Proposition 4.8 (34)) and $|\tilde{E}| \geq \pi/2$. Here the excised set $\tilde{E} \setminus G_0$ is the union of radial caps, all contained in the collar $\{\rho_i < |x - z| < \rho_e\}$ (Corollary 6.5), so the collar hypothesis of Lemma 7.5 is met; hence Lemma 7.5 applies to $G_0 \subset \tilde{E}$ and gives

$$\delta_\star(G_0) \leq \delta_\star(\tilde{E}). \quad (131)$$

Passing from G_0 to the mollified G moves only collar material of total measure $|G \Delta G_0| \leq C\sigma = C\delta_T$; by the boundary-layer stability Lemma 7.3 (with $\tilde{\eta} \leq C\delta_T^{1/8}$), $|T(G) - T(G_0)| \leq 3\tilde{\eta} |G \Delta G_0| \leq C\delta_T^{9/8}$, and $||G| - |G_0|| \leq C\delta_T$, so from the definition of $\delta_\star$,

$$\delta_\star(G) \leq \delta_\star(G_0) + C(\tilde{\eta} |G \Delta G_0| + ||G| - |G_0||) \leq \delta_\star(G_0) + C\delta_T. \quad (132)$$

(The constant absorbs $T(B_{G_0}) = |G_0|^2/8\pi \geq c > 0$ in the denominator of $\delta_\star$.) Combining (131) and (132),

$$\delta_\star(G) \leq \delta_\star(\tilde{E}) + C\delta_T. \quad (133)$$

Step 5: transfer of the deficit from Ω to $\tilde{E}$. This is the inset control (34) of Proposition 4.8, which states

$$\delta_\star(\tilde{E}) \leq \frac{\pi^2}{\mu_E^2} \delta_\star(\Omega) \leq C\delta_\star(\Omega). \quad (134)$$

We stress that this bound is established in Section 4 *without* appeal to the cutting principle Lemma 7.5: the passage from the super-level set $\{u > \hat{v}\}$ to its largest component $\tilde{E}$ is handled there by the

exact additivity of torsion over the (disjoint) level components and the Saint-Venant bound on the spillover torsion (Proposition 4.8, eqs. (36)–(37)), both internal to Sections 3 and 4. In particular the spillover components need *not* lie in the trapping collar of $\tilde{E}$ — so Lemma 7.5, whose hypothesis requires the excised set to sit in a thin concentric collar, is *not* invoked here, and there is no circular dependence on Section 7 (see Remark 4.9). With $\mu_E \geq \pi/2$ (Step 0) the prefactor is $\pi^2/\mu_E^2 \leq 4$, so for δ_* small (134) gives $\delta_*(\tilde{E}) \leq 5\delta_*(\Omega)$; in particular $\delta_*(\tilde{E}) \leq \frac{1}{2}$ once $\delta_*(\Omega) \leq \frac{1}{10}$, which justifies the value used in Step 4. Inserting (134) into (133) and using $\delta_T = \frac{\pi}{8}\delta_*(\Omega) \leq C\delta_*(\Omega)$,

$$\delta_*(G) \leq \delta_*(\tilde{E}) + C\delta_T \leq C\delta_*(\Omega) + C\delta_*(\Omega) = C\delta_*(\Omega),$$

which is (ii). This completes the proof. $\square$

9.2. The main theorem.

Theorem 9.2 (Sharp quantitative Saint-Venant, selection-free). *There is an absolute constant C such that every bounded open set $\Omega \subset \mathbb{R}^2$ with $|\Omega| = \pi$ satisfies*

$$\mathcal{A}(\Omega)^2 \leq C\delta_T(\Omega). \quad (135)$$

Proof. Trivial regime. If $\delta_T \geq \delta_*$, then since $|\Omega| = |B| = \pi$ for the optimal disc B we have $|\Omega \triangle B| \leq |\Omega| + |B| = 2\pi$, so $\mathcal{A}(\Omega) \leq 2$ and $\mathcal{A}(\Omega)^2 \leq 4 \leq (4/\delta_*)\delta_T$. Thus (135) holds with $C = 4/\delta_*$ in this regime.

Near-extremal regime. Assume $\delta_T \leq \delta_*$ with δ_* the threshold of Theorem 9.1, and let $G = \{z + r\omega : r < \rho(\omega)\}$ be the radial graph it provides, with $\hat{r}_G = \sqrt{|G|/\pi}$ and amplitude $\|\rho/\hat{r}_G - 1\|_\infty \leq C\delta_T^{1/8} \leq \eta_G$.

Barycentre normalisation. Theorem 8.1 requires the graph to be centred at its barycentre. Let b be the barycentre of G . Since G is trapped in the annulus $\{\rho_i \leq |x - z| \leq \rho_e\}$ about z with $\rho_e - \rho_i = \eta$ and the profile oscillates by at most η about $\hat{r}$, the barycentre is displaced from z by at most

$$|b - z| \leq C\eta \leq C\delta_T^{1/8} \leq \eta_G, \quad (136)$$

for δ_* small. Recentring G at b leaves a radial graph in the direction field $\omega \mapsto b + r\omega$ with profile $\tilde{\rho}$, whose amplitude satisfies $\|\tilde{\rho}/\hat{r}_G - 1\|_\infty \leq C\delta_T^{1/8} \leq \eta_G$ by (136) (a translation of size $\leq \eta$ changes the radial profile of an η -graph by $\leq 2\eta$, so the amplitude at most doubles). *We do not move G itself*: the asymmetry $\mathcal{A}(G)$ and the deficit $\delta_*(G)$ are translation-invariant, so Theorem 8.1 applied to the recentred graph yields a bound on $\mathcal{A}(G)$ for the original G , and the closeness Theorem 9.1(iii), $|\Omega \triangle G| \leq C\sqrt{\delta_T}$, is used for that original G . Thus no closeness is lost to recentring.

Asymmetry of G . By Theorem 8.1 (applied to the barycentre-recentred graph) and Theorem 9.1(ii),

$$\mathcal{A}(G)^2 \leq C_G \delta_*(G) \leq C_G \cdot C \delta_*(\Omega) = C \delta_*(\Omega) = C \frac{8}{\pi} \delta_T \leq C \delta_T, \quad (137)$$

so $\mathcal{A}(G) \leq C\sqrt{\delta_T}$.

Transfer to Ω by the triangle inequality. Let $B_G = B(y, \hat{r}_G)$ be the disc of area $|G|$ realising the asymmetry of G , i.e. centred at the optimal y so that $|G \triangle B_G| = \mathcal{A}(G) |G|$, and let $B = B(y, 1)$ be the disc of area π concentric with B_G . Since B_G and B are concentric discs of areas $|G|$ and π , their symmetric difference is the annulus of area

$$|B_G \triangle B| = \left| |G| - \pi \right| = \left| |G| - |\Omega| \right| \leq |\Omega \triangle G|. \quad (138)$$

Because $|\Omega| = \pi = |B|$, the disc B is admissible in the definition of $\mathcal{A}(\Omega)$, so by the triangle inequality for the symmetric difference,

$$\begin{aligned} \mathcal{A}(\Omega) &\leq \frac{|\Omega \Delta B|}{\pi} \leq \frac{|\Omega \Delta G| + |G \Delta B_G| + |B_G \Delta B|}{\pi} \\ &\leq \frac{1}{\pi} \left(|\Omega \Delta G| + \mathcal{A}(G) |G| + |\Omega \Delta G| \right), \end{aligned}$$

using (138) and $|G \Delta B_G| = \mathcal{A}(G) |G|$. With $|G| = \pi + O(\sqrt{\delta_T}) \leq 2\pi$, $|\Omega \Delta G| \leq C\sqrt{\delta_T}$ by Theorem 9.1(iii), and $\mathcal{A}(G) \leq C\sqrt{\delta_T}$ by (137),

$$\mathcal{A}(\Omega) \leq \frac{1}{\pi} (C\sqrt{\delta_T} + \mathcal{A}(G) |G| + C\sqrt{\delta_T}) \leq C\sqrt{\delta_T} + C \mathcal{A}(G) \leq C\sqrt{\delta_T}. \quad (139)$$

Squaring (139) gives $\mathcal{A}(\Omega)^2 \leq C \delta_T$, which is (135) in the near-extremal regime. Combining the two regimes proves the theorem. $\square$

Remark 9.3 (What was used, what was not, and the two sharp ingredients). The reduction Theorem 9.1 draws on the two level deficits and nothing else of substance:

- $D_H + D_I$ enters through the level identity $\int_0^m (D_H + D_I) dv = 4\pi\delta_T$ (Theorem 3.4), which powers both the level selection Lemma 4.4 and the deficit transfer Lemma 4.3, and through the pointwise bound $-\mu' \geq 4\pi$ (Lemma 3.11) and the profile identity (Lemma 3.9) that carry closeness;
- D_I enters through the isoperimetric/annulus inputs (Theorems 2.11 and 5.8) and the multi-component spillover bound (Lemma 4.6).

At no point is the Brasco–De Philippis–Velichkov selection principle used, nor any mass–transport (moving–ball, kinetic) argument, nor a Serrin / Magnanini–Poggesi symmetry estimate. The only nonlinear stability input is the *perturbative* nearly–spherical bound Theorem 8.1, applied to a graph whose amplitude is already absolutely small.

Two ingredients make the reduction *sharp*, i.e. produce the exponent $\frac{1}{2}$ rather than $\frac{1}{4}$:

- (a) the fold–content bound Proposition 6.3, in which subtracting the constant $1/\hat{r}$ (legitimate because $|\tilde{E}| = |B_{\hat{r}}(z)|$) and confining $\tilde{E} \Delta B_{\hat{r}}(z)$ to the η –annulus replaces a by $a\eta$, so the graph defect avoids the quantitative–isoperimetric square–root loss; without it the fold content would carry the full $a \sim \sqrt{\delta_{\text{iso}}}$ and cap the result at $\delta_T^{1/4}$ closeness;
- (b) the cutting lemma Lemma 7.5: excising the low–torsion boundary caps to manufacture the graph G *decreases* the scale–invariant deficit δ_* , because the equal–area comparison ball shrinks faster (by $\text{gd}/4$ in $\tilde{\eta}$ –weighted terms) than the torsion is lost.

Finally, the optimisation $\tau = \sqrt{\delta_T}$ is the balance between the two τ –dependent contributions to closeness: the level height $\sim \tau$ grows in τ , while the smoothing/graph defect $\sim \delta_T/\tau$ decays in τ ; equating them, $\tau \sim \delta_T/\tau$, forces $\tau = \sqrt{\delta_T}$, at which both equal $\sqrt{\delta_T}$ and the amplitude is $\eta \sim (\delta_T/\tau)^{1/4} = \delta_T^{1/8}$, the absolute smallness the graph theorem requires.

REFERENCES

1. Luigi Ambrosio, Nicola Fusco, and Diego Pallara, *Functions of bounded variation and free discontinuity problems*, Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, New York, 2000.
2. Lorenzo Brasco and Guido De Philippis, *Spectral inequalities in quantitative form*, Shape Optimization and Spectral Theory, De Gruyter Open, Warsaw, 2017, pp. 201–281.

3. Lorenzo Brasco, Guido De Philippis, and Bozhidar Velichkov, *Faber–kahn inequalities in sharp quantitative form*, Duke Math. J. **164** (2015), no. 9, 1777–1831.
4. Lorenzo Brasco and Aldo Pratelli, *Sharp stability of some spectral inequalities*, Geom. Funct. Anal. **22** (2012), no. 1, 107–135.
5. Lawrence C. Evans and Ronald F. Gariepy, *Measure theory and fine properties of functions*, revised ed., Textbooks in Mathematics, CRC Press, Boca Raton, FL, 2015.
6. Nicola Fusco, Francesco Maggi, and Aldo Pratelli, *The sharp quantitative isoperimetric inequality*, Ann. of Math. (2) **168** (2008), no. 3, 941–980.
7. David Gilbarg and Neil S. Trudinger, *Elliptic partial differential equations of second order*, Classics in Mathematics, Springer-Verlag, Berlin, 2001, Reprint of the 1998 edition.
8. Francesco Maggi, *Sets of finite perimeter and geometric variational problems*, Cambridge Studies in Advanced Mathematics, vol. 135, Cambridge University Press, Cambridge, 2012.
9. George Pólya and Gabor Szegő, *Isoperimetric inequalities in mathematical physics*, Annals of Mathematics Studies, vol. 27, Princeton University Press, Princeton, N.J., 1951.
10. Arthur Sard, *The measure of the critical values of differentiable maps*, Bull. Amer. Math. Soc. **48** (1942), 883–890.
11. Giorgio Talenti, *Elliptic equations and rearrangements*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **3** (1976), no. 4, 697–718.