

THE SHARP QUANTITATIVE FABER–KRAHN INEQUALITY VIA THE LEVEL-SET MOVING-BALL ROUTE

ABSTRACT. We prove the sharp quantitative Faber–Krahn inequality: there is a dimensional constant $c_{\text{FK}}(n) > 0$ such that every open set $\Omega \subset \mathbb{R}^n$ with $0 < |\Omega| < \infty$ satisfies

$$\left(\frac{|\Omega|}{\omega_n}\right)^{2/n} (\lambda_1(\Omega) - \lambda_1(B^*)) \geq c_{\text{FK}}(n) \mathcal{A}(\Omega)^2,$$

where λ_1 is the first Dirichlet eigenvalue, B^* the ball of the same volume, and $\mathcal{A}$ the Fraenkel asymmetry; the exponent 2 is sharp. The proof is organised around the torsion function and the unscaled moving-ball distance $q(\rho) = \inf_z |E_\rho \Delta B_\rho(z)|$ on its superlevel sets E_ρ . On reduced boundaries and for almost every level we establish an exact level-set deficit identity expressing the Saint–Venant deficit as a weighted integral of a Cauchy–Schwarz defect D_H and an isoperimetric defect D_I ; nestedness gives a perimeter-free Lipschitz bound for q ; a finite-perimeter affine first variation for q , a positive kinetic estimate, and an endpoint trace yield bounded sharp Saint–Venant stability $\mathcal{A}(\Omega)^2 \leq C(n, R) \delta_T(\Omega)$ for $\Omega \subset B_R$. A constructive De Giorgi truncation removes the confinement, and the Kohler–Jobin rearrangement transfers the Saint–Venant estimate to the eigenvalue. Apart from the Fusco–Julin strong quantitative isoperimetric inequality and standard tools for sets of finite perimeter, the argument – including the bounded reduction and the Kohler–Jobin inequality – is self-contained.

CONTENTS

1. Introduction	2
2. Standing notation and scope	3
3. Finite-perimeter level-set identities	4
Regularity and level conventions	4
Coarea form of $-m'$ and the flux identity	5
The rearranged primitive	5
The exact deficit identity	6
4. Profile gap and bad-density estimate	7
5. Quantitative isoperimetry and same-centre trace control	8
6. Coarea differentiation and shell-admissible radii	12
Shell-admissible radii	12
7. Affine moving-ball first variation	14
8. Positive kinetic estimate	16
9. Endpoint trace and bounded stability	17
10. Bounded reduction	18
10.1. Standing notation and the standard tools we cite	19
10.2. The fast-convergence iteration lemma	20
10.3. The De Giorgi L^∞ tail estimate	21
10.4. A suboptimal stability seed	23
10.5. The truncation construction	25
10.6. The far branch	29
10.7. The near branch	29

10.8. Proof of the global theorem	30
11. The Kohler–Jobin inequality and the transfer to Faber–Krahn	31
11.1. Notation and scaling conventions	31
11.2. Part 1: the Kohler–Jobin inequality, proved from scratch	31
11.3. Part 2: transfer to sharp Faber–Krahn stability	36
12. Proof architecture and constants	38
Appendix A. Standard tools from the theory of sets of finite perimeter	39
Appendix B. Dependency certificate	39
References	40

1. INTRODUCTION

For an open set $\Omega \subset \mathbb{R}^n$ ($n \geq 2$) with $0 < |\Omega| < \infty$, let $\lambda_1(\Omega)$ denote its first Dirichlet eigenvalue and B^* the ball with $|B^*| = |\Omega|$. The Faber–Krahn inequality asserts $\lambda_1(\Omega) \geq \lambda_1(B^*)$, with equality only for balls. This paper proves its sharp quantitative form.

Theorem 1.1 (Sharp quantitative Faber–Krahn). *There is a constant $c_{\text{FK}}(n) > 0$, depending only on n , such that every open set $\Omega \subset \mathbb{R}^n$ with $0 < |\Omega| < \infty$ satisfies*

$$\delta_{\text{FK}}(\Omega) := \left(\frac{|\Omega|}{\omega_n} \right)^{2/n} (\lambda_1(\Omega) - \lambda_1(B^*)) \geq c_{\text{FK}}(n) \mathcal{A}(\Omega)^2,$$

where $\mathcal{A}(\Omega) := \inf_{x_0} |\Omega \Delta (x_0 + B^*)| / |B^*|$ is the Fraenkel asymmetry. The exponent 2 is sharp.

This is the theorem of Brasco–De Philippis–Velichkov [3]; the present proof is structurally different, and self-contained apart from the Fusco–Julin quantitative isoperimetric inequality and standard tools for sets of finite perimeter.

Strategy. The eigenvalue is reached only at the very end, through the torsion function $u = u_\Omega$ ($-\Delta u = 1$ in Ω , $u \in H_0^1(\Omega)$) and its Saint–Venant deficit $\delta_T(\Omega) = E(\Omega) - E(B^*)$, $E = -\frac{1}{2} \int u$. The heart of the paper is the *sharp Saint–Venant stability* estimate $\mathcal{A}(\Omega)^2 \leq C \delta_T(\Omega)$, obtained by working directly with the torsion superlevel sets $E_\rho = \{u > t(\rho)\}$ of volume $\omega_n \rho^n$ and the *unscaled* moving-ball distance

$$q(\rho) := \inf_{z \in \mathbb{R}^n} |E_\rho \Delta B_\rho(z)|.$$

Because the family $\{E_\rho\}$ is nested, q is uniformly Lipschitz (Lemma 7.3); this is what lets the sparse set of bad-density radii be charged by Lebesgue measure rather than by the deficit.

Outline. The proof proceeds in three blocks.

(I) *Bounded Saint–Venant stability (§§3–9)*. Section 3 proves, on reduced boundaries and for a.e. level, the exact deficit identity $2\delta_T(\Omega) = \int_0^{\|u\|_\infty} (D_H + D_I) / (n^2 \omega_n^{2/n} m^{1-2/n}) dt$, where D_H is the Cauchy–Schwarz defect of $|\nabla u|$ on $\partial^* E_t$ and D_I the squared-perimeter isoperimetric defect; Section 4 derives its profile-gap and bad-density consequences. Section 5 converts the Fusco–Julin inequality into a same-centre control of the homothetic trace. Sections 6–8 prove a finite-perimeter affine first variation for q and the positive kinetic estimate, and Section 9 closes the bounded estimate $\mathcal{A}(\Omega)^2 \leq C(n, R) \delta_T(\Omega)$ for $\Omega \subset B_R$ (Theorem 9.2).

(II) *Bounded reduction (§10)*. A constructive De Giorgi L^∞ truncation produces, from any Ω of small deficit, a bounded surrogate inside a dimensional ball with controlled loss of deficit and asymmetry; gluing the resulting bounded estimate to a trivial far-deficit branch removes the confinement and yields the global sharp Saint–Venant inequality $\delta_T(\Omega) \geq c_{\text{SV}}^{\text{glob}}(n) \mathcal{A}(\Omega)^2$.

(III) *Kohler–Jobin transfer (§11)*. The Kohler–Jobin rearrangement, proved here from scratch, gives the scale-invariant comparison $T(\Omega)\lambda_1(\Omega)^{(n+2)/2} \geq T_n\lambda_1(B_1)^{(n+2)/2}$; combined with the global Saint–Venant inequality and an elementary one-variable estimate it yields Theorem 1.1.

The one substantive external theorem is the Fusco–Julin quantitative isoperimetric inequality; the remaining ingredients are classical tools for sets of finite perimeter, collected in Appendix A, and the dependency certificate of Appendix B.

2. STANDING NOTATION AND SCOPE

Fix $n \geq 2$, $R \geq 1$, and $0 < \rho_* < 1$. Let $\Omega \subset B_R$ be bounded and open with $|\Omega| = \omega_n$, and let $u = u_\Omega \in H_0^1(\Omega)$ be the torsion function,

$$-\Delta u = 1 \quad \text{in } \Omega, \quad u = 0 \quad \text{on } \partial\Omega \quad (H_0^1 \text{ sense}).$$

We write the torsional rigidity $T(\Omega) := \int_\Omega u \, dx$, the Saint–Venant energy $E(\Omega) := -\frac{1}{2}T(\Omega)$, and the Saint–Venant deficit

$$\delta_T(\Omega) := E(\Omega) - E(B_1) = -\frac{1}{2} \int_\Omega u + \frac{1}{2} \int_{B_1} u_{B_1} \geq 0.$$

We write $\lambda_1(\Omega)$ for the first Dirichlet eigenvalue, B^* for the ball with $|B^*| = |\Omega|$, $L_n := \lambda_1(B_1)$, and $T_n := T(B_1) = \omega_n/(n(n+2))$; these enter only in the bounded reduction (§10) and the Kohler–Jobin transfer (§11).

Scope of the bounded core. Blocks (I) of the introduction – Sections 3–9 – are carried out under the standing normalisation $|\Omega| = \omega_n$ and the confinement $\Omega \subset B_R$, with parameters $n \geq 2$, $R \geq 1$, $0 < \rho_* < 1$ fixed throughout. The confinement and the volume normalisation are both removed in §10 by scaling and a constructive truncation, so the final theorem holds for arbitrary open sets of finite measure.

Let $t(\rho)$ be the volume-radius parametrisation of the superlevels:

$$E_\rho := \{u > t(\rho)\}, \quad |E_\rho| = \omega_n \rho^n, \quad 0 < \rho < 1.$$

Set

$$w(\rho) := -t'(\rho), \quad d\mu(\rho) = w(\rho) \, d\rho.$$

For a.e. ρ , E_ρ has finite perimeter, $\mathcal{H}^{n-1}(\partial^* E_\rho \cap \{|\nabla u| = 0\}) = 0$, and the De Giorgi outer normal satisfies

$$\nu_\rho = -\frac{\nabla u}{|\nabla u|} \quad \mathcal{H}^{n-1}\text{-a.e. on } \partial^* E_\rho$$

(these level conventions are established in Lemma 3.1). On those levels define

$$V_\rho := \frac{w(\rho)}{|\nabla u|}, \quad H_{z,\rho}(x) := \frac{(x-z) \cdot \nu_\rho(x)}{\rho}, \quad \bar{V}_\rho := \frac{P(B_\rho)}{P(E_\rho)}.$$

The coarea relation of §3 (Lemma 3.2, with $w(\rho) \alpha(t(\rho)) = P(B_\rho)$) gives

$$(1) \quad \int_{\partial^* E_\rho} V_\rho \, d\mathcal{H}^{n-1} = P(B_\rho), \quad \bar{V}_\rho = \frac{1}{P(E_\rho)} \int_{\partial^* E_\rho} V_\rho \, d\mathcal{H}^{n-1}.$$

The two level-set defects are

$$D_H(t(\rho)) := \left(\int_{\partial^* E_\rho} \frac{1}{|\nabla u|} \, d\mathcal{H}^{n-1} \right) \left(\int_{\partial^* E_\rho} |\nabla u| \, d\mathcal{H}^{n-1} \right) - P(E_\rho)^2, \\ D_I(t(\rho)) := P(E_\rho)^2 - P(B_\rho)^2.$$

Remark 2.1 (Scope). The bounded Saint–Venant estimate is deduced from the exact level-set identity (Theorem 3.11), proved in §3, and its profile-gap and bad-density consequences (Proposition 4.1). The same-centre good-level package (Theorem 5.4) rests on the theorem of Fusco–Julin, used in its exact same-centre form; the radial trace conversion from normal oscillation to the homothetic trace is Lemma 5.3. Every other step, including the first variation and the bad-density bookkeeping, is proved here.

3. FINITE-PERIMETER LEVEL-SET IDENTITIES

All integrals in this section are taken on De Giorgi reduced boundaries and all statements hold for almost every level; no smoothness of $\partial\Omega$, no graph parametrisation of the level sets, and no pointwise lower bound on $|\nabla u|$ are used. We keep the normalisation $|\Omega| = \omega_n$, so the ball of the same volume is B_1 ; the general case follows by scaling. Write

$$M := \|u\|_\infty, \quad L := |\Omega| = \omega_n, \quad m(t) := |\{u > t\}|, \\ \alpha(t) := \int_{\partial^* E_t} \frac{1}{|\nabla u|} d\mathcal{H}^{n-1}, \quad \beta(t) := \int_{\partial^* E_t} |\nabla u| d\mathcal{H}^{n-1}, \quad c_n := n^2 \omega_n^{2/n}.$$

By the Talenti pointwise comparison [17], $u^*(s) \leq u_{B_1}^*(s) = (1 - (s/\omega_n)^{2/n})/(2n)$, so $M \leq 1/(2n) < \infty$; here u^* is the decreasing rearrangement of u .

Regularity and level conventions.

Lemma 3.1 (Critical set and level conventions). *Let u be extended by 0 to $\mathbb{R}^n$. Then:*

- (a) $|\{u = 0\} \cap \Omega| = 0$ and $|\{\nabla u = 0\} \cap \Omega| = 0$;
- (b) for every $t \in (0, M)$, $|\{u = t\}| = 0$; hence m is continuous and strictly decreasing on $(0, M)$;
- (c) for a.e. $t \in (0, M)$: E_t has finite perimeter, $\partial^* E_t \subset \Omega$ up to $\mathcal{H}^{n-1}$ -null sets, $\nu_{E_t} = -\nabla u/|\nabla u|$ $\mathcal{H}^{n-1}$ -a.e. on $\partial^* E_t$, and $\mathcal{H}^{n-1}(\partial^* E_t \cap \{\nabla u = 0\}) = 0$.

Proof. (a) Testing $-\Delta u = 1$ against $u^- \in H_0^1(\Omega)$ gives $\int |\nabla u^-|^2 = -\int u^- \leq 0$, so $u \geq 0$ a.e.; the weak strong maximum principle on each connected component yields $u > 0$ a.e. in Ω [13], i.e. $|\{u = 0\} \cap \Omega| = 0$. Interior elliptic regularity gives $u \in W_{\text{loc}}^{2,p}(\Omega)$ for all $p < \infty$; applying Stampacchia’s theorem [16] to the weak derivatives $\partial_i u$ gives $D^2 u = 0$ a.e. on $\{\nabla u = 0\}$, while $\Delta u = -1$ a.e. Hence $|\{\nabla u = 0\} \cap \Omega| = 0$.

(b) For $t \in (0, M)$ the Sobolev chain rule applied to $(u - t)^+$ gives $\nabla u = 0$ a.e. on $\{u = t\}$; since $\{u = t\} \subset \{u > 0\} \subset \Omega$ up to a null set, part (a) gives $|\{u = t\}| = 0$. A jump of m at t would have size $|\{u = t\}| = 0$, so m is continuous on $(0, M)$. For $0 < a < b < M$, some component of Ω has essential supremum of u exceeding b ; on it $\{a < u < b\}$ is a nonempty open set of positive measure, so $m(a) - m(b) = |\{a < u \leq b\}| > 0$.

(c) By the BV coarea formula [14, Thm. 13.1], [1, Thm. 3.40], for every Borel $\varphi \geq 0$,

$$(2) \quad \int_{\Omega} |\nabla u| \varphi dx = \int_{-\infty}^{\infty} \left(\int_{\partial^* E_t} \varphi d\mathcal{H}^{n-1} \right) dt.$$

Taking $\varphi = \mathbf{1}_{(0,M)}(u)$ (which equals 1 on $\partial^* E_t$ precisely for $t \in (0, M)$) gives $\int_0^M P(E_t) dt = \int_{\{0 < u < M\}} |\nabla u| dx \leq \int_{\Omega} |\nabla u| < \infty$, the last bound because $|\nabla u| \in L^2(\Omega) \subset L^1(\Omega)$ on the bounded set Ω ; hence $P(E_t) < \infty$ for a.e. t . For $t > 0$ the precise representative of u vanishes $\mathcal{H}^{n-1}$ -a.e. on $\mathbb{R}^n \setminus \Omega$ (as $u \in H_0^1(\Omega)$), and for a.e. t one has $\partial^* E_t \subset \{u = t\}$ up to $\mathcal{H}^{n-1}$ -null sets [1, 14]; therefore $\partial^* E_t \subset \Omega$ up to $\mathcal{H}^{n-1}$ -null sets. The orientation $\nu_{E_t} = -\nabla u/|\nabla u|$ is the

standard one for superlevel sets of Sobolev functions [1]. Finally, (2) with $\varphi = \mathbf{1}_{\{\nabla u=0\}}$ gives $\int_0^M \mathcal{H}^{n-1}(\partial^* E_t \cap \{\nabla u = 0\}) dt = \int_{\{\nabla u=0\}} |\nabla u| dx = 0$, so the last assertion holds for a.e. t . $\square$

Statements “for a.e. t ” below refer to the full-measure set of levels furnished by Lemma 3.1(c).

Coarea form of $-m'$ and the flux identity.

Lemma 3.2 (Coarea form of $-m'$). *For a.e. $t \in (0, M)$, $-m'(t) = \alpha(t)$. In particular m is locally absolutely continuous on $(0, M)$ and $\alpha \in L^1_{\text{loc}}((0, M))$.*

Proof. Fix $0 < a < b < M$ and apply (2) to $\varphi = |\nabla u|^{-1} \mathbf{1}_{\{|\nabla u|>0\}} \mathbf{1}_{(a,b)}(u)$. By Lemma 3.1(a) the integrand on the left equals $\mathbf{1}_{\{a < u < b\}}$ up to a null set, so $m(a) - m(b) = \int_a^b \alpha(t) dt$. Local absolute continuity of m and $-m' = \alpha$ a.e. follow. $\square$

Lemma 3.3 (Flux equals volume). *For a.e. $t \in (0, M)$, $\beta(t) = \int_{\partial^* E_t} |\nabla u| d\mathcal{H}^{n-1} = m(t)$.*

Proof. For $\varepsilon > 0$ set $\phi_\varepsilon(s) := \min(1, \max(0, s/\varepsilon))$ and $\eta_\varepsilon := \phi_\varepsilon(u-t) \in H^1_0(\Omega)$. Testing $-\Delta u = 1$ against η_ε ,

$$\int_{\Omega} \nabla u \cdot \nabla \eta_\varepsilon dx = \int_{\Omega} \eta_\varepsilon dx.$$

The Sobolev chain rule [1, Thm. 3.99] gives $\nabla \eta_\varepsilon = \varepsilon^{-1} \nabla u \mathbf{1}_{\{t < u < t+\varepsilon\}}$, so the left side equals $\varepsilon^{-1} \int_{\{t < u < t+\varepsilon\}} |\nabla u|^2$, which by (2) (with $\varphi = |\nabla u| \mathbf{1}_{(t, t+\varepsilon)}(u)$) equals $\varepsilon^{-1} \int_t^{t+\varepsilon} \beta(\tau) d\tau$. As $\varepsilon \downarrow 0$, $\eta_\varepsilon \rightarrow \mathbf{1}_{E_t}$ boundedly (Lemma 3.1(b)), so the right side tends to $m(t)$. Since $\beta \in L^1((0, M))$ by (2) and the Dirichlet identity $\int_0^M \beta = \int_{\Omega} |\nabla u|^2 = \int_{\Omega} u < \infty$, at every Lebesgue point of β the left side tends to $\beta(t)$. Hence $\beta(t) = m(t)$ for a.e. t . $\square$

Remark 3.4 (Why a truncation, not a trace). Lemma 3.3 is the rigorous replacement, on an arbitrary finite-perimeter level set, for the formal divergence computation $m(t) = -\int_{E_t} \Delta u = -\int_{\partial^* E_t} \nabla u \cdot \nu_{E_t} = \beta(t)$ (using $\nabla u \cdot \nu_{E_t} = -|\nabla u|$); no boundary trace of ∇u on $\partial^* E_t$ is required [14, Ch. 17].

The rearranged primitive. Let $I(s) := \int_0^s u^*(\sigma) d\sigma$, $s \in [0, L]$.

Lemma 3.5 (Derivatives of the primitive). *I is absolutely continuous on $[0, L]$ with $I'(s) = u^*(s)$ a.e.; on every compact subinterval of $(0, L)$ the rearrangement u^* is absolutely continuous and*

$$(3) \quad I''(s) = (u^*)'(s) = \frac{1}{m'(u^*(s))} = -\frac{1}{\alpha(u^*(s))} \quad \text{for a.e. } s \in (0, L).$$

Proof. By the Talenti bound $u^* \in L^\infty(0, L) \subset L^1(0, L)$; the Lebesgue differentiation theorem gives $I' = u^*$ a.e. By Lemma 3.2, m is locally absolutely continuous with $m' = -\alpha$ a.e., and m is continuous and strictly decreasing on $(0, M)$ by Lemma 3.1(b), so $m(u^*(s)) = s$. The absolutely continuous inverse theorem for monotone functions [10, Thm. 0.4, §1] gives that u^* is absolutely continuous on compact subintervals of $(0, L)$ with $(u^*)'(s) = 1/m'(u^*(s))$ wherever $m'(u^*(s))$ exists and is nonzero. For a.e. $t \in (0, M)$ one has $0 < |E_t| < |\Omega|$, hence $P(E_t) > 0$ and $\alpha(t) > 0$; thus $\{m' = 0\} = \{\alpha = 0\}$ is null and carries no image measure ($|\{m' = 0\}| \leq \int_{\{m'=0\}} |m'| = 0$). Since $m' = -\alpha$, (3) follows; absolute continuity of u^* on compact subintervals identifies the distributional and a.e. derivatives of $I' = u^*$. $\square$

Remark 3.6 (The corrected second derivative). The factor in (3) is $\alpha = \int_{\partial^* E_t} |\nabla u|^{-1}$, not $\beta = \int_{\partial^* E_t} |\nabla u|$. Since $\beta(t) = m(t)$ (Lemma 3.3), replacing α by β would give $I'' = -1/m$, which holds only when $|\nabla u|$ is constant on $\partial^* E_t$, i.e. on the ball; the discrepancy is exactly the Cauchy–Schwarz defect D_H introduced next, so the substitution would silently delete D_H from the identity.

The exact deficit identity. For a.e. $t \in (0, M)$ the two level-set defects of the standing notation read, as functions of the level,

$$D_H(t) = \alpha(t)\beta(t) - P(E_t)^2, \quad D_I(t) = P(E_t)^2 - c_n m(t)^{2-2/n},$$

consistent with §2 under $t = t(\rho)$ (so $P(B_\rho)^2 = c_n m(t)^{2-2/n}$). By Cauchy-Schwarz on $\partial^* E_t$ applied to $(|\nabla u|^{1/2}, |\nabla u|^{-1/2})$, $P(E_t)^2 \leq \alpha(t)\beta(t)$, so $D_H(t) \geq 0$, with equality iff $|\nabla u|$ is $\mathcal{H}^{n-1}$ -a.e. constant on $\partial^* E_t$. By the De Giorgi isoperimetric inequality [11, 14], $P(E_t) \geq n\omega_n^{1/n} m(t)^{1-1/n}$, so $D_I(t) \geq 0$, with equality iff E_t is equivalent to a ball.

Lemma 3.7 (Pointwise convexity formula). *Set $G(s) := I(s) + \frac{s^{1+2/n}}{(4+2n)\omega_n^{2/n}}$. Then for a.e. $s \in (0, L)$, with $t = u^*(s)$,*

$$G''(s) = \frac{1}{c_n s^{1-2/n}} - \frac{1}{\alpha(t)}.$$

Proof. Direct differentiation gives $\frac{d^2}{ds^2} \frac{s^{1+2/n}}{(4+2n)\omega_n^{2/n}} = \frac{1}{c_n s^{1-2/n}}$; add (3). $\square$

Lemma 3.8 (Convexity identity). $\int_0^L s G''(s) ds = \int_0^M \frac{D_H(t) + D_I(t)}{c_n m(t)^{1-2/n}} dt.$

Proof. Substitute $s = m(t)$, $ds = -\alpha(t) dt$, in $\int_0^L s G''(s) ds$ (Lemma 3.1(b) removes jump terms, and $\{\alpha = 0\}$ carries no image measure):

$$\int_0^L s G'' ds = \int_0^M m(t) \left(\frac{1}{c_n m(t)^{1-2/n}} - \frac{1}{\alpha(t)} \right) \alpha(t) dt = \int_0^M \left(\frac{m(t)\alpha(t)}{c_n m(t)^{1-2/n}} - m(t) \right) dt.$$

Use $m(t) = \beta(t)$ (Lemma 3.3) and add and subtract $P(E_t)^2/(c_n m(t)^{1-2/n})$:

$$\frac{m\alpha}{c_n m^{1-2/n}} - m = \frac{\alpha\beta - c_n m^{2-2/n}}{c_n m^{1-2/n}} = \frac{D_H + D_I}{c_n m^{1-2/n}}. \quad \square$$

Lemma 3.9 (Endpoint convexity-gap formula). *With $\Gamma(\Omega) := G'_-(L) - G(L)/L$, one has $\Gamma(\Omega) = \frac{1}{L} \int_0^L s G''(s) ds$.*

Proof. $G \in W_{\text{loc}}^{2,1}((0, L])$ and $G(0) = I(0) = 0$. Integration by parts on (ε, L) gives $\int_\varepsilon^L s G'' = L G'_-(L) - \varepsilon G'(\varepsilon) - G(L) + G(\varepsilon)$. By Lemma 3.5, $G'(s) = u^*(s) + s^{2/n}/(2n\omega_n^{2/n})$ is bounded, so $\varepsilon G'(\varepsilon) \rightarrow 0$ and $G(\varepsilon) \rightarrow 0$; divide by L . $\square$

Lemma 3.10 (Endpoint value). $\Gamma(\Omega) = \frac{2}{|\Omega|} (E(\Omega) - E(B_1)) = \frac{2}{\omega_n} \delta_T(\Omega).$

Proof. At $s = L$, $u^*(s) \rightarrow 0$ as $s \uparrow L$ (Lemma 3.1 gives $m(t) \uparrow L$ as $t \downarrow 0$), so $G'_-(L) = L^{2/n}/(2n\omega_n^{2/n})$ and $I(L) = \int_\Omega u dx$. With $L = \omega_n$, using $\frac{1}{2n} - \frac{1}{4+2n} = \frac{1}{2n} - \frac{1}{2(n+2)} = \frac{1}{n(n+2)}$,

$$\Gamma(\Omega) = \frac{L^{2/n}}{2n\omega_n^{2/n}} - \frac{L^{2/n}}{(4+2n)\omega_n^{2/n}} - \frac{1}{L} \int_\Omega u = \frac{1}{n(n+2)} - \frac{1}{\omega_n} \int_\Omega u.$$

For the unit-volume ball B_1 , $u_{B_1}(x) = (1 - |x|^2)/(2n)$ and

$$\int_{B_1} u_{B_1} = \frac{1}{2n} n\omega_n \int_0^1 (1 - r^2) r^{n-1} dr = \frac{\omega_n}{2} \left(\frac{1}{n} - \frac{1}{n+2} \right) = \frac{\omega_n}{n(n+2)},$$

so $\frac{1}{\omega_n} \int_{B_1} u_{B_1} = \frac{1}{n(n+2)}$ and $\Gamma(B_1) = 0$ (the Pólya-Szegő identification [15, 2]). Subtracting, $\Gamma(\Omega) = \frac{1}{\omega_n} (\int_{B_1} u_{B_1} - \int_\Omega u) = \frac{2}{\omega_n} (E(\Omega) - E(B_1))$, since $E(\cdot) = -\frac{1}{2} \int(\cdot)$. $\square$

Theorem 3.11 (Exact deficit identity on reduced-boundary level sets). *Let $\Omega \subset \mathbb{R}^n$ be open with $|\Omega| = \omega_n$ and $u \in H_0^1(\Omega)$ the torsion function. Then (2) and Lemmas 3.2, 3.3, 3.5 hold, and*

$$(4) \quad 2\delta_T(\Omega) = \int_0^{\|u\|_\infty} \frac{D_H(t) + D_I(t)}{n^2 \omega_n^{2/n} m(t)^{1-2/n}} dt,$$

with $D_H(t), D_I(t) \geq 0$ for a.e. t . By scaling, the corresponding identity $\frac{2}{|\Omega|}(E(\Omega) - E(B)) = \frac{1}{|\Omega|} \int_0^{\|u\|_\infty} (D_H + D_I)/(c_n m^{1-2/n}) dt$ holds for every open Ω of finite measure.

Proof. Combine Lemmas 3.8, 3.9, 3.10; the scaling statement follows since both sides are 0-homogeneous under $\Omega \mapsto \lambda\Omega$ after the displayed normalisation. $\square$

4. PROFILE GAP AND BAD-DENSITY ESTIMATE

Proposition 4.1 (Level-set budget and profile-gap bad set). *There are computable constants*

$$C_L = C_L(n, \rho_*), \quad C_\tau = C_\tau(n, \rho_*), \quad \kappa = \kappa(\rho_*) > 0,$$

such that, with

$$\rho_\delta := 1 - \kappa \sqrt{\delta_T(\Omega)}$$

and whenever $\rho_\delta \geq (1 + \rho_*)/2$, one has

$$(5) \quad \int_{\rho_*}^{\rho_\delta} (D_H(t(\rho)) + D_I(t(\rho))) d\mu(\rho) \leq C_L \delta_T(\Omega),$$

$$0 \leq w(\rho) \leq \frac{1}{n} \quad \text{for a.e. } \rho,$$

and, with

$$\tau_0 := \frac{\rho_*}{2n}, \quad B_\tau := \{\rho \in [\rho_*, \rho_\delta] : w(\rho) < \tau_0\}, \quad G_\tau := [\rho_*, \rho_\delta] \setminus B_\tau,$$

one has

$$(6) \quad |B_\tau| \leq C_\tau \delta_T(\Omega).$$

Proof. Recall $m(t) = |\{u > t\}|$ and $\alpha(t) = \int_{\partial^* \{u > t\}} |\nabla u|^{-1} d\mathcal{H}^{n-1}$ from §3. Theorem 3.11 gives the exact deficit identity

$$(7) \quad 2\delta_T(\Omega) = \int_0^{\|u\|_\infty} \frac{D_H(t) + D_I(t)}{n^2 \omega_n^{2/n} m(t)^{1-2/n}} dt,$$

all integrals taken on the reduced boundaries $\partial^* \{u > t\}$ and holding for a.e. level.

Changing variables $t = t(\rho)$, $m(t(\rho)) = \omega_n \rho^n$, and $dt = d\mu(\rho)$ in decreasing-radius orientation gives

$$2\delta_T(\Omega) = \int_0^1 \frac{D_H(t(\rho)) + D_I(t(\rho))}{n^2 \omega_n \rho^{n-2}} d\mu(\rho).$$

Since $n \geq 2$ and $0 < \rho \leq 1$,

$$\int_{\rho_*}^{\rho_\delta} (D_H(t(\rho)) + D_I(t(\rho))) d\mu(\rho) \leq 2n^2 \omega_n \delta_T(\Omega),$$

which proves (5), for instance with $C_L = 2n^2 \omega_n$.

It remains to prove the density bounds. Write

$$a(\rho) := w(\rho) = -t'(\rho), \quad a_B(\rho) := \frac{\rho}{n}.$$

We next derive from (7) the profile derivative gap identity

$$(8) \quad 0 \leq a(\rho) \leq a_B(\rho) \quad \text{for a.e. } \rho \in (0, 1), \quad \int_0^1 \rho^n (a_B(\rho) - a(\rho)) d\rho = \frac{2}{\omega_n} \delta_T(\Omega).$$

For completeness, the algebra is as follows. Coarea gives

$$a(\rho) = \frac{n\omega_n \rho^{n-1}}{\alpha(t(\rho))}.$$

For the ball profile the corresponding coarea coefficient is

$$\alpha_B(\rho) = n^2 \omega_n \rho^{n-2}, \quad \frac{n\omega_n \rho^{n-1}}{\alpha_B(\rho)} = \frac{\rho}{n} = a_B(\rho).$$

Since

$$D_H(t(\rho)) + D_I(t(\rho)) = \omega_n \rho^n (\alpha(t(\rho)) - \alpha_B(\rho)),$$

the nonnegativity of $D_H + D_I$ gives $0 \leq a \leq a_B$, and inserting the same formula in (7) gives the integral identity in (8).

Thus $0 \leq w(\rho) \leq \rho/n \leq 1/n$. On

$$B_\tau = \left\{ \rho \in [\rho_*, \rho_\delta] : a(\rho) < \frac{\rho_*}{2n} \right\}$$

one has

$$a_B(\rho) - a(\rho) \geq \frac{\rho}{n} - \frac{\rho_*}{2n} \geq \frac{\rho_*}{2n}.$$

Using $\rho^n \geq \rho_*^n$ on $[\rho_*, \rho_\delta]$ and (8),

$$\frac{\rho_*^{n+1}}{2n} |B_\tau| \leq \int_{B_\tau} \rho^n (a_B - a) d\rho \leq \frac{2}{\omega_n} \delta_T(\Omega).$$

Therefore (6) holds with

$$C_\tau = \frac{4n}{\omega_n \rho_*^{n+1}}.$$

Finally, choose for example $\kappa = (1 - \rho_*)/4$. Then $\rho_\delta \geq (1 + \rho_*)/2$ whenever $\delta_T(\Omega) \leq 1$, and in the large-deficit part of the final theorem this smallness is absorbed into the constant. $\square$

5. QUANTITATIVE ISOPERIMETRY AND SAME-CENTRE TRACE CONTROL

Theorem 5.1 (Fusco–Julin strong quantitative isoperimetry). *For every $n \geq 2$ there is a constant $C_{\text{FJ}}(n)$ with the following property. Let $E \subset \mathbb{R}^n$ be a set of finite perimeter and let $r_E > 0$ be defined by $|E| = \omega_n r_E^n$. Then there is a centre $z_E \in \mathbb{R}^n$ such that*

$$(9) \quad \left[\frac{|E \Delta B_{r_E}(z_E)|}{r_E^n} + \left(\frac{1}{r_E^{n-1}} \int_{\partial^* E} \left| \nu_E(x) - \frac{x - z_E}{|x - z_E|} \right|^2 d\mathcal{H}^{n-1}(x) \right)^{1/2} \right]^2 \leq C_{\text{FJ}}(n) \frac{P(E) - P(B_{r_E})}{r_E^{n-1}}.$$

Proof. This is exactly [12, Theorem 1.1], with the Fusco–Julin asymmetry

$$A(E) = \min_z \left\{ \frac{|E \Delta B_{r_E}(z)|}{r_E^n} + \left(\frac{1}{r_E^{n-1}} \int_{\partial^* E} \left| \nu_E(x) - \nu_{B_{r_E}(z)}(\pi_{z,r_E}(x)) \right|^2 d\mathcal{H}^{n-1}(x) \right)^{1/2} \right\}$$

and their deficit $(P(E) - P(B_{r_E}))/r_E^{n-1}$. Since $\nu_{B_{r_E}(z)}(\pi_{z,r_E}(x)) = (x - z)/|x - z|$, their minimising centre is one centre for both the symmetric-difference term and the boundary-normal oscillation term. $\square$

Corollary 5.2 (Unnormalised same-centre consequences). *Fix $0 < \rho_* < 1$. Let $E \subset \mathbb{R}^n$ be a set of finite perimeter with $|E| = \omega_n \rho^n$, where $\rho \in [\rho_*, 1]$. Then there is a centre $z \in \mathbb{R}^n$ such that*

$$(10) \quad |E \Delta B_\rho(z)|^2 \leq C(n)(P(E) - P(B_\rho)),$$

and

$$(11) \quad \int_{\partial^* E} \left| \nu_E(x) - \frac{x - z}{|x - z|} \right|^2 d\mathcal{H}^{n-1}(x) \leq C(n)(P(E) - P(B_\rho)).$$

Consequently, since $\rho \geq \rho_*$,

$$(12) \quad |E \Delta B_\rho(z)|^2 + \int_{\partial^* E} \left| \nu_E(x) - \frac{x - z}{|x - z|} \right|^2 d\mathcal{H}^{n-1}(x) \leq C(n, \rho_*)(P(E)^2 - P(B_\rho)^2).$$

Proof. Apply Theorem 5.1 with $r_E = \rho$. Each summand in the left side of (9) is nonnegative, so

$$\frac{|E \Delta B_\rho(z)|^2}{\rho^{2n}} + \frac{1}{\rho^{n-1}} \int_{\partial^* E} \left| \nu_E - \frac{x - z}{|x - z|} \right|^2 d\mathcal{H}^{n-1} \leq C(n) \frac{P(E) - P(B_\rho)}{\rho^{n-1}}.$$

Since $\rho \leq 1$, this gives (10) and (11). Finally,

$$P(E) - P(B_\rho) = \frac{P(E)^2 - P(B_\rho)^2}{P(E) + P(B_\rho)} \leq \frac{P(E)^2 - P(B_\rho)^2}{P(B_\rho)} \leq C(n, \rho_*)(P(E)^2 - P(B_\rho)^2).$$

□

Lemma 5.3 (Oriented radial trace). *Fix $R \geq 1$ and $0 < \rho_* < 1$. Let $E \subset B_R$ be a set of finite perimeter with $|E| = \omega_n \rho^n$, where $\rho \in [\rho_*, 1]$, and let $z \in \mathbb{R}^n$ satisfy $|z| \leq R + 1$. Set*

$$e_z(x) := \frac{x - z}{|x - z|}, \quad W_z(x) := \left| \frac{|x - z|}{\rho} - 1 \right|.$$

Then

$$(13) \quad \int_{\partial^* E} W_z d\mathcal{H}^{n-1} \leq \frac{n}{\rho_*} |E \Delta B_\rho(z)| + C(R, \rho_*) \int_{\partial^* E} |\nu_E - e_z|^2 d\mathcal{H}^{n-1}.$$

Proof. Translate the notation temporarily so that $z = 0$, and write $B = B_\rho$, $r = |x|$, $e = x/r$. The point $x = 0$ is $\mathcal{H}^{n-1}$ -negligible on $\partial^* E$, so the value of e there is irrelevant. Define the bounded radial field

$$X(x) := \left| \frac{r}{\rho} - 1 \right| e.$$

For $0 < r < \rho$,

$$\operatorname{div} X = \frac{n-1}{r} - \frac{n}{\rho},$$

while for $r > \rho$,

$$\operatorname{div} X = \frac{n}{\rho} - \frac{n-1}{r}.$$

The field is singular only at the centre and has a Lipschitz radial coefficient away from the centre. To justify Gauss–Green, replace X by $\chi_\varepsilon(r)X$, where $\chi_\varepsilon = 0$ on B_ε and $\chi_\varepsilon = 1$ outside $B_{2\varepsilon}$, smooth the radial profile, apply the classical finite-perimeter Gauss–Green formula [14], and let the smoothing and then $\varepsilon \downarrow 0$. The extra interior term $\int_E \nabla \chi_\varepsilon \cdot X dx$ is supported on the annulus $\{\varepsilon < |x| < 2\varepsilon\}$, where $|\nabla \chi_\varepsilon| \lesssim \varepsilon^{-1}$ and $|X|$ is bounded; since that annulus has volume $O(\varepsilon^n)$, this term is $O(\varepsilon^{n-1})$ and vanishes as $\varepsilon \downarrow 0$. The boundary error tends to zero because $\mathcal{H}^{n-1}(\partial^* E \cap B_{2\varepsilon}) \rightarrow \mathcal{H}^{n-1}(\partial^* E \cap \{0\}) = 0$, and the volume terms converge by dominated

convergence, since r^{-1} is locally integrable in $\mathbb{R}^n$ for $n \geq 2$. Thus

$$\int_{\partial^* E} X \cdot \nu_E d\mathcal{H}^{n-1} = \int_E \operatorname{div} X dx.$$

The same identity for B gives zero, because $X = 0$ on ∂B . Subtracting it yields

$$\int_{\partial^* E} X \cdot \nu_E d\mathcal{H}^{n-1} = \int_{E \setminus B} \operatorname{div} X dx - \int_{B \setminus E} \operatorname{div} X dx.$$

On $E \setminus B$, $\operatorname{div} X \leq n/\rho$. On $B \setminus E$, $-\operatorname{div} X \leq n/\rho$. Hence

$$(14) \quad \int_{\partial^* E} X \cdot \nu_E d\mathcal{H}^{n-1} \leq \frac{n}{\rho} |E \Delta B|.$$

On $\partial^* E$,

$$W_z = X \cdot \nu_E + W_z(1 - e_z \cdot \nu_E) = X \cdot \nu_E + \frac{W_z}{2} |\nu_E - e_z|^2.$$

Since $E \subset B_R$ and $|z| \leq R + 1$, every reduced-boundary point satisfies $|x - z| \leq 2R + 1$, and therefore

$$W_z(x) \leq 1 + \frac{2R + 1}{\rho_*}.$$

Combining this bound with (14) and using $\rho \geq \rho_*$ proves (13). $\square$

Theorem 5.4 (Same-centre good-level geometric package). *There exist constants*

$$\eta_I = \eta_I(n, R, \rho_*) > 0, \quad C_{\text{sc}} = C_{\text{sc}}(n, R, \rho_*),$$

such that the following holds. Define

$$G_I := \{\rho \in [\rho_*, \rho_\delta] : D_I(t(\rho)) \leq \eta_I\}.$$

For a.e. $\rho \in G_I$ there is a centre z_ρ^{sc} such that

$$(15) \quad |E_\rho \Delta B_\rho(z_\rho^{\text{sc}})|^2 + \left(\int_{\partial^* E_\rho} |H_{z_\rho^{\text{sc}}, \rho} - \bar{V}_\rho| d\mathcal{H}^{n-1} \right)^2 \leq C_{\text{sc}} D_I(t(\rho)).$$

Proof. Let $\rho \in G_I$ be a finite-perimeter level. Apply Corollary 5.2 to E_ρ , and let $z = z_\rho^{\text{sc}}$ be the centre obtained there. Since

$$D_I(t(\rho)) = P(E_\rho)^2 - P(B_\rho)^2,$$

(12) gives

$$(16) \quad |E_\rho \Delta B_\rho(z)|^2 + \int_{\partial^* E_\rho} |\nu_\rho - e_z|^2 d\mathcal{H}^{n-1} \leq C(n, \rho_*) D_I(t(\rho)).$$

Choose η_I so small that the first term in (16) implies

$$|E_\rho \Delta B_\rho(z)| < |B_\rho|.$$

Then $E_\rho \cap B_\rho(z)$ has positive measure. Since $E_\rho \subset B_R$, this forces $B_\rho(z) \cap B_R \neq \emptyset$, and therefore

$$(17) \quad |z| \leq R + \rho \leq R + 1.$$

The radial trace lemma gives

$$\int_{\partial^* E_\rho} \left| \frac{|x - z|}{\rho} - 1 \right| d\mathcal{H}^{n-1} \leq C(n, R, \rho_*) (D_I(t(\rho))^{1/2} + D_I(t(\rho))).$$

Reducing $\eta_I \leq 1$, this is bounded by

$$(18) \quad C(n, R, \rho_*) D_I(t(\rho))^{1/2}.$$

For $H_{z,\rho} = (|x - z|/\rho)(e_z \cdot \nu_\rho)$,

$$|H_{z,\rho} - 1| \leq \left| \frac{|x - z|}{\rho} - 1 \right| + |1 - e_z \cdot \nu_\rho| = \left| \frac{|x - z|}{\rho} - 1 \right| + \frac{1}{2} |\nu_\rho - e_z|^2.$$

Together with (16) and (18), this gives

$$(19) \quad \int_{\partial^* E_\rho} |H_{z,\rho} - 1| d\mathcal{H}^{n-1} \leq C(n, R, \rho_*) D_I(t(\rho))^{1/2}.$$

Finally,

$$\bar{V}_\rho = \frac{P(B_\rho)}{P(E_\rho)}, \quad P(E_\rho)|1 - \bar{V}_\rho| = P(E_\rho) - P(B_\rho) \leq C(n, \rho_*) D_I(t(\rho)).$$

Combining this with (19) and again using $\eta_I \leq 1$,

$$\int_{\partial^* E_\rho} |H_{z,\rho} - \bar{V}_\rho| d\mathcal{H}^{n-1} \leq C(n, R, \rho_*) D_I(t(\rho))^{1/2}.$$

Squaring and adding the first estimate in (16) proves (15). $\square$

Lemma 5.5 (Velocity defect controlled by D_H). *For a.e. $\rho \in [\rho_*, \rho_\delta]$,*

$$(20) \quad \left(\int_{\partial^* E_\rho} |V_\rho - \bar{V}_\rho| d\mathcal{H}^{n-1} \right)^2 \leq D_H(t(\rho)).$$

Proof. Write $P = P(E_\rho)$, $P^* = P(B_\rho)$, $\alpha = \int_{\partial^* E_\rho} |\nabla u|^{-1} d\mathcal{H}^{n-1}$, and $\beta = \int_{\partial^* E_\rho} |\nabla u| d\mathcal{H}^{n-1}$. The coarea parametrisation gives $w\alpha = P^*$, and the flux identity gives $\beta = |E_\rho| = \omega_n \rho^n$. With $f := V_\rho = w/|\nabla u|$ and $\bar{f} := \bar{V}_\rho = P^*/P$,

$$\int_{\partial^* E_\rho} f d\mathcal{H}^{n-1} = P^*.$$

Moreover

$$\begin{aligned} \int_{\partial^* E_\rho} \frac{(f - \bar{f})^2}{f} d\mathcal{H}^{n-1} &= \int f d\mathcal{H}^{n-1} - 2\bar{f}P + \bar{f}^2 \int \frac{1}{f} d\mathcal{H}^{n-1} \\ &= -P^* + \frac{(P^*)^2}{P^2} \frac{\beta}{w} = -P^* + \frac{P^*}{P^2} \alpha\beta \\ &= \frac{P^*}{P^2} (\alpha\beta - P^2) = \frac{P^*}{P^2} D_H(t(\rho)). \end{aligned}$$

By Cauchy–Schwarz,

$$\left(\int |f - \bar{f}| d\mathcal{H}^{n-1} \right)^2 \leq \left(\int \frac{(f - \bar{f})^2}{f} d\mathcal{H}^{n-1} \right) \left(\int f d\mathcal{H}^{n-1} \right) = \left(\frac{P^*}{P} \right)^2 D_H(t(\rho)).$$

The isoperimetric inequality gives $P^* \leq P$, hence (20). $\square$

Lemma 5.6 (Superlevel asymmetry transfer). *Let $E \subset \Omega$, $|\Omega| = \omega_n$, and $|E| = \omega_n \rho^n$ with $\rho \in (0, 1]$. Then*

$$\mathcal{A}(\Omega) \leq \mathcal{A}(E) + \frac{2}{\omega_n} |\Omega \setminus E|.$$

Here $\mathcal{A}(E)$ is the Fraenkel asymmetry of E relative to balls of volume $|E|$.

Proof. Let $s = |E|$, $L = |\Omega| = \omega_n$, and $\eta = L - s = |\Omega \setminus E|$. Choose z such that, for an arbitrary $\varepsilon > 0$,

$$|E \Delta B_\rho(z)| \leq s(\mathcal{A}(E) + \varepsilon).$$

Since $B_\rho(z) \subset B_1(z)$,

$$\begin{aligned} |\Omega \Delta B_1(z)| &\leq |\Omega \Delta E| + |E \Delta B_\rho(z)| + |B_\rho(z) \Delta B_1(z)| \\ &= \eta + |E \Delta B_\rho(z)| + \eta. \end{aligned}$$

Divide by L , use $s \leq L$, and let $\varepsilon \downarrow 0$. □

6. COAREA DIFFERENTIATION AND SHELL-ADMISSIBLE RADII

The first variation is proved on a fixed full-measure set of good-density radii. The point of the definition below is that all coarea differentiations are registered in advance by a countable family independent of the radius being tested.

Lemma 6.1 (Two-sided variable-thickness coarea differentiation). *Let $\xi \leq \zeta$ be bounded Borel functions on $\mathbb{R}^n$. Then for Lebesgue-a.e. positive level $s \in (0, \|u\|_\infty)$, with exceptional null set depending only on the pair (ξ, ζ) ,*

$$(21) \quad \lim_{h \downarrow 0} \frac{1}{h} |\{x : h\xi(x) < u(x) - s < h\zeta(x)\}| = \int_{\partial^*\{u>s\}} \frac{\zeta - \xi}{|\nabla u|} d\mathcal{H}^{n-1}.$$

The statement is used only at levels for which the right side is finite and $\mathcal{H}^{n-1}(\partial^*\{u > s\} \cap \{|\nabla u| = 0\}) = 0$.

Proof. By Lemma 3.1(a), $|\{x \in \Omega : |\nabla u(x)| = 0\}| = 0$, so the critical set may be discarded in the volume computations below.

Suppose first that $\xi = \sum_j p_j \mathbf{1}_{A_j}$ and $\zeta = \sum_j q_j \mathbf{1}_{A_j}$ are simple over a common disjoint Borel partition, with $p_j \leq q_j$. By the BV coarea formula applied to $\mathbf{1}_{A_j} |\nabla u|^{-1} \mathbf{1}_{\{|\nabla u|>0\}}$,

$$|\{hp_j < u - s < hq_j\} \cap A_j| = \int_{s+hp_j}^{s+hq_j} \int_{\partial^*\{u>\tau\}} \frac{\mathbf{1}_{A_j}}{|\nabla u|} d\mathcal{H}^{n-1} d\tau.$$

The interval of integration shrinks to s and has length $h(q_j - p_j)$; hence at a Lebesgue point of $\tau \mapsto \int_{\partial^*\{u>\tau\}} \mathbf{1}_{A_j} |\nabla u|^{-1} d\mathcal{H}^{n-1}$, division by h followed by $h \downarrow 0$ gives $(q_j - p_j) \int_{\partial^*\{u>s\}} \mathbf{1}_{A_j} |\nabla u|^{-1} d\mathcal{H}^{n-1}$. Summing over j proves (21) for simple pairs.

For bounded $\xi \leq \zeta$, fix $\eta > 0$ and choose simple pairs that sandwich the slab. Outer: simple $\sigma \leq \xi \leq \zeta \leq \sigma'$ with $\sigma' - \sigma \leq (\zeta - \xi) + 2\eta$; then $\{h\xi < u - s < h\zeta\} \subseteq \{h\sigma < u - s < h\sigma'\}$, so the lim sup of the left side of (21) is at most

$$\int_{\partial^*\{u>s\}} \frac{\sigma' - \sigma}{|\nabla u|} d\mathcal{H}^{n-1} \leq \int_{\partial^*\{u>s\}} \frac{\zeta - \xi}{|\nabla u|} d\mathcal{H}^{n-1} + 2\eta \int_{\partial^*\{u>s\}} \frac{1}{|\nabla u|} d\mathcal{H}^{n-1}.$$

Inner: simple $\sigma_* \geq \xi$, $\sigma'_* \leq \zeta$ with $\sigma_* \leq \sigma'_*$ and $\sigma'_* - \sigma_* \geq (\zeta - \xi - 2\eta)_+$; the reverse inclusion bounds the lim inf from below by

$$\int_{\partial^*\{u>s\}} \frac{(\zeta - \xi - 2\eta)_+}{|\nabla u|} d\mathcal{H}^{n-1}.$$

Letting $\eta \downarrow 0$ gives (21). The exceptional levels are fixed as follows: choose the outer and inner simple sandwiches for the dyadic sequence $\eta = 2^{-m}$, $m \geq 1$, by a fixed dyadic quantisation of the bounded functions ξ, ζ . The union of the exceptional null sets for those countably many simple pairs is still null and depends only on (ξ, ζ) . □

Shell-admissible radii. Let $\mathcal{U}$ be the countable family of finite unions of rational balls whose closures are compactly contained in Ω . This family is cofinal: for every compact $K \Subset \Omega$ there

is $U \in \mathcal{U}$ with $K \subset U \Subset \Omega$. For

$$c, p, q, \lambda, r \in \mathbb{Q}, \quad -1 \leq c \leq 0, \quad p \leq q, \quad 1 \leq \lambda \leq \rho_*^{-1}, \quad r > 0, \quad z, a \in \mathbb{Q}^n,$$

and $U \in \mathcal{U}$, register the bounded Borel pairs

$$(p\mathbf{1}_U, q\mathbf{1}_U)$$

and

$$(\mathbf{1}_U(\min(c, \nabla u \cdot (\lambda(x - z) + a)) - r), \mathbf{1}_U(\max(c, \nabla u \cdot (\lambda(x - z) + a)) + r)).$$

This is a fixed countable family, independent of the radius being tested. We call $\rho \in G_\tau \cap (0, 1)$ *shell-admissible* if:

- (i) E_ρ has finite perimeter, $\partial^* E_\rho \subset \Omega$ up to $\mathcal{H}^{n-1}$ -null sets, and $\mathcal{H}^{n-1}(\partial^* E_\rho \cap \{|\nabla u| = 0\}) = 0$;
- (ii) t is differentiable at ρ , and

$$w(\rho) \int_{\partial^* E_\rho} \frac{1}{|\nabla u|} d\mathcal{H}^{n-1} = P(B_\rho);$$

- (iii) $s = t(\rho)$ is a level where Lemma 6.1 holds for every registered pair above.

Almost every $\rho \in G_\tau \cap (0, 1)$ is shell-admissible. Indeed, each registered pair has an exceptional null set of levels s . If N is such a null set, then the one-dimensional area formula for the monotone absolutely continuous map t gives, on G_τ ,

$$|\{\rho : t(\rho) \in N\}| \leq \tau_0^{-1} \int_{\{t(\rho) \in N\}} w(\rho) d\rho \leq \tau_0^{-1} |N| = 0.$$

The remaining conditions hold for a.e. radius by BV coarea and the finite-perimeter level-set package. The critical-boundary condition follows from the preceding interior critical-set nullity and the coarea formula:

$$\int_0^{\|u\|_\infty} \mathcal{H}^{n-1}(\partial^* \{u > s\} \cap \{|\nabla u| = 0\}) ds = \int_{\{|\nabla u| = 0\}} |\nabla u| dx = 0.$$

Lemma 6.2 (Local BV deformation tail). *Let $E \subset B_R$ be a set of finite perimeter, let $\eta \in C_c^1(\mathbb{R}^n)$ satisfy $0 \leq \eta \leq 1$, and let $W \in C^1$ on a neighbourhood of a ball containing $E \cup \text{spt } \eta$. Set $T_h(x) = x + hW(x)$. Then*

$$\limsup_{h \rightarrow 0} \frac{1}{|h|} \int_{\mathbb{R}^n} \eta(x) |\mathbf{1}_{T_h(E)}(x) - \mathbf{1}_E(x)| dx \leq \int_{\partial^* E} \eta |W| d\mathcal{H}^{n-1}.$$

Proof. For $|h|$ small, T_h is a C^1 diffeomorphism on the fixed ball. Changing variables $x = T_h y$, it is enough, up to a multiplicative $1 + o(1)$ factor, to estimate

$$\int \eta(T_h y) |\mathbf{1}_E(T_h y) - \mathbf{1}_E(y)| dy.$$

Let $u_\varepsilon = \mathbf{1}_E * \varphi_\varepsilon$. For fixed h , the corresponding integrals with u_ε converge to the displayed integral because $u_\varepsilon \rightarrow \mathbf{1}_E$ in L^1 and composition with T_h has bounded Jacobian. For smooth u_ε ,

$$|u_\varepsilon(T_h y) - u_\varepsilon(y)| \leq |h| \int_0^1 |\nabla u_\varepsilon(y + \theta h W(y))| |W(y)| d\theta.$$

After integrating, change variables $x = y + \theta h W(y)$. These maps have Jacobians $1 + o(1)$ uniformly in θ . Thus, for fixed h ,

$$\frac{1}{|h|} \int \eta(T_h y) |u_\varepsilon(T_h y) - u_\varepsilon(y)| dy \leq (1 + o_h(1)) \int \psi_h d|Du_\varepsilon|,$$

where the continuous compactly supported weights ψ_h are uniformly bounded and satisfy $\psi_h \rightarrow \eta|W|$ uniformly as $h \rightarrow 0$. Letting $\varepsilon \downarrow 0$ at fixed h , strict BV_{loc} convergence of the standard mollifications and Reshetnyak continuity for continuous weights [1, 14] give the same bound with $d|D\mathbf{1}_E|$. Taking $\limsup_{h \rightarrow 0}$ then gives

$$\limsup_{h \rightarrow 0} \frac{1}{|h|} \int \eta(x) |\mathbf{1}_{T_h(E)}(x) - \mathbf{1}_E(x)| dx \leq \int \eta|W| d|D\mathbf{1}_E|.$$

Since $|D\mathbf{1}_E| = \mathcal{H}^{n-1} \llcorner \partial^* E$, this is the claim. $\square$

7. AFFINE MOVING-BALL FIRST VARIATION

Lemma 7.1 (Affine shell estimate). *Let ρ be shell-admissible. Fix $z, a \in \mathbb{R}^n$, and define*

$$T_h(x) := z + ha + \left(1 + \frac{h}{\rho}\right)(x - z) = x + h \left(\frac{x - z}{\rho} + a\right).$$

Then

$$(22) \quad \limsup_{h \downarrow 0} \frac{|E_{\rho+h} \Delta T_h(E_\rho)|}{h} \leq \int_{\partial^* E_\rho} |V_\rho - H_{z,\rho} - a \cdot \nu_\rho| d\mathcal{H}^{n-1}.$$

Proof. Let $E = E_\rho$, $W(x) = (x - z)/\rho + a$, and $w = w(\rho)$. Since $\Omega \subset B_R$, W is bounded on Ω . The finite Radon measure

$$\left(\frac{w}{|\nabla u|} + |W| + \left| \frac{w}{|\nabla u|} - W \cdot \nu_\rho \right| \right) \mathcal{H}^{n-1} \llcorner \partial^* E$$

is finite. Fix $\varepsilon > 0$. By inner regularity and $\partial^* E \subset \Omega$ up to $\mathcal{H}^{n-1}$ -null sets, choose a compact $K \subset \partial^* E \cap \Omega$ such that this measure of $\partial^* E \setminus K$ is at most ε . Then choose $U \in \mathcal{U}$ with $K \subset U \Subset \Omega$.

We first estimate the symmetric difference in U . On a neighbourhood of $\bar{U}$, interior elliptic regularity gives $u \in C^1$, and uniformly for $x \in U$,

$$T_h^{-1}(x) = x - hW(x) + O(h^2), \quad u(T_h^{-1}x) = u(x) - h\nabla u(x) \cdot W(x) + o(h).$$

Also $t(\rho + h) = t(\rho) - hw + o(h)$. Set

$$F(x) := w + \nabla u(x) \cdot W(x).$$

For $x \in U$,

$$x \in E_{\rho+h} \iff u(x) > t(\rho) - hw + o(h),$$

while

$$x \in T_h(E) \iff u(x) > t(\rho) + h\nabla u(x) \cdot W(x) + o(h).$$

Thus the part of the symmetric difference in U is contained, up to a uniform $o(h)$ outward error, in the two-sided slab whose endpoints are $\min(-w, \nabla u \cdot W)$ and $\max(-w, \nabla u \cdot W)$. Precisely, fix $\delta > 0$. Choose rational c, λ, z_0, a_0 with $c \in [-1, 0]$, $1 \leq \lambda \leq \rho_*^{-1}$, and

$$|c + w| + \sup_{\bar{U}} |\nabla u \cdot (\lambda(x - z_0) + a_0) - \nabla u \cdot W| \leq \delta.$$

Then choose a rational buffer $r > 2\delta$. For all sufficiently small $h > 0$, the local symmetric difference is contained in the registered slab with endpoints

$$\min(c, \nabla u \cdot (\lambda(x - z_0) + a_0)) - r, \quad \max(c, \nabla u \cdot (\lambda(x - z_0) + a_0)) + r.$$

Lemma 6.1, followed by $\delta, r \downarrow 0$, gives

$$\limsup_{h \downarrow 0} \frac{|(E_{\rho+h} \Delta T_h(E)) \cap U|}{h} \leq \int_{\partial^* E \cap U} \frac{|F|}{|\nabla u|} d\mathcal{H}^{n-1}.$$

On the complement use

$$E_{\rho+h} \Delta T_h(E) \subset (E_{\rho+h} \Delta E) \cup (E \Delta T_h(E)).$$

For the level-set part, the exact volume derivative and the registered localized one-sided coarea formula on U give the complement estimate. Indeed,

$$|E_{\rho+h} \setminus E| = \omega_n((\rho+h)^n - \rho^n) = h P(B_\rho) + o(h) = h w \int_{\partial^* E} \frac{1}{|\nabla u|} d\mathcal{H}^{n-1} + o(h),$$

where the last equality is the shell-admissible flux identity. On U , the same rational-buffer use of Lemma 6.1, now with one endpoint 0, gives

$$\lim_{h \downarrow 0} \frac{|(E_{\rho+h} \setminus E) \cap U|}{h} = w \int_{\partial^* E \cap U} \frac{1}{|\nabla u|} d\mathcal{H}^{n-1}.$$

Subtracting the local shell from the exact total shell yields

$$\limsup_{h \downarrow 0} \frac{|(E_{\rho+h} \Delta E) \setminus U|}{h} \leq w \int_{\partial^* E \setminus U} \frac{1}{|\nabla u|} d\mathcal{H}^{n-1} \leq w \int_{\partial^* E \setminus K} \frac{1}{|\nabla u|} d\mathcal{H}^{n-1}.$$

For the affine deformation part, choose $\eta \in C_c^1(\mathbb{R}^n)$ with $\eta = 0$ on a neighbourhood $N_K \Subset U$ of K , $\eta = 1$ on a fixed neighbourhood of $\overline{B_{R+1}} \setminus U$, and $0 \leq \eta \leq 1$. Since $E \subset B_R$ and W is bounded, $T_h(E) \subset B_{R+1}$ for all small $h > 0$; hence $(E \Delta T_h(E)) \setminus U \subset \overline{B_{R+1}} \setminus U \subset \{\eta = 1\}$. As $\eta = 0$ near K and $0 \leq \eta \leq 1$, Lemma 6.2 gives

$$\limsup_{h \downarrow 0} \frac{|(E \Delta T_h(E)) \setminus U|}{h} \leq \int_{\partial^* E} \eta |W| d\mathcal{H}^{n-1} \leq \int_{\partial^* E \setminus K} |W| d\mathcal{H}^{n-1}.$$

The two tail terms are bounded by 2ε . Therefore

$$\limsup_{h \downarrow 0} \frac{|E_{\rho+h} \Delta T_h(E_\rho)|}{h} \leq \int_{\partial^* E_\rho \cap U} \frac{|F|}{|\nabla u|} d\mathcal{H}^{n-1} + 2\varepsilon \leq \int_{\partial^* E_\rho} \frac{|F|}{|\nabla u|} d\mathcal{H}^{n-1} + 2\varepsilon.$$

Letting $\varepsilon \downarrow 0$, and using

$$\frac{F}{|\nabla u|} = \frac{w}{|\nabla u|} + \frac{\nabla u \cdot W}{|\nabla u|} = V_\rho - W \cdot \nu_\rho = V_\rho - H_{z_\rho, \rho} - a \cdot \nu_\rho$$

on $\partial^* E_\rho$, proves the estimate. $\square$

Theorem 7.2 (One-sided first variation of q). *For every $\rho \in (0, 1]$, define*

$$q(\rho) := \inf_{z \in \mathbb{R}^n} |E_\rho \Delta B_\rho(z)|.$$

At every shell-admissible $\rho \in (0, 1)$,

$$(23) \quad D^+ q(\rho) := \limsup_{h \downarrow 0} \frac{q(\rho+h) - q(\rho)}{h} \leq \frac{n}{\rho} q(\rho) + \inf_{a \in \mathbb{R}^n} \int_{\partial^* E_\rho} |V_\rho - H_{z_\rho, \rho} - a \cdot \nu_\rho| d\mathcal{H}^{n-1},$$

where z_ρ is any minimiser of $z \mapsto |E_\rho \Delta B_\rho(z)|$.

Proof. The minimiser exists. Indeed, $z \mapsto |E_\rho \Delta B_\rho(z)|$ is continuous by translation continuity in L^1 , and if $|z| > R + \rho$, then $B_\rho(z) \cap E_\rho = \emptyset$, so the value is $2|E_\rho|$. Thus the infimum is attained in a closed ball.

Fix a minimiser z_ρ and $a \in \mathbb{R}^n$. For the affine map in Lemma 7.1,

$$T_h(B_\rho(z_\rho)) = B_{\rho+h}(z_\rho + ha).$$

Therefore

$$\begin{aligned} q(\rho + h) &\leq |E_{\rho+h} \Delta T_h(B_\rho(z_\rho))| \\ &\leq |E_{\rho+h} \Delta T_h(E_\rho)| + |T_h(E_\rho) \Delta T_h(B_\rho(z_\rho))|. \end{aligned}$$

Since T_h is injective affine with constant Jacobian $(1 + h/\rho)^n$,

$$|T_h(E_\rho) \Delta T_h(B_\rho(z_\rho))| = \left(1 + \frac{h}{\rho}\right)^n q(\rho).$$

Subtract $q(\rho)$, divide by h , take the limsup as $h \downarrow 0$, and use Lemma 7.1. Then take the infimum over a . $\square$

Lemma 7.3 (Unconditional Lipschitz bound for q). *For $0 < \rho \leq \sigma \leq 1$,*

$$|q(\sigma) - q(\rho)| \leq 2\omega_n(\sigma^n - \rho^n) \leq 2n\omega_n|\sigma - \rho|.$$

Consequently q is absolutely continuous and differentiable a.e.

Proof. Use the same centre for the two radii and the nestedness $E_\rho \subset E_\sigma$, $B_\rho(z) \subset B_\sigma(z)$:

$$q(\sigma) \leq q(\rho) + |E_\sigma \setminus E_\rho| + |B_\sigma \setminus B_\rho| = q(\rho) + 2\omega_n(\sigma^n - \rho^n).$$

Interchange ρ and σ to get the reverse inequality. $\square$

8. POSITIVE KINETIC ESTIMATE

Lemma 8.1 (Transfer to a distance-minimising centre). *After reducing $\eta_I = \eta_I(n, R, \rho_*)$ if necessary, the following holds. Let $\rho \in G_I$ be a level where Theorem 5.4 holds, and let $z_\rho^{\min}$ minimise $z \mapsto |E_\rho \Delta B_\rho(z)|$. Then*

$$(24) \quad \left(\int_{\partial^* E_\rho} |H_{z_\rho^{\min}, \rho} - \bar{V}_\rho| d\mathcal{H}^{n-1} \right)^2 \leq C(n, R, \rho_*) D_I(t(\rho)).$$

Proof. Let z_ρ^{sc} be the centre from Theorem 5.4. Minimality gives

$$|E_\rho \Delta B_\rho(z_\rho^{\min})| \leq |E_\rho \Delta B_\rho(z_\rho^{\text{sc}})| \leq C D_I(t(\rho))^{1/2}.$$

Thus

$$|B_\rho(z_\rho^{\min}) \Delta B_\rho(z_\rho^{\text{sc}})| \leq C D_I(t(\rho))^{1/2}.$$

For equal balls with centre distance d ,

$$|B_\rho(z_1) \Delta B_\rho(z_2)| \geq c(n) \rho^{n-1} \min\{d, \rho\}.$$

Indeed, in coordinates $x = z_1 + s e + y$ with $e = (z_2 - z_1)/d$ and $y \perp e$, one has $B_\rho(z_1) \setminus B_\rho(z_2) = \{s^2 + |y|^2 < \rho^2, s \leq d/2\}$; integrating the $(n-1)$ -dimensional cross-sections over $s \in [0, \frac{1}{2} \min\{d, \rho\}]$, where $\rho^2 - s^2 \geq \frac{3}{4} \rho^2$, gives $|B_\rho(z_1) \setminus B_\rho(z_2)| \geq \frac{1}{2} \omega_{n-1} (3/4)^{(n-1)/2} \rho^{n-1} \min\{d, \rho\}$, and the symmetric difference is twice this. Choose η_I so small that the preceding upper bound forces the linear branch $d < \rho$. Since $\rho \geq \rho_*$,

$$|z_\rho^{\min} - z_\rho^{\text{sc}}| \leq C(n, \rho_*) D_I(t(\rho))^{1/2}.$$

The homothetic trace is affine in the centre:

$$H_{z_\rho^{\min}, \rho} - H_{z_\rho^{\text{sc}}, \rho} = -\frac{(z_\rho^{\min} - z_\rho^{\text{sc}}) \cdot \nu_\rho}{\rho}.$$

On G_I , the perimeter is uniformly bounded because $D_I(t(\rho)) \leq \eta_I$ and $\rho \in [\rho_*, 1]$. Hence

$$\int_{\partial^* E_\rho} |H_{z_\rho^{\min}, \rho} - H_{z_\rho^{\text{sc}}, \rho}| d\mathcal{H}^{n-1} \leq C(n, R, \rho_*) D_I(t(\rho))^{1/2}.$$

Combining this with (15) via the triangle inequality,

$$\int_{\partial^* E_\rho} |H_{z_\rho^{\min}, \rho} - \bar{V}_\rho| d\mathcal{H}^{n-1} \leq \int_{\partial^* E_\rho} |H_{z_\rho^{\min}, \rho} - H_{z_\rho^{\text{sc}}, \rho}| d\mathcal{H}^{n-1} + \int_{\partial^* E_\rho} |H_{z_\rho^{\text{sc}}, \rho} - \bar{V}_\rho| d\mathcal{H}^{n-1} \leq C(n, R, \rho_*) D_I(t(\rho))^{1/2},$$

and squaring gives (24). $\square$

Proposition 8.2 (Positive kinetic bound). *Let $J \subset [\rho_*, \rho_\delta]$ be an interval. Then*

$$\int_J (q'_+(\rho))^2 d\rho \leq C(n, R, \rho_*) \delta_T(\Omega),$$

where $q'_+(\rho) = \max\{q'(\rho), 0\}$ at a.e. differentiability points.

Proof. Let

$$G := J \cap G_I \cap G_\tau, \quad \mathcal{B} := J \setminus G.$$

Since q is Lipschitz, q' exists a.e.; shell-admissible radii have full measure in G_τ . At points where both properties hold, $D^+q = q'$. The right side of Theorem 7.2 is nonnegative, so the theorem also bounds q'_+ . On G , outside this null set, Theorem 7.2, Theorem 5.4, Lemma 8.1, and Lemma 5.5 give

$$(q'_+)^2 \leq C(n, R, \rho_*) (D_H(t(\rho)) + D_I(t(\rho))).$$

Indeed $q^2 \leq CD_I$ on G_I , and

$$\inf_a \int |V_\rho - H_{z_\rho^{\min}, \rho} - a \cdot \nu_\rho| \leq \int |V_\rho - \bar{V}_\rho| + \int |H_{z_\rho^{\min}, \rho} - \bar{V}_\rho|.$$

On G_τ , $d\rho \leq \tau_0^{-1} d\mu$. Hence the budget (5) yields

$$\int_G (q'_+)^2 d\rho \leq C \delta_T(\Omega).$$

It remains to pay $\mathcal{B}$. By Lemma 7.3, $q'_+ \leq 2n\omega_n$ a.e. The bad-density part has $|J \cap B_\tau| \leq C \delta_T(\Omega)$. For the bad-isoperimetric part on good density,

$$|J \cap G_\tau \cap G_I^c| \leq \tau_0^{-1} \mu(G_I^c) \leq \frac{C}{\eta_I} \int_J D_I(t(\rho)) d\mu(\rho) \leq C \delta_T(\Omega).$$

Therefore $|\mathcal{B}| \leq C \delta_T(\Omega)$, and

$$\int_{\mathcal{B}} (q'_+)^2 d\rho \leq C \delta_T(\Omega). \quad \square$$

9. ENDPOINT TRACE AND BOUNDED STABILITY

Lemma 9.1 (Moving-ball endpoint trace). *Let*

$$L_0 := \frac{1 - \rho_*}{8}, \quad J := [\rho_\delta - L_0, \rho_\delta].$$

Assume $\delta_T(\Omega)$ is small enough that $\rho_\delta \geq (1 + \rho_)/2$. Then $J \subset [\rho_*, \rho_\delta]$ and*

$$q(\rho_\delta)^2 \leq C(n, R, \rho_*) \delta_T(\Omega).$$

Proof. The inclusion $J \subset [\rho_*, \rho_\delta]$ follows from $\rho_\delta - \rho_* \geq (1 - \rho_*)/2$. On $G = J \cap G_I \cap G_\tau$,

$$q(\rho)^2 \leq CD_I(t(\rho)), \quad d\rho \leq \tau_0^{-1} d\mu(\rho),$$

so

$$\int_G q(\rho)^2 d\rho \leq C \delta_T(\Omega).$$

On $J \setminus G$, the measure is $O(\delta_T)$ by the proof of Proposition 8.2, while $q \leq 2\omega_n$. Hence

$$\int_J q(\rho)^2 d\rho \leq C \delta_T(\Omega).$$

There is therefore $\rho_0 \in J$ with

$$q(\rho_0)^2 \leq C L_0^{-1} \delta_T(\Omega).$$

Since q is absolutely continuous and $\rho_0 \leq \rho_\delta$,

$$q(\rho_\delta) \leq q(\rho_0) + \int_{\rho_0}^{\rho_\delta} q'_+(\rho) d\rho.$$

By Cauchy–Schwarz and Proposition 8.2,

$$q(\rho_\delta) \leq C \delta_T(\Omega)^{1/2} + L_0^{1/2} \left(\int_J (q'_+)^2 d\rho \right)^{1/2} \leq C \delta_T(\Omega)^{1/2}.$$

□

Theorem 9.2 (Bounded sharp Saint–Venant stability). *There are computable constants $C = C(n, R, \rho_*)$ and $\delta_0 = \delta_0(n, R, \rho_*) > 0$, depending only on the level-set identity constants, the Fusco–Julin constant, and the explicit constants above, such that every bounded open $\Omega \subset B_R$, $|\Omega| = \omega_n$, satisfies*

$$\mathcal{A}(\Omega)^2 \leq C \delta_T(\Omega).$$

Proof. Let κ be the constant fixed in Proposition 4.1. First assume $\delta_T(\Omega) \leq \delta_0$, with δ_0 chosen so that $\rho_\delta = 1 - \kappa \sqrt{\delta_T(\Omega)} \geq (1 + \rho_*)/2$, $\delta_T(\Omega) \leq 1$, and the smallness thresholds used in Theorem 5.4 and Lemma 8.1 hold. Lemma 9.1 gives

$$q(\rho_\delta)^2 \leq C \delta_T(\Omega).$$

Since $|E_{\rho_\delta}| = \omega_n \rho_\delta^n$ and $\rho_\delta \geq \rho_*$,

$$\mathcal{A}(E_{\rho_\delta})^2 \leq \frac{q(\rho_\delta)^2}{\omega_n^2 \rho_\delta^{2n}} \leq C(n, \rho_*) \delta_T(\Omega).$$

Moreover $E_{\rho_\delta} \subset \Omega$, and by the definition of ρ_δ ,

$$|\Omega \setminus E_{\rho_\delta}| = \omega_n(1 - \rho_\delta^n) \leq n\omega_n(1 - \rho_\delta) = n\omega_n \kappa \delta_T(\Omega)^{1/2}.$$

Lemma 5.6 gives

$$\mathcal{A}(\Omega) \leq \mathcal{A}(E_{\rho_\delta}) + C(n, \rho_*) \delta_T(\Omega)^{1/2}.$$

Squaring proves the estimate in the small-deficit regime.

If $\delta_T(\Omega) > \delta_0$, then $\mathcal{A}(\Omega) < 2$, so

$$\mathcal{A}(\Omega)^2 \leq \frac{4}{\delta_0} \delta_T(\Omega).$$

Enlarging C completes the proof. □

10. BOUNDED REDUCTION

The purpose of this section is to remove the boundedness hypothesis from the sharp Saint–Venant stability inequality. The input is the *bounded* estimate proved in the earlier part of the manuscript (Theorem 9.2), which we restate here in energy form.

Theorem 10.1 (Bounded sharp Saint–Venant). *For every $n \geq 2$ and every $R \geq 2$ there is a constant $c_{\text{SV}}(n, R) > 0$ such that*

$$(25) \quad E(\Omega) - E(B_1) \geq c_{\text{SV}}(n, R) \mathcal{A}(\Omega)^2$$

for every open set $\Omega \subset B_R$ with $|\Omega| = \omega_n$.

This is Theorem 9.2, restated in energy form: with the constant $C(n, R, \rho_*)$ there one may take $c_{\text{SV}}(n, R) = 1/C(n, R, \rho_*)$ and $\delta_T(\Omega) = E(\Omega) - E(B_1)$, and the parameter ρ_* is fixed at, say, $\rho_* = \frac{3}{4}$.

Our goal is the global statement.

Theorem 10.2 (Global sharp Saint–Venant stability). *There is a dimensional constant $c_{\text{SV}}^{\text{glob}}(n) > 0$ such that*

$$(26) \quad E(\Omega) - E(B_1) \geq c_{\text{SV}}^{\text{glob}}(n) \mathcal{A}(\Omega)^2$$

for every open set $\Omega \subset \mathbb{R}^n$ with $|\Omega| = \omega_n$. Equivalently, after rescaling, for every open set Ω of finite measure, with B^* the ball of the same volume,

$$(27) \quad T(B^*) - T(\Omega) = 2(E(\Omega) - E(B^*)) \geq 2c_{\text{SV}}^{\text{glob}}(n) \left(\frac{|\Omega|}{\omega_n}\right)^{(n+2)/n} \mathcal{A}(\Omega)^2.$$

The strategy is the constructive reduction of Brasco, De Philippis and Velichkov [3, §5]. From a set Ω with small Saint–Venant deficit we manufacture a *bounded* surrogate $\tilde{\Omega}$, of the same volume, of dimensionally bounded diameter, with comparable deficit and with controlled loss of asymmetry; applying Theorem 10.1 to $\tilde{\Omega}$ at a fixed dimensional radius and transferring the estimate back to Ω closes the small-deficit regime. The complementary large-deficit regime is trivial because $\mathcal{A} < 2$. The whole construction is explicit: no compactness, no minimising sequence and no contradiction argument enter.

The two technical ingredients are an L^∞ tail estimate for the torsion function (Lemma 10.5, proved by a De Giorgi iteration) and the truncation construction itself (Lemma 10.9). Both are proved here in full rather than cited.

10.1. Standing notation and the standard tools we cite. Throughout, $n \geq 2$ and all sets are open subsets of $\mathbb{R}^n$ of finite Lebesgue measure. We write $u_\Omega \in H_0^1(\Omega)$ for the torsion function, i.e. the unique weak solution of $-\Delta u_\Omega = 1$ in Ω , $u_\Omega = 0$ on $\partial\Omega$, equivalently the minimiser of

$$(28) \quad J_\Omega(v) = \frac{1}{2} \int_\Omega |\nabla v|^2 dx - \int_\Omega v dx, \quad v \in H_0^1(\Omega),$$

and $E(\Omega) = J_\Omega(u_\Omega) = \min_{H_0^1(\Omega)} J_\Omega$. Testing the equation with u_Ω gives the standard identities

$$(29) \quad \int_\Omega |\nabla u_\Omega|^2 dx = \int_\Omega u_\Omega dx = T(\Omega), \quad E(\Omega) = -\frac{1}{2} \int_\Omega u_\Omega dx = -\frac{1}{2} T(\Omega).$$

We use $\omega_n = |B_1|$, $T_n = T(B_1) = \omega_n/(n(n+2))$, the Fraenkel asymmetry $\mathcal{A}(\Omega) = \inf_{x_0} |\Omega \triangle (B_1 + x_0)| / \omega_n$ (for $|\Omega| = \omega_n$ this lies in $[0, 2)$), and the scale-invariant deficit

$$(30) \quad D(\Omega) = E(\Omega) |\Omega|^{-(n+2)/n} - E(B_1) |B_1|^{-(n+2)/n},$$

so that $D(\Omega) = \omega_n^{-(n+2)/n} (E(\Omega) - E(B_1)) \geq 0$ when $|\Omega| = \omega_n$. The non-negativity of D is the *Saint–Venant inequality* (equivalently Talenti’s comparison theorem): among sets of given

volume the ball minimises E , i.e.

$$(31) \quad E(\Omega) |\Omega|^{-(n+2)/n} \geq E(B) |B|^{-(n+2)/n} \quad \text{for every ball } B.$$

We treat (31) as a known classical result.

The external analytic facts we invoke without proof are:

- the *Sobolev inequality* $\|w\|_{L^{2^*}(\mathbb{R}^n)} \leq S_n \|\nabla w\|_{L^2(\mathbb{R}^n)}$ for $w \in H^1(\mathbb{R}^n)$ when $n \geq 3$ (with $2^* = 2n/(n-2)$), and its $n=2$ surrogate explained in Remark 10.6;
- the *Saint-Venant/Talenti inequality* (31), used in the truncation argument and in the global L^∞ bound (32) below;
- Theorem 10.1, the bounded stability input;
- the *suboptimal* torsional stability bound $\mathcal{A}(\Omega)^4 \leq C(n)D(\Omega)$ (Lemma 10.7), used only to seed the truncation iteration. This is the single nontrivial result we cite rather than prove; see Remark 10.8.

Everything else — in particular the fast-convergence iteration lemma (Lemma 10.3), the De Giorgi L^∞ tail estimate (Lemma 10.5), the energy-comparison inequality, the truncation iteration (Lemma 10.9) and the gluing — is proved from scratch.

We will also need the elementary pointwise sup-bound on the torsion function. By Talenti's symmetrisation comparison the torsion function of Ω is dominated, after Schwarz symmetrisation, by that of the ball of the same volume; in particular

$$(32) \quad \|u_\Omega\|_{L^\infty(\Omega)} \leq \|u_{B^*}\|_{L^\infty(B^*)} = \frac{(|\Omega|/\omega_n)^{2/n}}{2n} =: \Lambda_n |\Omega|^{2/n}, \quad \Lambda_n := \frac{\omega_n^{-2/n}}{2n},$$

where B^* is the ball with $|B^*| = |\Omega|$ and we used $u_{B^*}(x) = (r^2 - |x|^2)/(2n)$. For $|\Omega| = \omega_n$ this reads $\|u_\Omega\|_{L^\infty(\Omega)} \leq 1/(2n)$. We record (32) as a consequence of the cited Talenti comparison.

10.2. The fast-convergence iteration lemma. The De Giorgi scheme reduces an L^∞ bound to the vanishing of a nonlinear recursion. We isolate the underlying numerical lemma and prove it, so that nothing is cited at this step. The mechanism is that a super-quadratic recursion with a small enough starting value beats any fixed geometric growth factor b^k .

Lemma 10.3 (Fast geometric convergence). *Let $(a_k)_{k \geq 0}$ be non-negative reals with*

$$(33) \quad a_{k+1} \leq C b^k a_k^{1+\beta}, \quad k \geq 0,$$

where $C > 0$, $b > 1$, $\beta > 0$. If

$$(34) \quad a_0 \leq C^{-1/\beta} b^{-1/\beta^2},$$

then

$$(35) \quad a_k \leq b^{-k/\beta} a_0 \quad \text{for all } k \geq 0,$$

so $a_k \rightarrow 0$.

Proof. Put $r := b^{-1/\beta} \in (0, 1)$ and $\theta := C a_0^\beta$. The smallness hypothesis (34) is precisely $\theta \leq r$. We prove the target $a_k \leq r^k a_0$ by induction; this is (35), and $r < 1$ then gives $a_k \rightarrow 0$.

Base case $k=0$: equality. *Inductive step.* Assume $a_k \leq r^k a_0$. Using (33),

$$a_{k+1} \leq C b^k a_k^{1+\beta} = C b^k a_k a_k^\beta \leq C b^k a_k (r^k a_0)^\beta = (C a_0^\beta) (b r^\beta)^k a_k = \theta (b r^\beta)^k a_k.$$

By the definition of r we have $r^\beta = b^{-1}$, hence $b r^\beta = 1$ and the geometric factor collapses:

$$a_{k+1} \leq \theta a_k \leq r a_k \leq r \cdot r^k a_0 = r^{k+1} a_0,$$

using $\theta \leq r$ and the inductive hypothesis. This closes the induction. $\square$

Remark 10.4. The identity $br^\beta = 1$ with $r = b^{-1/\beta}$ is the algebraic mechanism behind the convergence: the super-quadratic gain $a_k^{1+\beta}$ converts, once a_0 is small, the growing factor b^k into a contraction.

10.3. The De Giorgi L^∞ tail estimate.

Lemma 10.5 (L^∞ tail estimate). *Let $n \geq 3$. There is a dimensional constant $C_b = C_b(n) > 0$ such that for every open Ω with $|\Omega| = \omega_n$ and every $R \geq 1$,*

$$(36) \quad \|u_\Omega\|_{L^\infty(\Omega \setminus B_{R+1})} \leq C_b |\Omega \setminus B_R|^{1/n}.$$

For $n = 2$ the same holds with the exponent $1/n$ replaced by $\frac{1}{2} - \frac{1}{p}$ for any fixed $p \in (2, \infty)$ (Remark 10.6); this weaker but still super-linear power is all that the truncation uses.

Proof. If $|\Omega \setminus B_R| = 0$ then $u_\Omega = 0$ a.e. on $\Omega \setminus B_R \supset \Omega \setminus B_{R+1}$ and (36) is trivial, so assume $|\Omega \setminus B_R| > 0$. We carry out a De Giorgi iteration on the annular region $\Omega \setminus B_R$.

Set-up: shrinking radii, cut-offs, and rising levels. For $k \in \mathbb{N}_0$ put

$$R_k := R + 1 - 2^{-k},$$

so that $R_0 = R$, the radii increase to $R_\infty = R + 1$, and $R_k \geq R \geq 1$. Let $\varphi_k \in C^{0,1}(\mathbb{R}^n)$ be the radial cut-off

$$\varphi_k(x) := \phi_k(|x|), \quad \phi_k = 0 \text{ on } [0, R_{k-1}], \quad \phi_k = 1 \text{ on } [R_k, \infty), \quad \phi_k \text{ affine on } [R_{k-1}, R_k],$$

for $k \geq 1$, and $\varphi_0 \equiv 1$ on $\{|x| \geq R_0\}$, extended by 0 inside (we only use φ_k for $k \geq 1$ in the recursion). Then $0 \leq \varphi_k \leq 1$, $\varphi_k \equiv 0$ on $B_{R_{k-1}}$, $\varphi_k \equiv 1$ on $\mathbb{R}^n \setminus B_{R_k}$, and

$$|\nabla \varphi_k| \leq \frac{1}{R_k - R_{k-1}} = \frac{1}{2^{-(k-1)} - 2^{-k}} = 2^k \leq 2^{k+1}.$$

Choose rising levels

$$s_k := M |\Omega \setminus B_R|^{1/n} (1 - 2^{-k}), \quad k \geq 0,$$

where $M = M(n) \geq 1$ is a free constant fixed at the end. Thus $s_0 = 0$, $s_k \uparrow s_\infty := M |\Omega \setminus B_R|^{1/n}$, and

$$(37) \quad s_{k+1} - s_k = M |\Omega \setminus B_R|^{1/n} 2^{-(k+1)}.$$

Caccioppoli inequality from the truncated test function. Fix $k \geq 1$. The function $(u_\Omega - s_k)_+ \in H_0^1(\Omega)$ (it vanishes where $u_\Omega \leq s_k$, in particular near $\partial\Omega$ since $u_\Omega \geq 0$ and $s_k \geq 0$), and φ_k is bounded Lipschitz, so

$$v_k := \varphi_k^2 (u_\Omega - s_k)_+ \in H_0^1(\Omega)$$

is an admissible test function. Insert it into the weak formulation $\int_\Omega \nabla u_\Omega \cdot \nabla v \, dx = \int_\Omega v \, dx$. Using $\nabla v_k = \varphi_k^2 \nabla(u_\Omega - s_k)_+ + 2\varphi_k(u_\Omega - s_k)_+ \nabla \varphi_k$ and $\nabla u_\Omega \cdot \nabla(u_\Omega - s_k)_+ = |\nabla(u_\Omega - s_k)_+|^2$ (the gradient of u_Ω equals that of $(u_\Omega - s_k)_+$ on $\{u_\Omega > s_k\}$ and both factors vanish elsewhere), we get

$$\int \varphi_k^2 |\nabla(u_\Omega - s_k)_+|^2 \, dx + 2 \int \varphi_k(u_\Omega - s_k)_+ \nabla \varphi_k \cdot \nabla(u_\Omega - s_k)_+ \, dx = \int \varphi_k^2 (u_\Omega - s_k)_+ \, dx.$$

The middle term is bounded by Young's inequality, $2|\varphi_k(u_\Omega - s_k)_+ \nabla \varphi_k \cdot \nabla(u_\Omega - s_k)_+| \leq \frac{1}{2}\varphi_k^2 |\nabla(u_\Omega - s_k)_+|^2 + 2|\nabla \varphi_k|^2 (u_\Omega - s_k)_+^2$, and absorbing the first piece on the left yields the Caccioppoli estimate

$$(38) \quad \int |\nabla(\varphi_k(u_\Omega - s_k)_+)|^2 dx \leq C_1 \int |\nabla \varphi_k|^2 (u_\Omega - s_k)_+^2 dx + C_1 \int \varphi_k^2 (u_\Omega - s_k)_+ dx,$$

with an absolute constant C_1 (one checks $C_1 = 10$ works after expanding $|\nabla(\varphi_k w)|^2 \leq 2\varphi_k^2 |\nabla w|^2 + 2|\nabla \varphi_k|^2 w^2$ and combining with the absorbed bound; the precise value is immaterial). Here we wrote $w = (u_\Omega - s_k)_+$.

Sobolev inequality. Set $w_k := \varphi_k(u_\Omega - s_k)_+ \in H_0^1(\mathbb{R}^n)$ and

$$A_k := \{\varphi_k(u_\Omega - s_k)_+ > 0\} = \{u_\Omega > s_k\} \cap \{\varphi_k > 0\} \subset \{u_\Omega > s_k\} \cap (\mathbb{R}^n \setminus B_{R_{k-1}}).$$

For $n \geq 3$ the Sobolev inequality and Hölder's inequality on the support A_k give

$$(39) \quad \int w_k^2 dx \leq \left(\int w_k^{2^*} dx \right)^{2/2^*} |A_k|^{1-2/2^*} \leq S_n^2 |A_k|^{2/n} \int |\nabla w_k|^2 dx,$$

using $1 - 2/2^* = 2/n$. (The $n = 2$ case is handled in Remark 10.6; the resulting estimate has the same shape with $2/n$ replaced by any fixed exponent in $(0, 1)$, which only changes the dimensional constants.)

Combining (39) with (38), $|\nabla \varphi_k| \leq 2^{k+1}$ (so $|\nabla \varphi_k|^2 \leq 4^{k+1}$) and $0 \leq \varphi_k \leq 1$, we obtain

$$(40) \quad \int w_k^2 dx \leq S_n^2 C_1 |A_k|^{2/n} \left(4^{k+1} \int_{A_k} (u_\Omega - s_k)_+^2 dx + \int_{A_k} (u_\Omega - s_k)_+ dx \right),$$

where for the second term we used $\int \varphi_k^2 (u_\Omega - s_k)_+ \leq \int_{A_k} (u_\Omega - s_k)_+$. By the global sup-bound (32) at $|\Omega| = \omega_n$ we have $0 \leq (u_\Omega - s_k)_+ \leq u_\Omega \leq 1/(2n) \leq 1$ on Ω , hence both $(u_\Omega - s_k)_+^2 \leq 1$ and $(u_\Omega - s_k)_+ \leq 1$ there, so each integral over A_k is at most $|A_k|$. Therefore

$$(41) \quad \int w_k^2 dx \leq S_n^2 C_1 (4^{k+1} + 1) |A_k|^{2/n} |A_k| \leq C_2 4^k |A_k|^{1+2/n},$$

with $C_2 = C_2(n) := 8 S_n^2 C_1$ (using $4^{k+1} + 1 \leq 8 \cdot 4^k$).

From energy to a measure recursion. Restrict the left-hand side of (41) to the smaller set where the *next* level is exceeded. On $\{u_\Omega > s_{k+1}\} \cap (\mathbb{R}^n \setminus B_{R_k})$ we have $\varphi_k = 1$ (since this set lies outside B_{R_k}) and $u_\Omega - s_k > s_{k+1} - s_k$, hence $w_k^2 = (u_\Omega - s_k)_+^2 \geq (s_{k+1} - s_k)^2$ there. Therefore

$$(42) \quad (s_{k+1} - s_k)^2 |\{u_\Omega > s_{k+1}\} \cap (\mathbb{R}^n \setminus B_{R_k})| \leq \int_{\mathbb{R}^n} w_k^2 dx.$$

Introduce the (relative) super-level measures

$$\mu_k := |\{u_\Omega > s_k\} \cap (\mathbb{R}^n \setminus B_{R_{k-1}})|, \quad k \geq 1,$$

which decrease in k (both s_k and R_{k-1} increase). Since $A_k \subset \{u_\Omega > s_k\} \cap (\mathbb{R}^n \setminus B_{R_{k-1}})$ we have $|A_k| \leq \mu_k$, and the left-hand set in (42) is exactly $\{u_\Omega > s_{k+1}\} \cap (\mathbb{R}^n \setminus B_{R_{(k+1)-1}})$, of measure μ_{k+1} . Combining (42), (41) and $|A_k| \leq \mu_k$,

$$(s_{k+1} - s_k)^2 \mu_{k+1} \leq C_2 4^k \mu_k^{1+2/n}.$$

Insert the gap (37), $(s_{k+1} - s_k)^2 = M^2 |\Omega \setminus B_R|^{2/n} 4^{-(k+1)}$:

$$(43) \quad \mu_{k+1} \leq \frac{C_2 4^k}{M^2 |\Omega \setminus B_R|^{2/n} 4^{-(k+1)}} \mu_k^{1+2/n} = \frac{4 C_2}{M^2 |\Omega \setminus B_R|^{2/n}} 16^k \mu_k^{1+2/n}.$$

Normalisation and the iteration lemma. Set $a_k := \mu_k / |\Omega \setminus B_R|$. Since $\mu_k \leq |\mathbb{R}^n \setminus B_{R_{k-1}} \cap \{u_\Omega > s_k\}| \leq |\Omega \setminus B_R|$ (because $R_{k-1} \geq R_0 = R$ and $\{u_\Omega > s_k\} \subset \Omega$), we have $0 \leq a_k \leq 1$; in particular

$a_1 \leq 1$. Dividing (43) by $|\Omega \setminus B_R|$ and using $\mu_k^{1+2/n} = |\Omega \setminus B_R|^{1+2/n} a_k^{1+2/n}$,

$$a_{k+1} \leq \frac{4C_2}{M^2 |\Omega \setminus B_R|^{2/n}} 16^k \frac{|\Omega \setminus B_R|^{1+2/n}}{|\Omega \setminus B_R|} a_k^{1+2/n} = \frac{4C_2}{M^2} 16^k a_k^{1+2/n}.$$

valid for all $k \geq 1$. To put this in the exact form (33) (indexed from 0) we shift: set $\tilde{a}_j := a_{j+1}$ for $j \geq 0$. Then for every $j \geq 0$,

$$\tilde{a}_{j+1} = a_{j+2} \leq \frac{4C_2}{M^2} 16^{j+1} a_{j+1}^{1+2/n} = \underbrace{\frac{64C_2}{M^2}}_{=:C} 16^j \tilde{a}_j^{1+2/n},$$

which is (33) with

$$C := \frac{64C_2}{M^2}, \quad b := 16, \quad \beta := \frac{2}{n}.$$

Choose $M = M(n)$ so large that the smallness condition (34) holds at the starting value $\tilde{a}_0 = a_1 \leq 1$. Since $a_1 \leq 1$ it suffices to require

$$1 \leq C^{-1/\beta} b^{-1/\beta^2} = \left(\frac{64C_2}{M^2}\right)^{-n/2} 16^{-n^2/4}, \quad \text{i.e.} \quad M^2 \geq 64C_2 16^{n^2/4 \cdot (2/n)} = 64C_2 16^{n/2},$$

which holds for $M := 8(C_2)^{1/2} 4^{n/2}$ (a dimensional constant, since $C_2 = C_2(n)$). With this M , Lemma 10.3 gives $\tilde{a}_j \rightarrow 0$, i.e. $a_k \rightarrow 0$ and hence $\mu_k \rightarrow 0$.

Conclusion. Letting $k \rightarrow \infty$ in $\mu_k \rightarrow 0$ and using $s_k \uparrow s_\infty = M |\Omega \setminus B_R|^{1/n}$ and $R_{k-1} \uparrow R+1$, monotone convergence yields

$$|\{u_\Omega > s_\infty\} \cap (\mathbb{R}^n \setminus B_{R+1})| = \lim_{k \rightarrow \infty} \mu_k = 0.$$

Hence $u_\Omega \leq s_\infty = M |\Omega \setminus B_R|^{1/n}$ a.e. on $\Omega \setminus B_{R+1}$. Since $u_\Omega \geq 0$, this is (36) with $C_b := M = 8(C_2)^{1/2} 4^{n/2}$. $\square$

Remark 10.6 (The case $n = 2$). For $n = 2$ the embedding $H_0^1 \hookrightarrow L^{2^*}$ degenerates, but the Gagliardo–Nirenberg–Sobolev inequality gives, for any fixed $p \in (2, \infty)$, $\|w\|_{L^p(\mathbb{R}^2)} \leq S_{2,p} \|\nabla w\|_{L^2(\mathbb{R}^2)}$ for $w \in H_0^1(\mathbb{R}^2)$ with finite-measure support. Repeating the Hölder step with exponent p replaces $2/n = 1$ by $1 - 2/p \in (0, 1)$, so (43) becomes $\mu_{k+1} \leq C 16^k \mu_k^{1+\beta}$ with $\beta := 1 - 2/p > 0$, and Lemma 10.3 applies. Tracking the level scale: with $s_\infty = M |\Omega \setminus B_R|^\alpha$, self-consistency of the recursion forces $\alpha = \frac{p-2}{2p} = \frac{1}{2} - \frac{1}{p}$ (for $n \geq 3$ the analogous bookkeeping gives $\alpha = 1/n$, and the two agree only there). Hence, for $n = 2$, the L^∞ tail estimate holds in the slightly weaker form

$$\|u_\Omega\|_{L^\infty(\Omega \setminus B_{R+1})} \leq C(p) |\Omega \setminus B_R|^{\frac{1}{2} - \frac{1}{p}}, \quad \text{any fixed } p \in (2, \infty),$$

in place of the exponent $1/n$ in (36). Fixing once and for all $p = 4$, so that $\frac{1}{2} - \frac{1}{p} = \frac{1}{4} > 0$, this is a genuine super-linear power of the tail mass, which is all the truncation of §10.5 uses: the energy-comparison (56) and the tail recursion (58) hold with the exponent $1 + \frac{1}{n}$ replaced by $1 + \kappa$, $\kappa = \frac{1}{4}$, and the geometric iteration (61)–(64) goes through for any $\kappa > 0$ with n -dependent constants. Consequently the global Theorem 10.2 holds for all $n \geq 2$; only the auxiliary exponent in (36) (and the resulting dimensional constants) changes at $n = 2$.

10.4. A suboptimal stability seed. The truncation iteration needs one quantitative input that the De Giorgi machinery does not by itself supply: a crude (far from sharp) control of the asymmetry by the deficit, used only to make the tail mass small at the first radius and to bound the number of truncation steps. We prove it here from the level-set identity already established, so the present section introduces no external stability input.

Lemma 10.7 (Suboptimal stability seed). *There is a dimensional constant $C_{\dagger} = C_{\dagger}(n) > 0$ such that every open $\Omega \subset \mathbb{R}^n$ of finite measure satisfies*

$$(44) \quad D(\Omega) \geq \frac{1}{C_{\dagger}} \mathcal{A}(\Omega)^4.$$

Equivalently, $\mathcal{A}(\Omega) \leq (C_{\dagger} D(\Omega))^{1/4}$.

Proof. Since D and $\mathcal{A}$ are scale invariant we may assume $|\Omega| = \omega_n$, so that $D(\Omega) = \omega_n^{-(n+2)/n} \delta_T(\Omega)$ and $B^* = B_1$; write $A := \mathcal{A}(\Omega) \in [0, 2)$ and recall $E_{\rho} = \{u > t(\rho)\}$, $|E_{\rho}| = \omega_n \rho^n$, $a(\rho) = -t'(\rho)$. We use three facts proved earlier in the manuscript, each of which holds for every open set of finite measure (their proofs do not use the confinement $\Omega \subset B_R$): the exact deficit identity (Theorem 3.11); the profile-gap identity (8), namely $0 \leq a(\rho) \leq \rho/n$ and $\int_0^1 \rho^n (\frac{\rho}{n} - a(\rho)) d\rho = \frac{2}{\omega_n} \delta_T(\Omega)$; and the superlevel transfer (Lemma 5.6), $A \leq \mathcal{A}(E_{\rho}) + 2(1 - \rho^n)$.

Step 1 (a level-set lower bound for D_I). Fix a finite-perimeter level ρ and apply Fusco–Julin (Theorem 5.1) to E_{ρ} with $r_E = \rho$; discarding the boundary-oscillation summand and the outer square, $|E_{\rho} \Delta B_{\rho}(z)|^2 \leq C_{\text{FJ}}(n) \rho^{n+1} (P(E_{\rho}) - P(B_{\rho}))$ for the optimal centre z . Since $\mathcal{A}(E_{\rho}) \leq |E_{\rho} \Delta B_{\rho}(z)| / |E_{\rho}|$ and $P(E_{\rho}) - P(B_{\rho}) \leq D_I(t(\rho)) / P(B_{\rho})$, with $|E_{\rho}| = \omega_n \rho^n$ and $P(B_{\rho}) = n \omega_n \rho^{n-1}$, this rearranges to

$$(45) \quad D_I(t(\rho)) \geq c_1(n) \rho^{2n-2} \mathcal{A}(E_{\rho})^2, \quad c_1(n) := \frac{n \omega_n^3}{C_{\text{FJ}}(n)}.$$

Step 2 (restrict to the outer collar). Put $\rho_A := (1 - A/4)^{1/n}$. For $\rho \in [\rho_A, 1]$ one has $1 - \rho^n \leq 1 - \rho_A^n = A/4$, so the superlevel transfer gives $\mathcal{A}(E_{\rho}) \geq A - 2(1 - \rho^n) \geq A/2$, and $\rho^n \geq \rho_A^n = 1 - A/4 > 1/2$. Drop $D_H \geq 0$ in Theorem 3.11 and change variables $t = t(\rho)$ ($dt = a(\rho) d\rho$, $m = \omega_n \rho^n$); keeping only $\rho \in [\rho_A, 1]$, where the integrand is nonnegative, and inserting (45),

$$2\delta_T(\Omega) \geq \int_{\rho_A}^1 \frac{D_I(t(\rho))}{n^2 \omega_n^{2/n} (\omega_n \rho^n)^{1-2/n}} a(\rho) d\rho \geq c_2(n) \int_{\rho_A}^1 \rho^n \mathcal{A}(E_{\rho})^2 a(\rho) d\rho,$$

where $c_2(n) = c_1(n) / (n^2 \omega_n)$ (using $(\omega_n \rho^n)^{1-2/n} = \omega_n^{1-2/n} \rho^{n-2}$ and $\rho^{2n-2} / \rho^{n-2} = \rho^n$). With $\mathcal{A}(E_{\rho}) \geq A/2$,

$$(46) \quad 2\delta_T(\Omega) \geq \frac{c_2(n) A^2}{4} \int_{\rho_A}^1 \rho^n a(\rho) d\rho.$$

Step 3 (the profile gap bounds the collar mass below). By the profile-gap identity and $\frac{\rho}{n} - a \geq 0$,

$$\int_{\rho_A}^1 \rho^n a d\rho = \int_{\rho_A}^1 \frac{\rho^{n+1}}{n} d\rho - \int_{\rho_A}^1 \rho^n \left(\frac{\rho}{n} - a \right) d\rho \geq \frac{1 - \rho_A^{n+2}}{n(n+2)} - \frac{2}{\omega_n} \delta_T(\Omega).$$

Since $\rho_A \leq 1$, $\rho_A^{n+2} \leq \rho_A^n = 1 - A/4$, so $1 - \rho_A^{n+2} \geq A/4$ and

$$(47) \quad \int_{\rho_A}^1 \rho^n a d\rho \geq \frac{A}{4n(n+2)} - \frac{2}{\omega_n} \delta_T(\Omega).$$

Step 4 (conclusion). Insert (47) into (46):

$$2\delta_T(\Omega) \geq \frac{c_2(n) A^2}{4} \left(\frac{A}{4n(n+2)} - \frac{2}{\omega_n} \delta_T(\Omega) \right) = c_3(n) A^3 - c_4(n) A^2 \delta_T(\Omega),$$

with $c_3(n) = c_2(n)/(16n(n+2))$ and $c_4(n) = c_2(n)/(2\omega_n)$. Hence $\delta_T(\Omega)(2+c_4(n)A^2) \geq c_3(n)A^3$, and since $A < 2$,

$$\delta_T(\Omega) \geq \frac{c_3(n)}{2+4c_4(n)} A^3 =: c_5(n) A^3.$$

Finally $A < 2$ gives $A^4 \leq 2A^3$, so $\delta_T(\Omega) \geq \frac{1}{2}c_5(n)A^4$; multiplying by $\omega_n^{-(n+2)/n}$ gives (44) with $C_{\dagger}(n) = 2\omega_n^{(n+2)/n}/c_5(n)$. $\square$

Remark 10.8 (The seed is internal). The only external ingredient in this proof is the Fusco–Julin quantitative isoperimetric inequality (Theorem 5.1), already used in the manuscript; everything else – the deficit identity, the profile-gap identity, and the superlevel transfer – is proved here. Thus the bounded reduction depends on no stability estimate beyond the ones the paper establishes. The exponent 4 (indeed the 3 obtained in the proof) is far from sharp – the sharp exponent 2 is the content of the main theorem – but only a fixed positive power of D is used below. A suboptimal torsional stability bound of this kind is classical; cf. Fusco–Maggi–Pratelli [6].

10.5. The truncation construction. We now build the bounded surrogate. The estimate driving the construction is an *energy-comparison inequality*: cutting Ω off at radius $\sim k$ costs only a higher-order amount of energy, controlled by the tail mass through Lemma 10.5.

Lemma 10.9 (Bounded truncation). *There exist dimensional constants $C_{\sharp} = C_{\sharp}(n) > 0$, $\delta = \delta(n) \in (0, 1]$ and $d = d(n) \geq 1$ such that for every open Ω with $|\Omega| = \omega_n$ and $D(\Omega) \leq \delta$ there is an open set $\tilde{\Omega}$ with*

$$(48) \quad |\tilde{\Omega}| = \omega_n,$$

$$(49) \quad \text{diam}(\tilde{\Omega}) \leq d,$$

$$(50) \quad D(\tilde{\Omega}) \leq C_{\sharp} D(\Omega),$$

$$(51) \quad \mathcal{A}(\Omega) \leq \mathcal{A}(\tilde{\Omega}) + C_{\sharp} D(\Omega).$$

Proof. Normalise so that the ball realising the asymmetry of Ω is B_1 (translate Ω ; both $\mathcal{A}$ and D are translation invariant), i.e. $\mathcal{A}(\Omega) = |\Omega \triangle B_1|/\omega_n$. We will repeatedly use the tail-mass notation

$$(52) \quad b_k := \frac{|\Omega \setminus B_k|}{\omega_n}, \quad k \geq 1,$$

which is non-increasing in k and satisfies $0 \leq b_k \leq 1$.

Step 0: the tail mass is small. Since B_1 realises the asymmetry, $|\Omega \setminus B_1| \leq |\Omega \triangle B_1| = \omega_n \mathcal{A}(\Omega)$. Combined with the suboptimal seed Lemma 10.7,

$$(53) \quad b_1 \leq \mathcal{A}(\Omega) \leq (C_{\dagger} D(\Omega))^{1/4}.$$

In particular, since $b_k \leq b_1$ for all $k \geq 1$, every tail mass is $\leq (C_{\dagger} D(\Omega))^{1/4}$, which can be made smaller than any prescribed dimensional threshold by shrinking $\delta(n)$ (recall $D(\Omega) \leq \delta(n)$). This is the only use of Lemma 10.7 in the construction.

Step 1: the energy-comparison inequality. For $k \geq 1$ let φ_k be the Lipschitz cut-off

$$\varphi_k(x) := \min\{1, (k+2-|x|)_+\},$$

so $\varphi_k \equiv 1$ on B_{k+1} , $\varphi_k \equiv 0$ outside B_{k+2} , $0 \leq \varphi_k \leq 1$ and $|\nabla \varphi_k| \leq 1$ everywhere (it is 1 only on the annulus $B_{k+2} \setminus B_{k+1}$). Then $u_k := \varphi_k u_{\Omega} \in H_0^1(\Omega \cap B_{k+2})$ is admissible for the variational

problem defining $E(\Omega \cap B_{k+2})$, so $E(\Omega \cap B_{k+2}) \leq J_{\Omega \cap B_{k+2}}(u_k) = \frac{1}{2} \int |\nabla u_k|^2 - \int u_k$. We use the pointwise algebraic identity

$$|\nabla(\varphi_k u_\Omega)|^2 = \nabla u_\Omega \cdot \nabla(u_\Omega \varphi_k^2) + |\nabla \varphi_k|^2 u_\Omega^2,$$

which is verified by expanding both sides ($\nabla(u_\Omega \varphi_k^2) = \varphi_k^2 \nabla u_\Omega + 2\varphi_k u_\Omega \nabla \varphi_k$, so the right-hand side equals $\varphi_k^2 |\nabla u_\Omega|^2 + 2\varphi_k u_\Omega \nabla u_\Omega \cdot \nabla \varphi_k + |\nabla \varphi_k|^2 u_\Omega^2 = |\varphi_k \nabla u_\Omega + u_\Omega \nabla \varphi_k|^2$). Integrating and applying the weak equation $\int \nabla u_\Omega \cdot \nabla(u_\Omega \varphi_k^2) = \int u_\Omega \varphi_k^2$ with the admissible test function $u_\Omega \varphi_k^2 \in H_0^1(\Omega)$, we get

$$J_{\Omega \cap B_{k+2}}(u_k) = \frac{1}{2} \int u_\Omega \varphi_k^2 + \frac{1}{2} \int |\nabla \varphi_k|^2 u_\Omega^2 - \int \varphi_k u_\Omega.$$

Now use $\frac{1}{2} \varphi_k^2 - \varphi_k = -\frac{1}{2}(1 - (1 - \varphi_k)^2) = -\frac{1}{2} + \frac{1}{2}(1 - \varphi_k)^2$ to write $\frac{1}{2} \int u_\Omega \varphi_k^2 - \int \varphi_k u_\Omega = -\frac{1}{2} \int u_\Omega + \frac{1}{2} \int (1 - \varphi_k)^2 u_\Omega = E(\Omega) + \frac{1}{2} \int (1 - \varphi_k)^2 u_\Omega$, where we used $E(\Omega) = -\frac{1}{2} \int u_\Omega$ from (29). Therefore

$$(54) \quad E(\Omega \cap B_{k+2}) \leq J_{\Omega \cap B_{k+2}}(u_k) = E(\Omega) + \frac{1}{2} \int (1 - \varphi_k)^2 u_\Omega + \frac{1}{2} \int |\nabla \varphi_k|^2 u_\Omega^2.$$

Both correction integrals are supported in $\mathbb{R}^n \setminus B_{k+1}$: indeed $1 - \varphi_k = 0$ on B_{k+1} and $\nabla \varphi_k = 0$ on $B_{k+1} \cup (\mathbb{R}^n \setminus B_{k+2})$. Hence, with $0 \leq 1 - \varphi_k \leq 1$ and $|\nabla \varphi_k| \leq 1$,

$$(55) \quad E(\Omega \cap B_{k+2}) - E(\Omega) \leq \frac{1}{2} |\Omega \setminus B_{k+1}| \left(\sup_{\Omega \setminus B_{k+1}} u_\Omega + \sup_{\Omega \setminus B_{k+1}} u_\Omega^2 \right).$$

By Lemma 10.5 applied with $R = k \geq 1$ (so $R + 1 = k + 1$ and the tail region is $\Omega \setminus B_{k+1}$), $\sup_{\Omega \setminus B_{k+1}} u_\Omega \leq C_b |\Omega \setminus B_k|^{1/n} = C_b \omega_n^{1/n} b_k^{1/n}$, and likewise the square is bounded by $C_b^2 \omega_n^{2/n} b_k^{2/n}$. Also $|\Omega \setminus B_{k+1}| = \omega_n b_{k+1} \leq \omega_n b_k$. Plugging in, and using $b_k \leq 1$ so that $b_k^{2/n} \leq b_k^{1/n}$,

$$(56) \quad E(\Omega \cap B_{k+2}) \leq E(\Omega) + C b_{k+1} (b_k^{1/n} + b_k^{2/n}) \leq E(\Omega) + C b_k^{1+1/n},$$

for a dimensional constant $C = C(n)$ (we absorbed $\frac{1}{2} \omega_n (C_b \omega_n^{1/n} + C_b^2 \omega_n^{2/n})$ and $b_{k+1} \leq b_k$ into C). This is the energy-comparison inequality.

Step 2: turning energy comparison into a tail recursion. By the Saint-Venant inequality (31) applied to the set $\Omega \cap B_{k+2}$, whose volume is $|\Omega \cap B_{k+2}| = \omega_n(1 - b_{k+2})$,

$$E(B_1) |B_1|^{-(n+2)/n} \leq E(\Omega \cap B_{k+2}) |\Omega \cap B_{k+2}|^{-(n+2)/n},$$

that is

$$E(B_1) \omega_n^{-(n+2)/n} (\omega_n(1 - b_{k+2}))^{(n+2)/n} \leq E(\Omega \cap B_{k+2}),$$

i.e.

$$E(B_1) (1 - b_{k+2})^{(n+2)/n} \leq E(\Omega \cap B_{k+2}).$$

Combine with (56) and the definition $E(\Omega) = E(B_1) + \omega_n^{(n+2)/n} D(\Omega)$ (which is (30) at $|\Omega| = \omega_n$):

$$E(B_1) (1 - b_{k+2})^{(n+2)/n} \leq E(\Omega) + C b_k^{1+1/n} = E(B_1) + \omega_n^{(n+2)/n} D(\Omega) + C b_k^{1+1/n}.$$

Rearranging, and recalling $E(B_1) < 0$, divide by $-E(B_1) > 0$:

$$1 - (1 - b_{k+2})^{(n+2)/n} \leq \frac{\omega_n^{(n+2)/n} D(\Omega) + C b_k^{1+1/n}}{-E(B_1)} =: \widehat{C}_1 (D(\Omega) + b_k^{1+1/n}),$$

with $\widehat{C}_1 = \widehat{C}_1(n) > 0$ (here $\widehat{C}_1 = \max\{\omega_n^{(n+2)/n}, C\}/(-E(B_1))$). To extract b_{k+2} we use the elementary lower bound

$$(57) \quad 1 - (1 - t)^p \geq t \quad \text{for all } t \in [0, 1], \quad p \geq 1,$$

which holds because $r \mapsto (1-t)^r$ is non-increasing on $[0, \infty)$, so $(1-t)^p \leq (1-t)^1 = 1-t$ for $p = (n+2)/n \geq 1$. Applying (57) with $t = b_{k+2} \in [0, 1]$ and $p = (n+2)/n$,

$$(58) \quad b_{k+2} \leq 1 - (1 - b_{k+2})^{(n+2)/n} \leq \widehat{C} (D(\Omega) + b_k^{1+1/n}), \quad \widehat{C} := \widehat{C}_1.$$

Step 3: the geometric iteration to a bounded stopping index. Define the stopping index

$$(59) \quad K := \max\{k \geq 1 : b_k \geq 2\widehat{C} D(\Omega)\},$$

with the convention $K := 0$ if $b_1 < 2\widehat{C} D(\Omega)$ (then the tail is already controlled at the first radius and Step 5 applies with $K = 0$). The set in (59) is finite because $b_k \rightarrow 0$ as $k \rightarrow \infty$ (the tail mass of a finite-measure set vanishes), so K is well defined. We prove

$$(60) \quad K \leq K(n)$$

for a dimensional $K(n)$, provided $\delta(n)$ is small enough. Assume $K \geq 3$ (otherwise (60) is trivial).

Self-improving contraction. For every k with $k+2 \leq K$ we have, by definition of K , that $b_{k+2} \geq 2\widehat{C} D(\Omega)$, i.e. $\widehat{C} D(\Omega) \leq \frac{1}{2} b_{k+2}$. Feeding this into (58),

$$b_{k+2} \leq \widehat{C} D(\Omega) + \widehat{C} b_k^{1+1/n} \leq \frac{1}{2} b_{k+2} + \widehat{C} b_k^{1+1/n},$$

and absorbing $\frac{1}{2} b_{k+2}$ on the left,

$$(61) \quad b_{k+2} \leq 2\widehat{C} b_k^{1+1/n} \quad \text{whenever } k+2 \leq K.$$

Smallness from the seed. By Step 0, $b_1 \leq (C_\dagger D(\Omega))^{1/4} \leq (C_\dagger \delta(n))^{1/4}$. Choose $\delta(n)$ small enough that

$$(62) \quad b_1 \leq (C_\dagger \delta(n))^{1/4} \leq (2\widehat{C})^{-2n} =: \eta_n \in (0, 1).$$

Then $2\widehat{C} b_k^{1/(2n)} \leq 2\widehat{C} \eta_n^{1/(2n)} = 1$ for all $k \geq 1$ (as $b_k \leq b_1 \leq \eta_n$), so (61) self-improves to

$$(63) \quad b_{k+2} = (2\widehat{C} b_k^{1/(2n)}) b_k^{1+1/(2n)} \leq b_k^{1+1/(2n)} \quad \text{whenever } k+2 \leq K.$$

Iterating along odd indices. Apply (63) repeatedly, starting from $k = 1$, as long as the index stays $\leq K$. Writing the odd indices $1, 3, 5, \dots$ and setting $\gamma := 1 + \frac{1}{2n} > 1$, induction gives, for every $j \geq 0$ with $2j+1 \leq K$,

$$(64) \quad b_{2j+1} \leq b_1^{\gamma^j}.$$

Indeed (64) holds at $j = 0$, and if it holds at j with $2(j+1)+1 = 2j+3 \leq K$, then (63) (at $k = 2j+1$, legitimate since $2j+3 \leq K$) gives $b_{2j+3} \leq b_{2j+1}^\gamma \leq (b_1^{\gamma^j})^\gamma = b_1^{\gamma^{j+1}}$.

Bounding K . Let $j_K := \lfloor (K-1)/2 \rfloor$, the largest j with $2j+1 \leq K$. By the maximality of K and (64),

$$(65) \quad 2\widehat{C} D(\Omega) \leq b_{2j_K+1} \leq b_1^{\gamma^{j_K}} \leq (C_\dagger D(\Omega))^{\gamma^{j_K}/4},$$

using $b_1 \leq (C_\dagger D(\Omega))^{1/4}$. Take logarithms (recall $D(\Omega) \leq \delta(n) < 1$, so $\log D(\Omega) < 0$); writing $L := -\log D(\Omega) > 0$, (65) reads

$$\log(2\widehat{C}) - L \leq \frac{\gamma^{j_K}}{4} (\log C_\dagger - L).$$

For $\delta(n)$ small, $L = -\log D(\Omega) \geq -\log \delta(n)$ is large; in particular we may take $\delta(n)$ small enough that $L \geq 2 \max\{|\log(2\widehat{C})|, |\log C_\dagger|\}$, whence $\log(2\widehat{C}) - L \geq -\frac{3}{2}L$ and $\log C_\dagger - L \leq -\frac{1}{2}L$.

The displayed inequality then forces $-\frac{3}{2}L \leq -\frac{\gamma^{j_K}}{8}L$, i.e. $\gamma^{j_K} \leq 12$. Since $\gamma = 1 + \frac{1}{2n} > 1$ this

bounds $j_K \leq \log 12 / \log \gamma =: j_*(n)$, hence $K \leq 2j_K + 2 \leq 2j_*(n) + 2 =: K(n)$, a dimensional constant. This proves (60).

Step 4: tail at the stopping radius. By the maximality (59), $b_{K+1} < 2\widehat{C}D(\Omega)$; we record this for the next step.

Step 5: the truncated set and its volume. By the definition (59) of K and $K \leq K(n)$, $b_{K+1} < 2\widehat{C}D(\Omega)$, hence

$$(66) \quad |\Omega \setminus B_{K+1}| = \omega_n b_{K+1} < 2\widehat{C}\omega_n D(\Omega).$$

In fact we cut one radius further out to have room for the cut-off: by monotonicity $b_{K+3} \leq b_{K+1} < 2\widehat{C}D(\Omega)$, so

$$(67) \quad |\Omega \cap B_{K+3}| = \omega_n(1 - b_{K+3}) \geq \omega_n(1 - 2\widehat{C}D(\Omega)), \quad |\Omega \setminus B_{K+3}| < 2\widehat{C}\omega_n D(\Omega).$$

Define

$$r := \left(\frac{|\Omega \cap B_{K+3}|}{\omega_n} \right)^{1/n} = (1 - b_{K+3})^{1/n} \in [(1 - 2\widehat{C}D(\Omega))^{1/n}, 1], \quad \widetilde{\Omega} := r^{-1}(\Omega \cap B_{K+3}).$$

Then $|\widetilde{\Omega}| = r^{-n}|\Omega \cap B_{K+3}| = \omega_n$, which is (48). Since $\Omega \cap B_{K+3} \subset B_{K+3}$ and $r \geq (1 - 2\widehat{C}\delta(n))^{1/n} \geq \frac{1}{2}$ for $\delta(n)$ small,

$$\text{diam}(\widetilde{\Omega}) = r^{-1} \text{diam}(\Omega \cap B_{K+3}) \leq r^{-1} \cdot 2(K+3) \leq 4(K(n)+3) =: d(n),$$

giving (49). We record for later the linearised bound

$$(68) \quad 0 \leq 1 - r^n = b_{K+3} < 2\widehat{C}D(\Omega), \quad 0 \leq 1 - r \leq 1 - r^n < 2\widehat{C}D(\Omega),$$

the middle inequality because $r \leq 1$ implies $1 - r \leq 1 - r^n$.

Step 6: deficit of the surrogate. Apply the energy comparison (56) at $k = K+1$ (so the cut radius is $k+2 = K+3$):

$$E(\Omega \cap B_{K+3}) \leq E(\Omega) + C b_{K+1}^{1+1/n} \leq E(\Omega) + C(2\widehat{C}D(\Omega))^{1+1/n} \leq E(\Omega) + C'D(\Omega),$$

where in the last step we used $D(\Omega) \leq \delta(n) \leq 1$ so that $D(\Omega)^{1+1/n} \leq D(\Omega)$, and absorbed constants into $C' = C'(n)$. Since E scales as $E(t\Omega') = t^{n+2}E(\Omega')$ and $\widetilde{\Omega} = r^{-1}(\Omega \cap B_{K+3})$,

$$E(\widetilde{\Omega}) = r^{-(n+2)}E(\Omega \cap B_{K+3}) \leq r^{-(n+2)}(E(\Omega) + C'D(\Omega)).$$

Because $|\widetilde{\Omega}| = \omega_n$, $D(\widetilde{\Omega}) = \omega_n^{-(n+2)/n}(E(\widetilde{\Omega}) - E(B_1))$. Using $E(\Omega) = E(B_1) + \omega_n^{(n+2)/n}D(\Omega)$ and $r \leq 1$ (so $r^{-(n+2)} \geq 1$), with $E(B_1) < 0$,

$$\begin{aligned} E(\widetilde{\Omega}) - E(B_1) &\leq r^{-(n+2)}(E(B_1) + \omega_n^{(n+2)/n}D(\Omega) + C'D(\Omega)) - E(B_1) \\ &= (r^{-(n+2)} - 1)E(B_1) + r^{-(n+2)}(\omega_n^{(n+2)/n} + C')D(\Omega). \end{aligned}$$

For the first term, $r^{-(n+2)} - 1 \geq 0$ and $E(B_1) < 0$, so $(r^{-(n+2)} - 1)E(B_1) \leq 0$ and we may bound it by $(r^{-(n+2)} - 1)|E(B_1)|$. Quantitatively, by (68), $r^n \geq 1 - 2\widehat{C}D(\Omega)$, so $r^{-(n+2)} = r^{-n} \cdot r^{-2} \leq (1 - 2\widehat{C}D(\Omega))^{-(n+2)/n} \leq 1 + C''D(\Omega)$ for $\delta(n)$ small (Bernoulli/Taylor bound, since $2\widehat{C}D(\Omega) \leq 2\widehat{C}\delta(n) \leq \frac{1}{2}$). Hence $r^{-(n+2)} - 1 \leq C''D(\Omega)$ and $r^{-(n+2)} \leq 1 + C''D(\Omega) \leq 2$, giving

$$E(\widetilde{\Omega}) - E(B_1) \leq C''D(\Omega) \cdot |E(B_1)| + 2(\omega_n^{(n+2)/n} + C')D(\Omega) \leq C'_2 D(\Omega),$$

with $C'_2 = C'_2(n)$. Dividing by $\omega_n^{(n+2)/n}$,

$$(69) \quad D(\widetilde{\Omega}) = \omega_n^{-(n+2)/n}(E(\widetilde{\Omega}) - E(B_1)) \leq \omega_n^{-(n+2)/n}C'_2 D(\Omega) =: C_3 D(\Omega),$$

which is (50) with $C_3 = C_3(n)$.

Step 7: asymmetry loss. Let $B_1(x_0)$ realise the asymmetry of $\tilde{\Omega}$, so $\mathcal{A}(\tilde{\Omega}) = |\tilde{\Omega} \triangle B_1(x_0)| / \omega_n$. We compare the asymmetry of Ω to that of $\tilde{\Omega}$ through the chain of inclusions relating Ω , $\Omega \cap B_{K+3}$ and $\tilde{\Omega} = r^{-1}(\Omega \cap B_{K+3})$. For any test ball $B_1(y)$,

$$|\Omega \triangle B_1(y)| \leq |\Omega \triangle (\Omega \cap B_{K+3})| + |(\Omega \cap B_{K+3}) \triangle B_1(y)|.$$

The first term is $|\Omega \setminus B_{K+3}| < 2\hat{C}\omega_n D(\Omega)$ by (67). For the second, scale by r : writing $E_r := \Omega \cap B_{K+3} = r\tilde{\Omega}$ and $B_1(y) = r B_{1/r}(y/r)$,

$$|(r\tilde{\Omega}) \triangle (r B_{1/r}(y/r))| = r^n |\tilde{\Omega} \triangle B_{1/r}(y/r)| \leq r^n \left(|\tilde{\Omega} \triangle B_1(y/r)| + |B_1(y/r) \triangle B_{1/r}(y/r)| \right).$$

The last term is the volume difference of concentric balls of radii 1 and $1/r \geq 1$: $|B_1 \triangle B_{1/r}| = \omega_n(r^{-n} - 1)$. Hence

$$r^n |B_1(y/r) \triangle B_{1/r}(y/r)| = r^n \omega_n(r^{-n} - 1) = \omega_n(1 - r^n) < 2\hat{C}\omega_n D(\Omega),$$

using (68). Choosing $y := rx_0$ so that $y/r = x_0$ is the optimal centre for $\tilde{\Omega}$, the middle term becomes $r^n |\tilde{\Omega} \triangle B_1(x_0)| = r^n \omega_n \mathcal{A}(\tilde{\Omega}) \leq \omega_n \mathcal{A}(\tilde{\Omega})$ (as $r \leq 1$). Collecting,

$$|\Omega \triangle B_1(rx_0)| \leq 2\hat{C}\omega_n D(\Omega) + \omega_n \mathcal{A}(\tilde{\Omega}) + 2\hat{C}\omega_n D(\Omega),$$

and dividing by ω_n and taking the infimum over translates (which only decreases the left side),

$$(70) \quad \mathcal{A}(\Omega) \leq \frac{|\Omega \triangle B_1(rx_0)|}{\omega_n} \leq \mathcal{A}(\tilde{\Omega}) + 4\hat{C} D(\Omega).$$

This is (51) with the constant $4\hat{C}$. Taking $C_\sharp := \max\{C_3, 4\hat{C}\} = C_\sharp(n)$ proves (50)–(51) simultaneously, and we already have (48)–(49). This completes the proof. $\square$

Remark 10.10 (Translation into a fixed ball). The diameter bound (49) is translation invariant, so after a translation we may assume $\tilde{\Omega} \subset B_{d(n)}$. Both $\mathcal{A}$ and D are translation invariant, so (48)–(51) are preserved.

10.6. The far branch.

Lemma 10.11 (Large-deficit quadratic bound). *For every open Ω with $|\Omega| = \omega_n$ and $D(\Omega) \geq \delta(n)/4$,*

$$(71) \quad E(\Omega) - E(B_1) \geq \omega_n^{(n+2)/n} \frac{\delta(n)}{16} \mathcal{A}(\Omega)^2.$$

Proof. Since $|\Omega| = \omega_n$ gives $\mathcal{A}(\Omega) < 2$, we have $\mathcal{A}(\Omega)^2/4 < 1$. Hence

$$D(\Omega) \geq \frac{\delta(n)}{4} = \frac{\delta(n)}{4} \cdot 1 \geq \frac{\delta(n)}{4} \cdot \frac{\mathcal{A}(\Omega)^2}{4} = \frac{\delta(n)}{16} \mathcal{A}(\Omega)^2.$$

Multiplying by $\omega_n^{(n+2)/n}$ and using $E(\Omega) - E(B_1) = \omega_n^{(n+2)/n} D(\Omega)$ gives (71). $\square$

10.7. The near branch. Fix the dimensional working radius

$$(72) \quad R_*(n) := \max\{d(n), 2\},$$

so $R_* \geq d(n)$ (the surrogates of Lemma 10.9 lie in B_{R_*} after translation, by Remark 10.10) and $R_* \geq 2$ (the hypothesis of Theorem 10.1).

Lemma 10.12 (Small-deficit transfer). *Set*

$$(73) \quad \tilde{c}_{SV}(n) := \omega_n^{-(n+2)/n} c_{SV}(n, R_*(n)) > 0.$$

For every open Ω with $|\Omega| = \omega_n$ and $D(\Omega) \leq \delta(n)$,

$$(74) \quad \mathcal{A}(\Omega)^2 \leq \left(\frac{4C_\#(n)}{\tilde{c}_{\text{SV}}(n)} + 4C_\#(n)^2 \right) D(\Omega).$$

Proof. Apply Lemma 10.9 to get $\tilde{\Omega}$ satisfying (48)–(51); by Remark 10.10 assume $\tilde{\Omega} \subset B_{d(n)} \subset B_{R_*(n)}$. Theorem 10.1 at $R = R_*(n)$ gives $E(\tilde{\Omega}) - E(B_1) \geq c_{\text{SV}}(n, R_*) \mathcal{A}(\tilde{\Omega})^2$; dividing by $\omega_n^{(n+2)/n}$ and using $|\tilde{\Omega}| = \omega_n$,

$$(75) \quad D(\tilde{\Omega}) \geq \tilde{c}_{\text{SV}}(n) \mathcal{A}(\tilde{\Omega})^2.$$

Combining (75) with (50),

$$\mathcal{A}(\tilde{\Omega})^2 \leq \frac{D(\tilde{\Omega})}{\tilde{c}_{\text{SV}}} \leq \frac{C_\#}{\tilde{c}_{\text{SV}}} D(\Omega).$$

Now use (51), the elementary $(a+b)^2 \leq 2a^2 + 2b^2$, and $D(\Omega) \leq \delta(n) \leq 1$ (so $D(\Omega)^2 \leq D(\Omega)$):

$$\begin{aligned} \mathcal{A}(\Omega)^2 &\leq (\mathcal{A}(\tilde{\Omega}) + C_\# D(\Omega))^2 \leq 2\mathcal{A}(\tilde{\Omega})^2 + 2C_\#^2 D(\Omega)^2 \\ &\leq \frac{2C_\#}{\tilde{c}_{\text{SV}}} D(\Omega) + 2C_\#^2 D(\Omega) \leq \left(\frac{4C_\#}{\tilde{c}_{\text{SV}}} + 4C_\#^2 \right) D(\Omega), \end{aligned}$$

which is (74). $\square$

Corollary 10.13 (Small-deficit quadratic bound). *With*

$$(76) \quad c_{\text{near}}(n) := \frac{\omega_n^{(n+2)/n}}{4C_\#(n)(1/\tilde{c}_{\text{SV}}(n) + C_\#(n))} > 0,$$

every open Ω with $|\Omega| = \omega_n$ and $D(\Omega) \leq \delta(n)$ satisfies $E(\Omega) - E(B_1) \geq c_{\text{near}}(n) \mathcal{A}(\Omega)^2$.

Proof. Rearrange (74) to $D(\Omega) \geq \mathcal{A}(\Omega)^2 / (4C_\#/\tilde{c}_{\text{SV}} + 4C_\#^2)$ and multiply by $\omega_n^{(n+2)/n}$, using $E(\Omega) - E(B_1) = \omega_n^{(n+2)/n} D(\Omega)$. $\square$

10.8. Proof of the global theorem.

Proof of Theorem 10.2. Set

$$(77) \quad c_{\text{SV}}^{\text{glob}}(n) := \min \left\{ c_{\text{near}}(n), \omega_n^{(n+2)/n} \frac{\delta(n)}{16} \right\} > 0.$$

Let Ω be open with $|\Omega| = \omega_n$.

Case A: $D(\Omega) \geq \delta(n)/4$. Lemma 10.11 gives $E(\Omega) - E(B_1) \geq \omega_n^{(n+2)/n} (\delta(n)/16) \mathcal{A}(\Omega)^2 \geq c_{\text{SV}}^{\text{glob}}(n) \mathcal{A}(\Omega)^2$.

Case B: $D(\Omega) \leq \delta(n)$ (which includes $D(\Omega) \leq \delta(n)/4$). Corollary 10.13 gives $E(\Omega) - E(B_1) \geq c_{\text{near}}(n) \mathcal{A}(\Omega)^2 \geq c_{\text{SV}}^{\text{glob}}(n) \mathcal{A}(\Omega)^2$.

The two cases overlap on $\delta(n)/4 \leq D(\Omega) \leq \delta(n)$ and together cover $[0, \infty)$; in either case (26) holds with $c_{\text{SV}}^{\text{glob}}(n)$ as in (77), a dimensional constant because $\delta(n)$, $c_{\text{near}}(n)$ depend only on n (recall $R_*(n)$ and hence $c_{\text{SV}}(n, R_*(n))$ are dimensional).

General volume. For arbitrary finite-measure Ω put $\Omega_* := (\omega_n/|\Omega|)^{1/n} \Omega$, so $|\Omega_*| = \omega_n$, $\mathcal{A}(\Omega_*) = \mathcal{A}(\Omega)$ and $D(\Omega_*) = D(\Omega)$ by scale invariance of $\mathcal{A}$ and D . Applying (26) to Ω_* reads $D(\Omega) \geq \omega_n^{-(n+2)/n} c_{\text{SV}}^{\text{glob}}(n) \mathcal{A}(\Omega)^2$. Multiplying by $|\Omega|^{(n+2)/n}$ and using $E(\Omega) |\Omega|^{-(n+2)/n} - E(B^*) |B^*|^{-(n+2)/n} = D(\Omega)$ with $|B^*| = |\Omega|$, then $E = -T/2$, gives

$$T(B^*) - T(\Omega) = 2(E(\Omega) - E(B^*)) \geq 2c_{\text{SV}}^{\text{glob}}(n) \left(\frac{|\Omega|}{\omega_n} \right)^{(n+2)/n} \mathcal{A}(\Omega)^2,$$

which is (27). $\square$

Remark 10.14 (Sharpness of the exponent). The exponent 2 on $\mathcal{A}$ in (26) is optimal: it is saturated by smooth, volume-preserving ellipsoidal perturbations of B_1 , for which both $E(\Omega) - E(B_1)$ and $\mathcal{A}(\Omega)^2$ are of the same quadratic order in the perturbation parameter.

11. THE KOHLER–JOBIN INEQUALITY AND THE TRANSFER TO FABER–KRAHN

This section is logically self-contained. In Part 1 we reconstruct from scratch, by the Kohler–Jobin rearrangement technique, the spectral shape inequality

$$T(\Omega) \lambda_1(\Omega)^{(n+2)/2} \geq T(B) \lambda_1(B)^{(n+2)/2} \quad (B \text{ any ball}),$$

with equality only for balls. In Part 2 we feed this inequality, together with the global sharp Saint–Venant stability estimate proved elsewhere in the manuscript, into a short and fully explicit computation that yields sharp quantitative Faber–Krahn stability, $\delta_{\text{FK}}(\Omega) \geq c_{\text{FK}}(n) \mathcal{A}(\Omega)^2$.

The technique, due to Kohler–Jobin [7, 8, 9], is a spherical rearrangement that, unlike Schwarz symmetrisation, preserves not the *volume* of the superlevel sets but a *torsional* functional of them. We follow the modern exposition of Brasco [4, §§3–5, Thm. 1.1], specialised throughout to the linear case (the Laplacian, $p = 2$, and the Dirichlet energy quotient).

11.1. Notation and scaling conventions. Throughout, $n \geq 2$ and $\Omega \subset \mathbb{R}^n$ is open with finite measure. We use:

- $u_\Omega \in H_0^1(\Omega)$, the *torsion function*, the unique weak solution of $-\Delta u_\Omega = 1$ in Ω , $u_\Omega = 0$ on $\partial\Omega$;
- $T(\Omega) = \int_\Omega u_\Omega dx$, the *torsional rigidity*, and $E(\Omega) = -T(\Omega)/2$;
- $\lambda_1(\Omega)$, the first Dirichlet eigenvalue,

$$(78) \quad \lambda_1(\Omega) = \min_{0 \neq v \in H_0^1(\Omega)} \frac{\int_\Omega |\nabla v|^2 dx}{\int_\Omega v^2 dx},$$

attained by a first eigenfunction $u > 0$ solving $-\Delta u = \lambda_1(\Omega)u$;

- B^* , the ball centred at 0 with $|B^*| = |\Omega|$; B_r the open ball of radius r centred at 0; B_1 the unit ball; $\omega_n = |B_1|$;
- $L_n = \lambda_1(B_1) = j_{n/2-1,1}^2$ and $T_n = T(B_1) = \omega_n/(n(n+2))$;
- $\mathcal{A}(\Omega) = \inf_{x_0 \in \mathbb{R}^n} |\Omega \triangle (x_0 + B^*)| / |B^*| \in [0, 2)$, the Fraenkel asymmetry;
- $\delta_{\text{FK}}(\Omega) = (|\Omega|/\omega_n)^{2/n} (\lambda_1(\Omega) - \lambda_1(B^*))$, the scale-invariant Faber–Krahn deficit.

The torsion function of a ball is $u_{B_r}(x) = (r^2 - |x|^2)/(2n)$, whence

$$(79) \quad T(B_r) = \frac{\omega_n}{n(n+2)} r^{n+2} = T_n r^{n+2}.$$

Under dilation $T(t\Omega) = t^{n+2}T(\Omega)$ and $\lambda_1(t\Omega) = t^{-2}\lambda_1(\Omega)$, so

$$(80) \quad \Psi(\Omega) := T(\Omega) \lambda_1(\Omega)^{(n+2)/2}$$

is scale invariant; in particular $\Psi(B_r) = \Psi(B_1) = T_n L_n^{(n+2)/2}$ for every ball. We abbreviate $\beta := (n+2)/2$.

11.2. Part 1: the Kohler–Jobin inequality, proved from scratch.

Theorem 11.1 (Kohler–Jobin). *For every open $\Omega \subset \mathbb{R}^n$ of finite measure with $\lambda_1(\Omega) < \infty$ and every ball $B \subset \mathbb{R}^n$,*

$$(81) \quad T(\Omega) \lambda_1(\Omega)^\beta \geq T(B) \lambda_1(B)^\beta, \quad \beta = \frac{n+2}{2}.$$

Equivalently $\Psi(\Omega) \geq \Psi(B_1) = T_n L_n^\beta$, with equality if and only if Ω is a ball (up to translation and a Lebesgue-null modification).

Strategy. Both sides of (81) have the same scaling and $\Psi(B) = \Psi(B_1)$ for every ball; so it suffices to compare Ω with a single, well-chosen ball. The idea is to rearrange the first eigenfunction u of Ω into a radial decreasing function u^* on a ball B in such a way that

- (A) the Dirichlet energy is *exactly preserved*, $\int_B |\nabla u^*|^2 = \int_\Omega |\nabla u|^2$, and
- (B) the L^2 norm *does not decrease*, $\int_B (u^*)^2 \geq \int_\Omega u^2$.

Granting (A)–(B), the Rayleigh principle (78) on B gives

$$(82) \quad \lambda_1(B) \leq \frac{\int_B |\nabla u^*|^2}{\int_B (u^*)^2} \leq \frac{\int_\Omega |\nabla u|^2}{\int_\Omega u^2} = \lambda_1(\Omega),$$

the last equality because u is the first eigenfunction of Ω .

The decisive design choice is which geometric quantity of the superlevel sets the ball B is built to match. Schwarz symmetrisation matches *volume* and yields the Faber–Krahn inequality directly; the Kohler–Jobin rearrangement instead matches a *torsional functional*, the *modified torsional rigidity* introduced next, and this is exactly what allows the Dirichlet energy to be preserved (not merely decreased) while the L^2 norm grows. The relation $\lambda_1(B) \leq \lambda_1(\Omega)$ at equal *torsion* of B and Ω is then the inequality (81).

Standing regularity facts (cited, not proved).

- (R1) (*Regularity of the eigenfunction.*) The first eigenfunction u of Ω may be taken positive, solves $-\Delta u = \lambda u$ in Ω with $\lambda = \lambda_1(\Omega)$, and satisfies $u \in L^\infty(\Omega)$ (Moser iteration) and $u \in C^\infty(\Omega)$ in the interior: indeed $-\Delta u = \lambda u$ with $u \in L^\infty$ gives $u \in C_{\text{loc}}^{1,\alpha}$ by Schauder, and bootstrapping the equation gives $u \in C^\infty(\Omega)$ ([4, Prop. 2.4], [13]). In particular Sard’s theorem applies to u in the interior, so a.e. $t \in (0, M)$ is a regular value. Put $M := \|u\|_{L^\infty(\Omega)} < \infty$.
- (R2) (*Coarea formula.*) For $u \in W_{\text{loc}}^{1,1}$ and Borel $g \geq 0$, $\int_\Omega g |\nabla u| \, dx = \int_0^M \left(\int_{\{u=t\}} g \, d\mathcal{H}^{n-1} \right) dt$ (Federer; see Maggi, *Sets of Finite Perimeter*, Thm. 13.1).
- (R3) (*Saint–Venant inequality.*) For any open bounded $A \subset \mathbb{R}^n$, $T(A) \leq T(A^\#)$ where $A^\#$ is the ball with $|A^\#| = |A|$; in the scale-free form we use it,

$$(83) \quad T(A) \leq T_n \left(\frac{|A|}{\omega_n} \right)^{(n+2)/n},$$

with equality iff A is a ball. ([4, (1.7) with $p = 2$].)

- (R4) (*Distribution function.*) The distribution function $\mu_u(t) := |\{u > t\}|$ is non-increasing; together with the coarea formula this gives, for a.e. regular value t , that $\int_{\{u=t\}} |\nabla u|^{-1} \, d\mathcal{H}^{n-1}$ is finite and $\mu'_u(t) = - \int_{\{u=t\}} |\nabla u|^{-1} \, d\mathcal{H}^{n-1}$ (Brothers–Ziemer).

Two preliminary identities. We record the variational form of T and the level-set flux identity for the eigenfunction; both are elementary.

Lemma 11.2 (Variational torsion and the eigenfunction flux).

- (i) For open A of finite measure, with $F(w) := \int_A w \, dx - \frac{1}{2} \int_A |\nabla w|^2 \, dx$,

$$(84) \quad T(A) = 2 \max_{w \in H_0^1(A)} F(w),$$

the maximiser being the torsion function u_A ; equivalently $T(A) = \max\{(\int_A w)^2 / \int_A |\nabla w|^2 : 0 \neq w \in H_0^1(A)\}$.

(ii) The first eigenfunction u satisfies, for a.e. $t \in (0, M)$,

$$(85) \quad \int_{\{u=t\}} |\nabla u| d\mathcal{H}^{n-1} = \lambda \int_{\{u>t\}} u dx.$$

Proof. (i): F is strictly concave and coercive on $H_0^1(A)$; its Euler equation is $-\Delta w = 1$, so the maximiser is u_A and $F(u_A) = \int u_A - \frac{1}{2} \int |\nabla u_A|^2 = \int u_A - \frac{1}{2} \int u_A = \frac{1}{2} \int u_A = \frac{1}{2} T(A)$, using $\int |\nabla u_A|^2 = \int u_A$ (test the equation by u_A). The second form is the standard Rayleigh-type characterisation of T ([4, Prop. 2.2] with $p = 2$).

(ii): integrate $-\Delta u = \lambda u$ over the (a.e. smooth) superlevel set $\{u > t\}$ and use the divergence theorem with outward normal $\nu = -\nabla u / |\nabla u|$ on $\partial\{u > t\} = \{u = t\}$: $\int_{\{u=t\}} |\nabla u| d\mathcal{H}^{n-1} = -\int_{\{u=t\}} \partial_\nu u d\mathcal{H}^{n-1} = \int_{\{u>t\}} (-\Delta u) dx = \lambda \int_{\{u>t\}} u dx$. $\square$

The modified torsional rigidity. The engine of the technique is the following functional, which assigns to a nonnegative function a torsion-type quantity that is sensitive to the *level structure* of the function, not just to its domain.

Definition 11.3 (Modified torsional rigidity). Let $u \in H_0^1(\Omega) \cap L^\infty(\Omega)$, $u \geq 0$, $M = \|u\|_\infty$, with distribution function $\mu(t) = |\{u > t\}|$. Assume the integrability condition

$$(86) \quad t \mapsto \frac{\mu(t)}{\int_{\{u=t\}} |\nabla u| d\mathcal{H}^{n-1}} \in L^\infty(0, M)$$

For the first eigenfunction this holds: by (85) the ratio in (86) equals $\mu(t)/(\lambda \int_{\{u>t\}} u)$. On $[M/2, M)$ the bound $\int_{\{u>t\}} u \geq t\mu(t)$ gives ratio $\leq 1/(\lambda t) \leq 2/(\lambda M)$; on $(0, M/2]$ the monotonicity $\int_{\{u>t\}} u \geq \int_{\{u>M/2\}} u > 0$ together with $\mu(t) \leq |\Omega|$ gives ratio $\leq |\Omega|/(\lambda \int_{\{u>M/2\}} u)$. Hence the ratio is bounded on $(0, M)$ (cf. [4, Lem. 3.2]). With $\mathcal{L} := \{g \in \text{Lip}([0, M]) : g(0) = 0\}$, define the *modified torsional rigidity of Ω along u*

$$(87) \quad T_{\text{mod}}(\Omega; u) := 2 \sup_{g \in \mathcal{L}} F(g \circ u), \quad F(w) = \int_\Omega w dx - \frac{1}{2} \int_\Omega |\nabla w|^2 dx.$$

Since the competitors $g \circ u$ form a subclass of $H_0^1(\Omega)$, comparison with (84) gives $T_{\text{mod}}(\Omega; u) \leq T(\Omega)$ always.

Lemma 11.4 (Closed form of T_{mod}). *Under the hypotheses of Definition 11.3, writing $P(t) := \int_{\{u=t\}} |\nabla u| d\mathcal{H}^{n-1}$, the supremum in (87) is uniquely attained by the nondecreasing g_0 with $g_0'(t) = \mu(t)/P(t)$, and*

$$(88) \quad T_{\text{mod}}(\Omega; u) = \int_0^M \frac{\mu(t)^2}{P(t)} dt.$$

Proof. We may restrict to nondecreasing g : replacing g by $\tilde{g}(t) = \int_0^t |g'|$ leaves $\int_\Omega |\nabla(g \circ u)|^2$ unchanged (since $\tilde{g}' = |g'|$ a.e.) and increases $\int_\Omega g \circ u$ (since $\tilde{g} \geq g$), hence increases F . For nondecreasing $g \in \mathcal{L}$, Cavalieri's principle gives $\int_\Omega g(u) dx = \int_0^M g'(t) \mu(t) dt$, while the coarea formula (R2) gives $\int_\Omega |\nabla(g \circ u)|^2 dx = \int_0^M g'(t)^2 P(t) dt$. Thus

$$F(g \circ u) = \int_0^M \left[g'(t) \mu(t) - \frac{1}{2} g'(t)^2 P(t) \right] dt \leq \int_0^M \max_{s \geq 0} \left[s \mu(t) - \frac{1}{2} s^2 P(t) \right] dt = \int_0^M \frac{\mu(t)^2}{2P(t)} dt,$$

the pointwise maximum $\max_s (s\mu - \frac{1}{2}s^2P) = \mu^2/(2P)$ attained at $s = \mu(t)/P(t)$. Equality holds iff $g'(t) = \mu(t)/P(t)$ a.e., which by (86) is an L^∞ function, so $g_0 \in \mathcal{L}$. Multiplying by 2 gives (88). $\square$

The next proposition is the isoperimetric statement that makes the rearrangement decrease the Rayleigh quotient. It is the only place where the Saint–Venant inequality enters.

Proposition 11.5 (Isoperimetric inequality for T_{mod}). *Let $u \in H_0^1(\Omega) \cap L^\infty$, $u \geq 0$, satisfy (86). If $B \subset \mathbb{R}^n$ is a ball with $T(B) = T_{\text{mod}}(\Omega; u)$, then $|B| \leq |\Omega|$, with equality iff Ω is a ball and u is (a translate of) a radial function.*

Proof. $T(B) = T_{\text{mod}}(\Omega; u) \leq T(\Omega)$ by the remark after Definition 11.3. The Saint–Venant inequality (83) gives $T(\Omega) \leq T_n(|\Omega|/\omega_n)^{(n+2)/n}$, while $T(B) = T_n(|B|/\omega_n)^{(n+2)/n}$ by (79). Hence $(|B|/\omega_n)^{(n+2)/n} \leq (|\Omega|/\omega_n)^{(n+2)/n}$, i.e. $|B| \leq |\Omega|$. Equality forces $T_{\text{mod}}(\Omega; u) = T(\Omega)$ and equality in Saint–Venant, hence Ω a ball; the maximiser $g_0 \circ u$ of (87) then coincides with the (radial) torsion function, so u is radial. ([4, Prop. 3.8].) $\square$

The Kohler–Jobin rearrangement. We now build u^* . For $t \in [0, M]$ set $\Omega_t = \{u > t\}$ and let

$$(89) \quad \mathcal{T}(t) := T_{\text{mod}}(\Omega_t; (u - t)_+).$$

Since $|\{(u - t)_+ > s\}| = \mu(t + s)$ and the coarea factor of $(u - t)_+$ at level s is $P(t + s)$, formula (88) gives the explicit expression

$$(90) \quad \mathcal{T}(t) = \int_t^M \frac{\mu(\tau)^2}{P(\tau)} d\tau,$$

so that $\mathcal{T} \in \text{Lip}_{\text{loc}}([0, M])$, $\mathcal{T}(M) = 0$, $\mathcal{T}(0) = T_{\text{mod}}(\Omega; u) =: T_\star$, and for a.e. t

$$(91) \quad \mathcal{T}'(t) = -\frac{\mu(t)^2}{P(t)} < 0.$$

Thus $\mathcal{T}$ is a strictly decreasing bijection $[0, M] \rightarrow [0, T_\star]$.

Definition of u^* . Let $B = B_{R_\star}$ be the ball with $T(B) = T_\star$, i.e. $R_\star = (T_\star/T_n)^{1/(n+2)}$. For $\tau \in [0, T_\star]$ let $R(\tau)$ be the radius with $T(B_{R(\tau)}) = \tau$, i.e.

$$(92) \quad R(\tau) = (\tau/T_n)^{1/(n+2)}, \quad |B_{R(\tau)}| = \omega_n R(\tau)^n = \omega_n (\tau/T_n)^{n/(n+2)}.$$

Let $\varphi := \mathcal{T}^{-1} : [0, T_\star] \rightarrow [0, M]$ (so $\varphi(0) = M$, $\varphi(T_\star) = 0$, $\varphi' < 0$). We define u^* on B to be the radial decreasing function whose value on the sphere $\{|x| = R(\tau)\}$ is $\psi(\tau)$, where $\psi : [0, T_\star] \rightarrow [0, \infty)$ is the locally Lipschitz (monotone, absolutely continuous) function determined by

$$(93) \quad (-\psi'(\tau)) \mu^*(\psi(\tau)) = (-\varphi'(\tau)) \mu(\varphi(\tau)), \quad \psi(T_\star) = 0,$$

μ^* denoting the distribution function of u^* . Concretely the superlevel set $\{u^* > \psi(\tau)\} = B_{R(\tau)}$ has, by construction, torsional rigidity exactly τ , the value of the *modified* torsion that the corresponding superlevel set $\Omega_{\varphi(\tau)}$ of u carries. In particular $\{u^* > 0\} = B = B_{R_\star}$, so

$$(94) \quad T(B) = T_\star = T_{\text{mod}}(\Omega; u) \leq T(\Omega).$$

Since $\mu^*(\psi(\tau)) = |B_{R(\tau)}| = \omega_n R(\tau)^n$, equation (93) integrates to an explicit ψ ; we shall only need the two qualitative consequences (A), (B), proved next.

Lemma 11.6 (Dirichlet energy is preserved). $\int_B |\nabla u^*|^2 dx = \int_\Omega |\nabla u|^2 dx$.

Proof. The original side. By coarea and the change of variable $t = \varphi(\tau)$ (Lemma 2.5 of [4], the one-dimensional area formula for $\varphi = \mathcal{T}^{-1} \in \text{Lip}_{\text{loc}}$ with $\varphi' < 0$),

$$(95) \quad \int_\Omega |\nabla u|^2 dx = \int_0^M P(t) dt = \int_0^{T_\star} P(\varphi(\tau)) (-\varphi'(\tau)) d\tau.$$

Differentiating $\mathcal{T}(\varphi(\tau)) = \tau$ and using (91) gives the key relation

$$(96) \quad -\varphi'(\tau) = \frac{1}{-\mathcal{T}'(\varphi(\tau))} = \frac{P(\varphi(\tau))}{\mu(\varphi(\tau))^2}.$$

Hence, writing $P(\varphi)(-\varphi') = \mu(\varphi)^2 \cdot \frac{P(\varphi)}{\mu(\varphi)^2} \cdot (-\varphi')$ and using (96) for the middle factor, $P(\varphi)(-\varphi') = \mu(\varphi)^2(-\varphi')^2$, so (95) becomes

$$(97) \quad \int_{\Omega} |\nabla u|^2 dx = \int_0^{T_*} \mu(\varphi(\tau))^2 (-\varphi'(\tau))^2 d\tau.$$

The rearranged side. For the radial u^* , the level set $\{u^* = \psi(\tau)\}$ is the sphere of radius $R(\tau)$, on which $|\nabla u^*|$ is constant. A direct computation (differentiate $u^*(x) = \psi(\tau)$ on $|x| = R(\tau)$ and use $R(\tau) = (\tau/T_n)^{1/(n+2)}$) gives, with $P^*(\tau') := \int_{\{u^*=\tau'\}} |\nabla u^*| d\mathcal{H}^{n-1}$ the coarea factor of u^* ,

$$(98) \quad \int_B |\nabla u^*|^2 dx = \int_0^{T_*} \mu^*(\psi(\tau))^2 (-\psi'(\tau))^2 d\tau,$$

the exact radial analogue of (97); this is the specialisation to $p = 2$ of the identity (4.4) of [4], obtained there by the same coarea computation. Now the defining relation (93), $\mu^*(\psi)(-\psi') = \mu(\varphi)(-\varphi')$, makes the integrands of (98) and (97) *equal*:

$$\mu^*(\psi(\tau))^2 (-\psi'(\tau))^2 = [\mu^*(\psi(\tau))(-\psi'(\tau))]^2 = [\mu(\varphi(\tau))(-\varphi'(\tau))]^2 = \mu(\varphi(\tau))^2 (-\varphi'(\tau))^2.$$

Hence $\int_B |\nabla u^*|^2 = \int_{\Omega} |\nabla u|^2$. □

Lemma 11.7 (L^2 norm does not decrease). $\int_B (u^*)^2 dx \geq \int_{\Omega} u^2 dx$, and more generally $\int_B f(u^*) dx \geq \int_{\Omega} f(u) dx$ for every convex $f : [0, \infty) \rightarrow [0, \infty)$ with $f(0) = 0$, strictly if f is strictly convex unless Ω is a ball and u radial.

Proof. The point is the comparison of *distribution functions* along the rearrangement. Apply Proposition 11.5 not to u but to the truncations: for each $\tau \in [0, T_*]$ the ball $B_{R(\tau)}$ has $T(B_{R(\tau)}) = \tau = T_{\text{mod}}(\Omega_{\varphi(\tau)}; (u - \varphi(\tau))_+)$ by construction, so Proposition 11.5 (applied to $\Omega_{\varphi(\tau)}$ with reference function $(u - \varphi(\tau))_+$) yields

$$(99) \quad \mu^*(\psi(\tau)) = |B_{R(\tau)}| \leq |\Omega_{\varphi(\tau)}| = \mu(\varphi(\tau)), \quad \tau \in [0, T_*].$$

Combining (99) with the defining relation (93), $\mu^*(\psi)(-\psi') = \mu(\varphi)(-\varphi')$, gives

$$(100) \quad -\psi'(\tau) = \frac{\mu(\varphi(\tau))}{\mu^*(\psi(\tau))} (-\varphi'(\tau)) \geq -\varphi'(\tau) \quad \text{for a.e. } \tau,$$

since $\mu(\varphi) \geq \mu^*(\psi)$. Integrating from τ to T_* and using $\psi(T_*) = \varphi(T_*) = 0$,

$$(101) \quad \psi(\tau) = \int_{\tau}^{T_*} (-\psi'(s)) ds \geq \int_{\tau}^{T_*} (-\varphi'(s)) ds = \varphi(\tau), \quad \tau \in [0, T_*].$$

Now compute, for convex f with $f(0) = 0$ (so $f(u) = \int_0^u f'(t) dt$ and Cavalieri applies), using the change of variable $t = \varphi(\tau)$:

$$\begin{aligned} \int_{\Omega} f(u) dx &= \int_0^M f'(t) \mu(t) dt = \int_0^{T_{\star}} f'(\varphi(\tau)) (-\varphi'(\tau)) \mu(\varphi(\tau)) d\tau \\ &= \int_0^{T_{\star}} f'(\varphi(\tau)) (-\psi'(\tau)) \mu^*(\psi(\tau)) d\tau \quad \text{by (93)} \\ &\leq \int_0^{T_{\star}} f'(\psi(\tau)) (-\psi'(\tau)) \mu^*(\psi(\tau)) d\tau \quad \text{by (101) and } f' \text{ nondecreasing} \\ &= \int_0^M f'(s) \mu^*(s) ds = \int_B f(u^*) dx, \end{aligned}$$

the last change of variable being $s = \psi(\tau)$. Taking $f(s) = s^2$ proves $\int_B (u^*)^2 \geq \int_{\Omega} u^2$. If f is strictly convex, equality in the middle step forces $\psi = \varphi$ a.e., hence equality in (99), i.e. $|B_{R(\tau)}| = |\Omega_{\varphi(\tau)}|$ for a.e. τ ; by the equality case of the Saint-Venant inequality this makes a.e. superlevel set $\Omega_t = \{u > t\}$ a ball. The sets Ω_t are nested, so for $s > t$ the inclusion $\Omega_s \subset \Omega_t$ forces $|c_s - c_t| \leq r_t - r_s$ on the centres and radii; as $t \downarrow 0$ the radii increase to a limit $r_{\star}$ and the centres are therefore Cauchy, converging to some $c_{\star}$. Since the Ω_t increase to $\{u > 0\} = \Omega$ (up to a null set), $\Omega = B(c_{\star}, r_{\star})$; hence Ω is a ball and u is radial, consistently with the equality case of Proposition 11.5. $\square$

Proof of Theorem 11.1. Apply the construction to the first eigenfunction u of Ω (a valid reference function by (85) and (R1)). By Lemmas 11.6 and 11.7 the rearranged $u^* \in H_0^1(B)$ satisfies $\int_B |\nabla u^*|^2 = \int_{\Omega} |\nabla u|^2$ and $\int_B (u^*)^2 \geq \int_{\Omega} u^2$. By the Rayleigh principle on B (78) and these two facts,

$$\lambda_1(B) \leq \frac{\int_B |\nabla u^*|^2}{\int_B (u^*)^2} \leq \frac{\int_{\Omega} |\nabla u|^2}{\int_{\Omega} u^2} = \lambda_1(\Omega).$$

Multiplying $\lambda_1(B) \leq \lambda_1(\Omega)$ by $T(B)^{\beta} > 0$ and using $T(B) \leq T(\Omega)$ from (94) and the scale invariance $\Psi(B) = \Psi(B_1)$:

$$T(B_1) L_n^{\beta} = \Psi(B) = T(B) \lambda_1(B)^{\beta} \leq T(\Omega) \lambda_1(\Omega)^{\beta} = \Psi(\Omega).$$

This is (81) (for an arbitrary ball, $\Psi(B) = \Psi(B_1)$ on both sides). Equality forces equality throughout, in particular in Lemma 11.7 with the strictly convex $f(s) = s^2$, hence Ω is a ball. $\square$

Remark 11.8 (On the sign that distinguishes Kohler–Jobin from Faber–Krahn). The torsion-preserving rearrangement makes the ball $B_{R(\tau)}$ *smaller* than the superlevel set $\Omega_{\varphi(\tau)}$, (99) — the opposite of Schwarz symmetrisation. It is precisely this that lets the Dirichlet energy be *exactly preserved* (Lemma 11.6) rather than merely decreased, while the level-set volumes *grow under the inverse rearrangement*, forcing $\int (u^*)^2 \geq \int u^2$ (Lemma 11.7). The naive attempt to compare L^2 masses level by level fails because the rearranged superlevel balls are smaller; the correct comparison is through the *distribution-function inequality* (101), driven by the isoperimetric Proposition 11.5.

11.3. Part 2: transfer to sharp Faber–Krahn stability. We now use Theorem 11.1 as a tool. Throughout, $L := \lambda_1(B^*)$, $T_0 := T(B^*)$, $\beta = (n+2)/2$, $p := 1/\beta = 2/(n+2) \in (0, 1]$.

Lemma 11.9 (Pointwise Kohler–Jobin transfer). *For every open $\Omega \subset \mathbb{R}^n$ with $|\Omega| = |B^*|$,*

$$(102) \quad \lambda_1(\Omega) \geq L \left(\frac{T_0}{T(\Omega)} \right)^{1/\beta} = L \left(\frac{T_0}{T(\Omega)} \right)^{2/(n+2)}.$$

Proof. Theorem 11.1 with $B = B^*$ gives $T(\Omega)\lambda_1(\Omega)^\beta \geq T_0 L^\beta$; solve for $\lambda_1(\Omega)$. $\square$

By Saint–Venant (83) at fixed volume, $T(\Omega) \leq T_0$, so the torsion deficit $\sigma := \sigma(\Omega) := T_0 - T(\Omega) \geq 0$. Put $s := \sigma/T_0 \in [0, 1]$; then (102) reads

$$(103) \quad \lambda_1(\Omega) - L \geq L[(1-s)^{-p} - 1], \quad p = \frac{2}{n+2}.$$

Lemma 11.10 (An elementary one-variable inequality). *For $p \in (0, 1]$ and $s \in [0, 1]$, $(1-s)^{-p} - 1 \geq ps$.*

Proof. $f(s) = (1-s)^{-p} - 1$ has $f(0) = 0$ and $f'(s) = p(1-s)^{-(p+1)} \geq p$ on $[0, 1]$ because $(1-s)^{-(p+1)} \geq 1$. By the fundamental theorem of calculus $f(s) = \int_0^s f' \geq ps$. $\square$

Lemma 11.11 (Small-deficit linearisation). *If $|\Omega| = |B^*|$ and $\sigma \leq T_0/2$, then $\lambda_1(\Omega) - L \geq \frac{2L}{(n+2)T_0} \sigma$.*

Proof. With $s = \sigma/T_0 \leq 1/2$, $p = 2/(n+2)$, combine (103) and Lemma 11.10: $\lambda_1(\Omega) - L \geq Lps = Lp\sigma/T_0 = \frac{2L}{(n+2)T_0} \sigma$. $\square$

Lemma 11.12 (Large-deficit branch). *If $|\Omega| = |B^*|$ and $\sigma > T_0/2$, then $\lambda_1(\Omega) - L \geq L(2^{2/(n+2)} - 1)$, hence, since $\mathcal{A}(\Omega) < 2$,*

$$(104) \quad \lambda_1(\Omega) - L \geq \frac{L(2^{2/(n+2)} - 1)}{4} \mathcal{A}(\Omega)^2.$$

Proof. $\sigma > T_0/2$ gives $T(\Omega) < T_0/2$, so $T_0/T(\Omega) > 2$ and (102) yields $\lambda_1(\Omega) \geq L 2^{2/(n+2)}$. Since $\mathcal{A}(\Omega)^2 \leq 4$ at fixed volume, dividing by 4 gives (104). $\square$

Theorem 11.13 (Sharp Faber–Krahn via Kohler–Jobin). *By the global sharp Saint–Venant stability estimate of Theorem 10.2, with $c_{\text{SV}}^{\text{glob}}(n) > 0$,*

$$(105) \quad T(B^*) - T(\Omega) = 2(E(\Omega) - E(B^*)) \geq 2c_{\text{SV}}^{\text{glob}}(n) \left(\frac{|\Omega|}{\omega_n} \right)^{(n+2)/n} \mathcal{A}(\Omega)^2$$

for all open bounded $\Omega \subset \mathbb{R}^n$, $0 < |\Omega| < \infty$. Then

$$(106) \quad \delta_{\text{FK}}(\Omega) = \left(\frac{|\Omega|}{\omega_n} \right)^{2/n} (\lambda_1(\Omega) - \lambda_1(B^*)) \geq c_{\text{FK}}(n) \mathcal{A}(\Omega)^2,$$

with the explicit constant

$$(107) \quad c_{\text{FK}}(n) = \min \left\{ \frac{4L_n c_{\text{SV}}^{\text{glob}}(n)}{(n+2)T_n}, \frac{L_n(2^{2/(n+2)} - 1)}{4} \right\},$$

equivalently $\mathcal{A}(\Omega)^2 \leq C_{\text{FK}}(n) \delta_{\text{FK}}(\Omega)$ with $C_{\text{FK}}(n) = 1/c_{\text{FK}}(n)$.

Proof. Both δ_{FK} and $\mathcal{A}$ are scale invariant, and so is (105); rescale so that $|\Omega| = \omega_n$, whence $B^* = B_1$, $L = L_n$, $T_0 = T_n$, and the volume prefactor $(|\Omega|/\omega_n)^{2/n} = 1$. Split on the torsion deficit $\sigma = T_n - T(\Omega)$.

Case 1 ($\sigma \leq T_n/2$). By Lemma 11.11 and (105) (which now reads $\sigma \geq 2c_{\text{SV}}^{\text{glob}}(n)\mathcal{A}(\Omega)^2$),

$$\lambda_1(\Omega) - L_n \geq \frac{2L_n}{(n+2)T_n} \sigma \geq \frac{2L_n}{(n+2)T_n} \cdot 2c_{\text{SV}}^{\text{glob}}(n) \mathcal{A}(\Omega)^2 = \frac{4L_n c_{\text{SV}}^{\text{glob}}(n)}{(n+2)T_n} \mathcal{A}(\Omega)^2.$$

Case 2 ($\sigma > T_n/2$). By Lemma 11.12, $\lambda_1(\Omega) - L_n \geq \frac{L_n(2^{2/(n+2)}-1)}{4} \mathcal{A}(\Omega)^2$.

In both cases $\delta_{\text{FK}}(\Omega) = \lambda_1(\Omega) - L_n \geq c_{\text{FK}}(n) \mathcal{A}(\Omega)^2$ with $c_{\text{FK}}(n)$ the minimum (107). Undoing the scaling preserves the inequality since every term is scale invariant. $\square$

Remark 11.14 (Sharpness of the exponent). The exponent 2 on $\mathcal{A}$ in (106) is optimal: smooth ellipsoidal perturbations of the ball have $\lambda_1(\Omega) - L_n \asymp \mathcal{A}(\Omega)^2$ ([5, 4] and references therein). The transfer is non-perturbative: no compactness, contradiction, or Taylor expansion is used past the elementary Lemma 11.10.

Proof of Theorem 1.1. For an open set of finite measure with $\lambda_1(\Omega) = +\infty$ the inequality is trivial, since its left side is then $+\infty$ while $\mathcal{A}(\Omega)^2 \leq 4$. Otherwise $\lambda_1(\Omega) < \infty$, and Theorem 11.13 applies: its Saint-Venant hypothesis (105) is exactly the global estimate (27) of Theorem 10.2, so (106) holds with $c_{\text{FK}}(n)$ as in (107). This is the assertion of Theorem 1.1. $\square$

12. PROOF ARCHITECTURE AND CONSTANTS

The argument is organised around the following principal steps; the first five produce the bounded Saint-Venant estimate, the last two upgrade it to Faber-Krahn.

- (1) The exact level-set deficit identity (Theorem 3.11) writes $\delta_T(\Omega)$ as a weighted reduced-boundary integral of the Cauchy-Schwarz defect D_H and the isoperimetric defect D_I ; the profile-gap identity and the bad-density bound of Proposition 4.1 are its ρ -parametrised consequences.
- (2) The affine shell estimate (Lemma 7.1) is a finite-perimeter limsup estimate, proved by localized coarea differentiation on variable slabs (Lemma 6.1) together with the local BV deformation-tail lemma (Lemma 6.2). Its only bounded-variation input is Reshetnyak continuity for a continuous weight, applied as an *upper* bound to a swept-volume tail made arbitrarily small by inner regularity; no lower-semicontinuity passage is used.
- (3) The endpoint trace (Lemma 9.1) uses the upper Dini derivative D^+q together with the global Lipschitz bound for q (Lemma 7.3).
- (4) Bad-density levels are charged by Lebesgue measure rather than by the deficit, which is possible because q is uniformly Lipschitz by nestedness.
- (5) The same-centre good-level package (Theorem 5.4) is obtained from the Fusco-Julin strong quantitative isoperimetric inequality (Theorem 5.1); the passage from normal oscillation to the homothetic trace is the radial trace Lemma 5.3.
- (6) The bounded reduction (Theorem 10.2) removes the confinement $\Omega \subset B_R$: a constructive De Giorgi L^∞ tail estimate (Lemma 10.5) feeds a truncation construction (Lemma 10.9) that replaces Ω by a bounded surrogate of dimensional diameter with comparable deficit and controlled asymmetry loss; gluing the bounded estimate to the trivial large-deficit branch gives the global Saint-Venant inequality with a dimensional constant.
- (7) The Kohler-Jobin rearrangement (Theorem 11.1), built to preserve the modified torsional rigidity of superlevel sets, gives $T(\Omega)\lambda_1(\Omega)^{(n+2)/2} \geq T_n L_n^{(n+2)/2}$; combined with the global Saint-Venant inequality and an elementary one-variable estimate it yields Theorem 1.1 (Theorem 11.13).

All constants in Theorem 9.2 depend only on n , R , and ρ_* , together with the Fusco-Julin constant $C_{\text{FJ}}(n)$. Tracing them through the proof: $C_L = 2n^2\omega_n$ and $C_\tau = 4n/(\omega_n\rho_*^{n+1})$ (Proposition 4.1); $\kappa = (1-\rho_*)/4$ and $\tau_0 = \rho_*/(2n)$; the same-centre constant $C_{\text{sc}}(n, R, \rho_*)$ and threshold $\eta_I(n, R, \rho_*)$ (Theorem 5.4, Lemma 8.1); the ball-ball lower constant $\frac{1}{2}\omega_{n-1}(3/4)^{(n-1)/2}$; the slab

length $L_0 = (1 - \rho_*)/8$; and the large-deficit cutoff $4/\delta_0$. No constant depends on Ω beyond these parameters. The bounded reduction internalises the confinement at the dimensional radius $R_*(n) = \max\{d(n), 2\}$, producing the purely dimensional $c_{SV}^{\text{glob}}(n)$ (Theorem 10.2), and the Kohler–Jobin transfer outputs $c_{FK}(n) = \min\{4L_n c_{SV}^{\text{glob}}(n)/((n+2)T_n), L_n(2^{2/(n+2)} - 1)/4\}$ of Theorem 1.1, with $L_n = \lambda_1(B_1)$, $T_n = \omega_n/(n(n+2))$.

APPENDIX A. STANDARD TOOLS FROM THE THEORY OF SETS OF FINITE PERIMETER

For the reader’s convenience we record the three classical results used above, in the exact form needed; all hold for sets of finite perimeter with no graph or manifold regularity.

- (1) *BV coarea formula* [14, Thm. 13.1], [1, Thm. 3.40]: for $u \in BV_{\text{loc}}(\mathbb{R}^n)$ and Borel $\varphi \geq 0$, $\int |\nabla u| \varphi = \int_{\mathbb{R}} (\int_{\partial^* \{u > t\}} \varphi d\mathcal{H}^{n-1}) dt$. This underlies (2) and Lemma 6.1.
- (2) *Gauss–Green on sets of finite perimeter* [14, Ch. 17]: for a bounded Lipschitz vector field X and a set E of finite perimeter, $\int_E \text{div } X = \int_{\partial^* E} X \cdot \nu_E d\mathcal{H}^{n-1}$. This is the form used in Lemma 5.3.
- (3) *Strict BV convergence and Reshetnyak continuity* [1, Thm. 2.39], [14, Ch. 20]: if $v_k \rightarrow v$ in L_{loc}^1 with $|Dv_k|(\mathbb{R}^n) \rightarrow |Dv|(\mathbb{R}^n)$, then $\int \psi d|Dv_k| \rightarrow \int \psi d|Dv|$ for every continuous compactly supported ψ . Applied to mollifications $v_k = \mathbf{1}_E * \varphi_{1/k}$, this gives the upper bound in Lemma 6.2.

APPENDIX B. DEPENDENCY CERTIFICATE

Every step is either proved in this manuscript or quoted from a named published theorem.

Proved internally: the exact deficit identity and its inputs (coarea form of $-m'$, the Lipschitz-truncation flux identity, the rearranged primitive with $I'' = -1/\alpha$) in §3; the profile-gap and bad-density estimates in §4; the oriented radial trace, the same-centre package, the velocity defect, and the superlevel asymmetry transfer in §5; the two-sided coarea differentiation lemma, the shell-admissible registration, and the local BV deformation tail in §6; the affine shell estimate, the one-sided first variation, and the Lipschitz bound for q in §7; the centre transfer and the positive kinetic estimate in §8; the endpoint trace and the bounded stability theorem in §9; the De Giorgi L^∞ tail estimate, the truncation construction, and the global bounded reduction in §10; and the Kohler–Jobin inequality (via the modified-torsion rearrangement) together with the transfer to Faber–Krahn in §11.

Quoted with exact citation: the Fusco–Julin strong quantitative isoperimetric inequality [12]; the BV coarea formula, the finite-perimeter Gauss–Green theorem, and Reshetnyak continuity [1, 14]; the De Giorgi isoperimetric inequality [11]; the Talenti rearrangement comparison and L^∞ bound [17]; the classical Saint–Venant inequality; the Sobolev inequality; the Brothers–Ziemer regularity of the distribution function [1]; the absolutely continuous monotone inverse theorem [10]; eigenfunction regularity, interior elliptic regularity, and the weak maximum principle [13]; and the Pólya–Szegő identification of the ball energy [15, 2].

The suboptimal seed is proved internally. The truncation of §10 uses a *suboptimal* Saint–Venant stability bound $\delta_T(\Omega) \geq c\mathcal{A}(\Omega)^4$ (Lemma 10.7) only to bound the number of truncation steps. This is proved here (Lemma 10.7) from the exact deficit identity, the profile-gap identity, and the superlevel transfer established above, together with the Fusco–Julin inequality already in use; it relies on no stability estimate external to this paper. Consequently every ingredient of the argument is either proved above or is one of the standard tools just listed.

REFERENCES

- [1] L. Ambrosio, N. Fusco, and D. Pallara, *Functions of Bounded Variation and Free Discontinuity Problems*, Oxford University Press, 2000. (Thm. 3.40, coarea; Thm. 3.99, chain rule; §3.5, reduced boundaries.)
- [2] C. Bandle, *Isoperimetric Inequalities and Applications*, Pitman, 1980. Ch. II.
- [3] L. Brasco, G. De Philippis, and B. Velichkov, *Faber–Krahn inequalities in sharp quantitative form*, Duke Math. J. **164** (2015), 1777–1831.
- [4] L. Brasco, *On torsional rigidity and principal frequencies: an invitation to the Kohler–Jobin rearrangement technique*, ESAIM Control Optim. Calc. Var. **20** (2014), 315–338.
- [5] L. Brasco and G. De Philippis, *Spectral inequalities in quantitative form*, in *Shape Optimization and Spectral Theory*, De Gruyter, 2017, pp. 201–281.
- [6] N. Fusco, F. Maggi, and A. Pratelli, *Stability estimates for certain Faber–Krahn, isocapacitary and Cheeger inequalities*, Ann. Sc. Norm. Super. Pisa Cl. Sci. (5) **8** (2009), 51–71.
- [7] M.-T. Kohler-Jobin, *Une méthode de comparaison isopérimétrique de fonctionnelles de domaines de la physique mathématique*, Z. Angew. Math. Phys. **29** (1978), 757–766.
- [8] M.-T. Kohler-Jobin, *Démonstration de l’inégalité isopérimétrique $P\lambda^2 \geq \pi j_0^4/2$ conjecturée par Pólya et Szegő*, C. R. Acad. Sci. Paris **281** (1975), 119–121.
- [9] M.-T. Kohler-Jobin, *Symmetrization with equal Dirichlet integrals*, SIAM J. Math. Anal. **13** (1982), 153–161.
- [10] P. Bénilan, H. Brezis, and M. G. Crandall, *A semilinear equation in $L^1(\mathbb{R}^N)$* , J. Anal. Math. **27** (1975), 273–299. (Thm. 0.4 and §1: absolutely continuous monotone inverse.)
- [11] E. De Giorgi, *Sulla proprietà isoperimetrica dell’ipersfera, nella classe degli insiemi aventi frontiera orientata di misura finita*, Atti Accad. Naz. Lincei **5** (1958), 33–44.
- [12] N. Fusco and V. Julin, *A strong form of the quantitative isoperimetric inequality*, Calc. Var. Partial Differential Equations **50** (2014), 925–937.
- [13] D. Gilbarg and N. S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Classics in Mathematics, Springer, 2001. (Interior $W^{2,p}$ regularity; weak maximum principle.)
- [14] F. Maggi, *Sets of Finite Perimeter and Geometric Variational Problems*, Cambridge University Press, 2012. (Ch. 13, coarea; Ch. 17, Gauss–Green on finite-perimeter sets.)
- [15] G. Pólya and G. Szegő, *Isoperimetric Inequalities in Mathematical Physics*, Princeton University Press, 1951.
- [16] G. Stampacchia, *Le problème de Dirichlet pour les équations elliptiques du second ordre à coefficients discontinus*, Ann. Inst. Fourier **15** (1965), 189–258. ($D^2u = 0$ a.e. on $\{\nabla u = 0\}$ for Sobolev maps.)
- [17] G. Talenti, *Elliptic equations and rearrangements*, Ann. Mat. Pura Appl. **110** (1976), 353–372.