

SMT: Linear Arithmetic and Combination of Theories

ALC

IST, ULisboa

Outline

Theories: Real / Integer Linear Arithmetic

Theories Solver for RLA: The Simplex Method

Theories Solver for RLA: Fourier-Motzkin

Theories: Integer Linear Arithmetic

SMT: Recap

Theory Combinations

Outline

Theories: Real / Integer Linear Arithmetic

Theories Solver for RLA: The Simplex Method

Theories Solver for RLA: Fourier-Motzkin

Theories: Integer Linear Arithmetic

SMT: Recap

Theory Combinations

Context – SMT Lazy Solving

input : formula φ in theory \mathcal{T}

output: truth value

```
1  $\alpha \leftarrow \mathcal{T}2\mathcal{B}(\varphi)$  // abstract input formula
2 while true do
3    $(\text{res}, \tau) \leftarrow \text{SAT}(\alpha)$  // Boolean model
4   if  $\text{res} = \text{false}$  then return false
5    $\mathcal{L} \leftarrow \bigcup_{l \in \tau} \mathcal{B}2\mathcal{T}(l)$  // convert to theory model
6    $(\text{res}, \mathcal{L}') \leftarrow \mathcal{T}\text{-SAT}(\mathcal{L})$  // theory check
7   if  $\text{res} = \text{true}$  then return true
8    $\alpha \leftarrow \alpha \wedge \bigvee_{l \in \mathcal{L}'} \neg \mathcal{T}2\mathcal{B}(l)$  // block explanation
9 end
```

Linear arithmetic

atom : sum op sum

op : \leq | $<$ | $=$

sum : term + sum

term : uninterpretedConstant | interpretedConstant

- Integers: **ILA**
- Reals: **RLA**

A simple example

- Example RLA/ILA formula:

$$\begin{aligned} &((x_1 + x_2 \geq 2 \vee x_3 + x_4 \geq 3) \wedge \\ &((x_1 + x_2 \leq 1 \vee x_3 + x_4 \leq 2) \wedge \\ &((x_5 + x_6 \geq 2 \vee x_7 + x_8 \geq 3) \wedge \\ &((x_5 + x_6 \leq 1 \vee x_7 + x_8 \leq 2) \end{aligned}$$

- Represent Boolean structure as CNF formula:

$$(a \vee b) \wedge (c \vee d) \wedge (e \vee f) \wedge (g \vee h)$$

- Interaction between SAT solver & theory solver (RLA/ILA):

SAT Outcome	Model/Core	(R I)LA Outcome	Explanation clause (sent to SAT solver)
----------------	------------	----------------------	--

A simple example

- Example RLA/ILA formula:

$$\begin{aligned} &((x_1 + x_2 \geq 2 \vee x_3 + x_4 \geq 3) \wedge \\ &((x_1 + x_2 \leq 1 \vee x_3 + x_4 \leq 2) \wedge \\ &((x_5 + x_6 \geq 2 \vee x_7 + x_8 \geq 3) \wedge \\ &((x_5 + x_6 \leq 1 \vee x_7 + x_8 \leq 2) \end{aligned}$$

- Represent Boolean structure as CNF formula:

$$(a \vee b) \wedge (c \vee d) \wedge (e \vee f) \wedge (g \vee h)$$

- Interaction between SAT solver & theory solver (RLA/ILA):

SAT Outcome	Model/Core	(R I)LA Outcome	Explanation clause (sent to SAT solver)
SAT	$\{a, \bar{b}, c, \bar{d}, e, \bar{f}, g, \bar{h}\}$	UNSAT	$(\bar{a} \vee \bar{c})$

A simple example

- Example RLA/ILA formula:

$$\begin{aligned} &((x_1 + x_2 \geq 2 \vee x_3 + x_4 \geq 3) \wedge \\ &((x_1 + x_2 \leq 1 \vee x_3 + x_4 \leq 2) \wedge \\ &((x_5 + x_6 \geq 2 \vee x_7 + x_8 \geq 3) \wedge \\ &((x_5 + x_6 \leq 1 \vee x_7 + x_8 \leq 2) \end{aligned}$$

- Represent Boolean structure as CNF formula:

$$(a \vee b) \wedge (c \vee d) \wedge (e \vee f) \wedge (g \vee h)$$

- Interaction between SAT solver & theory solver (RLA/ILA):

SAT Outcome	Model/Core	(R I)LA Outcome	Explanation clause (sent to SAT solver)
SAT	$\{a, \bar{b}, c, \bar{d}, e, \bar{f}, g, \bar{h}\}$	UNSAT	$(\bar{a} \vee \bar{c})$
SAT	$\{a, \bar{b}, \bar{c}, d, e, \bar{f}, g, \bar{h}\}$	UNSAT	$(\bar{e} \vee \bar{g})$

A simple example

- Example RLA/ILA formula:

$$\begin{aligned} &((x_1 + x_2 \geq 2 \vee x_3 + x_4 \geq 3) \wedge \\ &((x_1 + x_2 \leq 1 \vee x_3 + x_4 \leq 2) \wedge \\ &((x_5 + x_6 \geq 2 \vee x_7 + x_8 \geq 3) \wedge \\ &((x_5 + x_6 \leq 1 \vee x_7 + x_8 \leq 2) \end{aligned}$$

- Represent Boolean structure as CNF formula:

$$(a \vee b) \wedge (c \vee d) \wedge (e \vee f) \wedge (g \vee h)$$

- Interaction between SAT solver & theory solver (RLA/ILA):

SAT Outcome	Model/Core	(R I)LA Outcome	Explanation clause (sent to SAT solver)
SAT	$\{a, \bar{b}, c, \bar{d}, e, \bar{f}, g, \bar{h}\}$	UNSAT	$(\bar{a} \vee \bar{c})$
SAT	$\{a, \bar{b}, \bar{c}, d, e, \bar{f}, g, \bar{h}\}$	UNSAT	$(\bar{e} \vee \bar{g})$
SAT	$\{a, \bar{b}, \bar{c}, d, e, \bar{f}, \bar{g}, h\}$	SAT	Done

Outline

Theories: Real / Integer Linear Arithmetic

Theories Solver for RLA: The Simplex Method

Theories Solver for RLA: Fourier-Motzkin

Theories: Integer Linear Arithmetic

SMT: Recap

Theory Combinations

Using Simplex

Recap. Ch. 29 of Introduction to Algorithms (Cormen et al. – 3rd Ed.)

$$\begin{array}{rclcl} x_1 & + & x_2 & \leq & 0 \\ x_1 & & & + & x_3 \leq 0 \\ & & & - & x_3 \leq -1 \\ & & x_1, x_2, x_3 & \geq & 0 \end{array}$$

Using Simplex

Recap. Ch. 29 of Introduction to Algorithms (Cormen et al. – 3rd Ed.)

$$\begin{array}{rclcl} x_1 & + & x_2 & & \leq & 0 \\ x_1 & & & + & x_3 & \leq & 0 \\ & & & - & x_3 & \leq & -1 \\ & & x_1, x_2, x_3 & & \geq & 0 \end{array}$$

Basic solution does not work. Build auxiliary linear program:

maximize $-x_0$
such that

$$\begin{array}{rclcl} x_1 & + & x_2 & & - & x_0 & \leq & 0 \\ x_1 & & & + & x_3 & - & x_0 & \leq & 0 \\ & & & - & x_3 & - & x_0 & \leq & -1 \\ & & x_1, x_2, x_3, x_0 & & & & \geq & 0 \end{array}$$

Using Simplex

Slack form:

$$\begin{array}{rclclclcl} z & = & & & & - & x_0 \\ s_1 & = & 0 & - & x_1 & - & x_2 & + & x_0 \\ s_2 & = & 0 & - & x_1 & & - & x_3 & + & x_0 \\ s_3 & = & -1 & & & & + & x_3 & + & x_0 \end{array}$$

Pivot between x_0 and s_3

$$\begin{array}{rclclclcl} z & = & -1 & & & + & x_3 & - & s_3 \\ s_1 & = & 1 & - & x_1 & - & x_2 & - & x_3 & + & s_3 \\ s_2 & = & 1 & - & x_1 & & - & 2x_3 & + & s_3 \\ x_0 & = & 1 & & & & - & x_3 & + & s_3 \end{array}$$

Using Simplex

Slack form:

$$\begin{array}{rclclclcl} z & = & -1 & & & + & x_3 & - & s_3 \\ s_1 & = & 1 & - & x_1 & - & x_2 & - & x_3 & + & s_3 \\ s_2 & = & 1 & - & x_1 & & & - & 2x_3 & + & s_3 \\ x_0 & = & 1 & & & & & - & x_3 & + & s_3 \end{array}$$

Pivot between x_3 and s_2

$$\begin{array}{rclclclcl} z & = & -\frac{1}{2} & - & \frac{x_1}{2} & & - & \frac{s_2}{2} & - & \frac{s_3}{2} \\ s_1 & = & \frac{1}{2} & - & \frac{x_1}{2} & - & x_2 & + & \frac{s_2}{2} & + & \frac{s_3}{2} \\ x_3 & = & \frac{1}{2} & - & \frac{x_1}{2} & & & - & \frac{s_2}{2} & + & \frac{s_3}{2} \\ x_0 & = & \frac{1}{2} & + & \frac{x_1}{2} & & & + & \frac{s_2}{2} & + & \frac{s_3}{2} \end{array}$$

Using Simplex

- Optimal solution was found with $x_0 \neq 0$. RLA formula is **unsatisfiable**
 - If $x_0 = 0$, then formula would be satisfiable
- Unsatisfiable core can be determined by analysing final slack form
 - Variable s_i is a non-basic variable (right-hand side) in final slack form means constraint i belongs to the unsatisfiable core
- In our example, the unsatisfiable core would be:

$$\begin{array}{rclcl} x_1 & & + & x_3 & \leq & 0 \\ & & - & x_3 & \leq & -1 \end{array}$$

Remarks on Simplex

Standard Simplex only supports weak inequalities (\leq).

But we also have $=, <$.

- Rewrite equalities with two inequalities
- Rewrite strict inequalities as weak with additional variable:
 1. To decide: $\bigwedge_i \alpha_{i1}x_1 + \cdots + \alpha_{in}x_n < \beta_i$
 2. Maximize x_{n+1} subject to $\bigwedge_i \alpha_{i1}x_1 + \cdots + \alpha_{in}x_n + x_{n+1} \leq \beta_i$
 3. If optimum **positive**, original satisfiable.

Outline

Theories: Real / Integer Linear Arithmetic

Theories Solver for RLA: The Simplex Method

Theories Solver for RLA: Fourier-Motzkin

Theories: Integer Linear Arithmetic

SMT: Recap

Theory Combinations

Using Fourier-Motzkin (FM)

- Alternative to Simplex
- Idea:
 - If $A \leq x$ and $x \leq B$ then $A \leq B$
 - Elimination by enforcing **all** bounds to be nonempty:

$$\left(\bigwedge_i A_i \leq x \wedge \bigwedge_j x \leq B_j \right) \Leftrightarrow \bigwedge_{i,j} A_i \leq B_j$$

- **Complexity:** Elimination of one variable is quadratic. Elimination of n variables: $m^2/4, m^4/16, \dots, m^{2^n}/4^n$

FM phase 1 eliminate equalities

$$\sum_{j=1}^n a_{ij} \cdot x_j = b_i$$

- Pick equality i and remove x_n

FM phase 1 eliminate equalities

$$\sum_{j=1}^n a_{ij} \cdot x_j = b_i$$

$$x_n = \frac{b_i}{a_{in}} - \sum_{j=1}^{n-1} \frac{a_{ij}}{a_{in}} \cdot x_j$$

- Pick equality i and remove x_n

FM phase 1 eliminate equalities

$$\sum_{j=1}^n a_{ij} \cdot x_j = b_i$$

$$x_n = \frac{b_i}{a_{in}} - \sum_{j=1}^{n-1} \frac{a_{ij}}{a_{in}} \cdot x_j$$

$$\bigwedge_{i=1}^m \sum_{j=1}^n a_{ij} \cdot x_j \leq b_i$$

- Pick equality i and remove x_n

FM phase 2: Variable Elimination

$$\sum_{j=1}^n a_{ij} \cdot x_j \leq b_i$$

- Pick inequality i and remove x_n

FM phase 2: Variable Elimination

$$\sum_{j=1}^n a_{ij} \cdot x_j \leq b_i$$

$$a_{in} \cdot x_n \leq b_i - \sum_{j=1}^{n-1} a_{ij} \cdot x_j$$

- Pick inequality i and remove x_n

FM phase 2: Variable Elimination

$$\sum_{j=1}^n a_{ij} \cdot x_j \leq b_i$$

$$a_{in} \cdot x_n \leq b_i - \sum_{j=1}^{n-1} a_{ij} \cdot x_j$$

$$x_n \leq \frac{b_i}{a_{in}} - \sum_{j=1}^{n-1} \frac{a_{ij}}{a_{in}} \cdot x_j$$

- Pick inequality i and remove x_n

FM phase 2: Variable Elimination

$$\sum_{j=1}^n a_{ij} \cdot x_j \leq b_i$$

$$a_{in} \cdot x_n \leq b_i - \sum_{j=1}^{n-1} a_{ij} \cdot x_j$$

$$x_n \leq \frac{b_i}{a_{in}} - \sum_{j=1}^{n-1} \frac{a_{ij}}{a_{in}} \cdot x_j$$

$$\beta_l \leq x_n \leq \beta_u$$

- Pick inequality i and remove x_n
- Combine **all** pairs l and u

FM phase 2: Variable Elimination

$$\sum_{j=1}^n a_{ij} \cdot x_j \leq b_i$$

$$a_{in} \cdot x_n \leq b_i - \sum_{j=1}^{n-1} a_{ij} \cdot x_j$$

$$x_n \leq \frac{b_i}{a_{in}} - \sum_{j=1}^{n-1} \frac{a_{ij}}{a_{in}} \cdot x_j$$

$$\beta_l \leq x_n \leq \beta_u$$

$$\beta_l \leq \beta_u$$

- Pick inequality i and remove x_n
- Combine **all** pairs l and u
- If variable unbounded, then **remove** variable

FM: An example

- Initial formula:

$$\begin{array}{rclclcl} x_1 & - & x_2 & & \leq & 0 \\ x_1 & & & - & x_3 & \leq 0 \\ -x_1 & + & x_2 & + & 2x_3 & \leq 0 \\ & & & & -x_3 & \leq -1 \end{array}$$

- Eliminate x_1 :

FM: An example

- Initial formula:

$$\begin{array}{rclcl} x_1 & - & x_2 & & \leq & 0 \\ x_1 & & & - & x_3 & \leq & 0 \\ -x_1 & + & x_2 & + & 2x_3 & \leq & 0 \\ & & & & -x_3 & \leq & -1 \end{array}$$

- Eliminate x_1 :

$$\text{Bounds: } (x_2 + 2x_3) \leq x_1 \leq \min(x_2, x_3)$$

$$\begin{array}{rclcl} & & 2x_3 & \leq & 0 \\ x_2 & + & x_3 & \leq & 0 \\ & & -x_3 & \leq & -1 \end{array}$$

FM contd.

- Eliminate x_2 (unbounded removed):

$$\begin{array}{rcl} 2x_3 & \leq & 0 \\ -x_3 & \leq & -1 \end{array}$$

- Eliminate x_3 :

$$\text{Bounds: } 1 \leq x_3 \leq 0$$

$$1 \leq 0$$

FM contd.

- Eliminate x_2 (unbounded removed):

$$\begin{array}{rcl} 2x_3 & \leq & 0 \\ -x_3 & \leq & -1 \end{array}$$

- Eliminate x_3 :

$$\text{Bounds: } 1 \leq x_3 \leq 0$$

$$1 \leq 0$$

- Formula is **unsatisfiable**
- Unsatisfiable core can be determined by a trace back on the dependencies of the final constraints

Outline

Theories: Real / Integer Linear Arithmetic

Theories Solver for RLA: The Simplex Method

Theories Solver for RLA: Fourier-Motzkin

Theories: Integer Linear Arithmetic

SMT: Recap

Theory Combinations

Decision procedures for integer domains

- Number of techniques, integer linear programming (ILP) solvers
- Several techniques based on
 - solving in reals (relaxation)
 - blocking non-integer solutions

Branch and Bound in Integer Linear Programming

- Main steps:

Branch and Bound in Integer Linear Programming

- Main steps:
 1. Solve relaxed linear program (e.g. Simplex)

Branch and Bound in Integer Linear Programming

- Main steps:
 1. Solve relaxed linear program (e.g. Simplex)
 2. Pick real valued variable $x_i = v_i$ and create **two** new constraints:
 - ▶ $x_i \leq \lfloor v_i \rfloor$
 - ▶ $x_i \geq \lceil v_i \rceil$
 3. Split on the two constraints (one and only one must hold)

Outline

Theories: Real / Integer Linear Arithmetic

Theories Solver for RLA: The Simplex Method

Theories Solver for RLA: Fourier-Motzkin

Theories: Integer Linear Arithmetic

SMT: Recap

Theory Combinations

SMT: Approaches

- Two main approaches: **eager** and **lazy**
- Eager techniques convert to a single SAT call
- Lazy techniques call SAT repeatedly

- **Small model property:** in some theories, if there is a model there is a small one (bounded)
- **Example:** Satisfiable formulas in *Integer difference logic*'s must have a bounded solution
- **Ackermann reduction:** encode congruence axioms as implications

- SAT handles the propositional structure
- **Theory solver** checks models from the SAT solver within the theory
- **Example:** *equivalence with uninterpreted functions* solved by congruence closure
- **Nelson-Oppen** closure enables combining multiple theory solvers

Outline

Theories: Real / Integer Linear Arithmetic

Theories Solver for RLA: The Simplex Method

Theories Solver for RLA: Fourier-Motzkin

Theories: Integer Linear Arithmetic

SMT: Recap

Theory Combinations

Context

Language

- Signature $\Sigma = \Sigma_F \cup \Sigma_P$

Language

- Signature $\Sigma = \Sigma_F \cup \Sigma_P$
- Every predicate $p \in \Sigma_P$ and function $f \in \Sigma_F$ has a fixed arity.

Language

- Signature $\Sigma = \Sigma_F \cup \Sigma_P$
- Every predicate $p \in \Sigma_P$ and function $f \in \Sigma_F$ has a fixed arity.
- Functions with arity 0 are called constants.

Language

- Signature $\Sigma = \Sigma_F \cup \Sigma_P$
- Every predicate $p \in \Sigma_P$ and function $f \in \Sigma_F$ has a fixed arity.
- Functions with arity 0 are called constants.
- Countable set of variables \mathcal{V} , disjoint from Σ .

Context

Language

- Signature $\Sigma = \Sigma_F \cup \Sigma_P$
- Every predicate $p \in \Sigma_P$ and function $f \in \Sigma_F$ has a fixed arity.
- Functions with arity 0 are called constants.
- Countable set of variables \mathcal{V} , disjoint from Σ .

Semantics

Context

Language

- Signature $\Sigma = \Sigma_F \cup \Sigma_P$
- Every predicate $p \in \Sigma_P$ and function $f \in \Sigma_F$ has a fixed arity.
- Functions with arity 0 are called constants.
- Countable set of variables \mathcal{V} , disjoint from Σ .

Semantics

- Σ -model (or structure): $\mathcal{A} = (A, \mathcal{I})$

Context

Language

- Signature $\Sigma = \Sigma_F \cup \Sigma_P$
- Every predicate $p \in \Sigma_P$ and function $f \in \Sigma_F$ has a fixed arity.
- Functions with arity 0 are called constants.
- Countable set of variables \mathcal{V} , disjoint from Σ .

Semantics

- Σ -model (or structure): $\mathcal{A} = (A, \mathcal{I})$
- A is the universe (or domain)

Context

Language

- Signature $\Sigma = \Sigma_F \cup \Sigma_P$
- Every predicate $p \in \Sigma_P$ and function $f \in \Sigma_F$ has a fixed arity.
- Functions with arity 0 are called constants.
- Countable set of variables \mathcal{V} , disjoint from Σ .

Semantics

- Σ -model (or structure): $\mathcal{A} = (A, \mathcal{I})$
- A is the universe (or domain)
- \mathcal{I} is the interpretation

Context – Models and Satisfiability

- A model \mathcal{A} satisfies ϕ iff ϕ evaluates to true under standard semantics

Context – Models and Satisfiability

- A model \mathcal{A} satisfies ϕ iff ϕ evaluates to true under standard semantics
- For convenience, free variables are treated as uninterpreted constants.

Context – Models and Satisfiability

- A model \mathcal{A} satisfies ϕ iff ϕ evaluates to true under standard semantics
- For convenience, free variables are treated as uninterpreted constants.
- In SMT, for a given theory \mathcal{T} , a formula ϕ is \mathcal{T} -satisfied by a model \mathcal{A} if \mathcal{A} satisfies ϕ and it also satisfies \mathcal{T} .

Context – Models and Satisfiability

- A model \mathcal{A} satisfies ϕ iff ϕ evaluates to true under standard semantics
- For convenience, free variables are treated as uninterpreted constants.
- In SMT, for a given theory \mathcal{T} , a formula ϕ is \mathcal{T} -satisfied by a model \mathcal{A} if \mathcal{A} satisfies ϕ and it also satisfies \mathcal{T} .
- A formula is \mathcal{T} -satisfiable if there is a model that \mathcal{T} -satisfies it.

Context – Models and Satisfiability

- A model \mathcal{A} satisfies ϕ iff ϕ evaluates to true under standard semantics
- For convenience, free variables are treated as uninterpreted constants.
- In SMT, for a given theory \mathcal{T} , a formula ϕ is \mathcal{T} -satisfied by a model \mathcal{A} if \mathcal{A} satisfies ϕ and it also satisfies \mathcal{T} .
- A formula is \mathcal{T} -satisfiable if there is a model that \mathcal{T} -satisfies it.
- $x < y < (x + 1)$ is \mathcal{T}_{LRA} -satisfiable (linear real arithmetic) with a model $x = 0, y = .5$.

Context – Models and Satisfiability

- A model \mathcal{A} satisfies ϕ iff ϕ evaluates to true under standard semantics
- For convenience, free variables are treated as uninterpreted constants.
- In SMT, for a given theory \mathcal{T} , a formula ϕ is \mathcal{T} -satisfied by a model \mathcal{A} if \mathcal{A} satisfies ϕ and it also satisfies \mathcal{T} .
- A formula is \mathcal{T} -satisfiable if there is a model that \mathcal{T} -satisfies it.
- $x < y < (x + 1)$ is \mathcal{T}_{LRA} -satisfiable (linear real arithmetic) with a model $x = 0, y = .5$.
- $x < y < (x + 1)$ is \mathcal{T}_{LIA} -unsatisfiable (linear integer arithmetic)

Context – Lazy solving

input : formula φ in theory \mathcal{T}

output: truth value

```
1  $\alpha \leftarrow \mathcal{T2B}(\varphi)$  // abstract input formula
2 while true do
3    $(\text{res}, \tau) \leftarrow \text{SAT}(\alpha)$  // Boolean model
4   if  $\text{res} = \text{false}$  then return false
5    $\mathcal{L} \leftarrow \bigcup_{l \in \tau} \mathcal{B2T}(l)$  // convert to theory model
6    $(\text{res}, \mathcal{L}') \leftarrow \mathcal{T}\text{-SAT}(\mathcal{L})$  // theory check
7   if  $\text{res} = \text{true}$  then return true
8    $\alpha \leftarrow \alpha \wedge \bigvee_{l \in \mathcal{L}'} \neg \mathcal{T2B}(l)$  // block explanation
9 end
```

Observe: Effectively enumerating sets of literals \mathcal{L} , s.t. $\mathcal{L} \Rightarrow \varphi$.
Theory solver decides if \mathcal{L} satisfiable.

Theory Combinations

- Until now we have assumed there is a single theory.

Theory Combinations

- Until now we have assumed there is a single theory.
- In practice important to combine theories.

$$(a = b+2) \wedge (A = \text{write}(B, a+1, 4)) \wedge (\text{read}(A, b+3) = 2 \vee f(a-1) \neq f(b+1))$$

Theory Combinations

- Until now we have assumed there is a single theory.
- In practice important to combine theories.

$$(a = b+2) \wedge (A = \textit{write}(B, a+1, 4)) \wedge (\textit{read}(A, b+3) = 2 \vee f(a-1) \neq f(b+1))$$

- **Nelson-Oppen method**: Use lazy solving and let the theory solvers communicate through equality.

Theory Combinations

- Until now we have assumed there is a single theory.
- In practice important to combine theories.

$$(a = b+2) \wedge (A = \text{write}(B, a+1, 4)) \wedge (\text{read}(A, b+3) = 2 \vee f(a-1) \neq f(b+1))$$

- **Nelson-Oppen method**: Use lazy solving and let the theory solvers communicate through equality.
- **Setup**: The theories \mathcal{T}_1 and \mathcal{T}_2 have the respective signatures:
 $\Sigma^1 = \Sigma_P^1 \cup \Sigma_F^1 \cup C$ and $\Sigma^2 = \Sigma_P^2 \cup \Sigma_F^2 \cup C$.

Theory Combinations

- Until now we have assumed there is a single theory.
- In practice important to combine theories.

$$(a = b+2) \wedge (A = \text{write}(B, a+1, 4)) \wedge (\text{read}(A, b+3) = 2 \vee f(a-1) \neq f(b+1))$$

- **Nelson-Oppen method**: Use lazy solving and let the theory solvers communicate through equality.
- **Setup**: The theories \mathcal{T}_1 and \mathcal{T}_2 have the respective signatures:
 $\Sigma^1 = \Sigma_P^1 \cup \Sigma_F^1 \cup C$ and $\Sigma^2 = \Sigma_P^2 \cup \Sigma_F^2 \cup C$.
- C is a set of constants and $\Sigma^1 \cap \Sigma^2 = C \cup \{=\}$.

Theory Combinations

- Until now we have assumed there is a single theory.
- In practice important to combine theories.

$$(a = b+2) \wedge (A = \text{write}(B, a+1, 4)) \wedge (\text{read}(A, b+3) = 2 \vee f(a-1) \neq f(b+1))$$

- **Nelson-Oppen method**: Use lazy solving and let the theory solvers communicate through equality.
- **Setup**: The theories \mathcal{T}_1 and \mathcal{T}_2 have the respective signatures:
 $\Sigma^1 = \Sigma_P^1 \cup \Sigma_F^1 \cup C$ and $\Sigma^2 = \Sigma_P^2 \cup \Sigma_F^2 \cup C$.
- C is a set of constants and $\Sigma^1 \cap \Sigma^2 = C \cup \{=\}$.
- Only equalities on C will be used to interact between the two respective theory solvers.

Theory Combinations

- Until now we have assumed there is a single theory.
- In practice important to combine theories.

$$(a = b+2) \wedge (A = \text{write}(B, a+1, 4)) \wedge (\text{read}(A, b+3) = 2 \vee f(a-1) \neq f(b+1))$$

- **Nelson-Oppen method**: Use lazy solving and let the theory solvers communicate through equality.
- **Setup**: The theories \mathcal{T}_1 and \mathcal{T}_2 have the respective signatures:
 $\Sigma^1 = \Sigma_P^1 \cup \Sigma_F^1 \cup C$ and $\Sigma^2 = \Sigma_P^2 \cup \Sigma_F^2 \cup C$.
- C is a set of constants and $\Sigma^1 \cap \Sigma^2 = C \cup \{=\}$.
- Only equalities on C will be used to interact between the two respective theory solvers.
- We want to decide a set of literals φ on the signature $\Sigma^1 \cup \Sigma^2$.

First Step: Purification

- Iteratively **replace** each sub-term t by a fresh constant $c \in C$ and add the literal $c = t$.

First Step: Purification

- Iteratively **replace** each sub-term t by a fresh constant $c \in C$ and add the literal $c = t$.
- **Separate** into two sets of literals φ_1, φ_2 , so that φ_i is on the signature Σ^i .

First Step: Purification

- Iteratively **replace** each sub-term t by a fresh constant $c \in C$ and add the literal $c = t$.
- **Separate** into two sets of literals φ_1, φ_2 , so that φ_i is on the signature Σ^i .

Example

First Step: Purification

- Iteratively **replace** each sub-term t by a fresh constant $c \in C$ and add the literal $c = t$.
- **Separate** into two sets of literals φ_1, φ_2 , so that φ_i is on the signature Σ^i .

Example

- \mathcal{T}_E ...theory of the equality (EUF), $\mathcal{T}_{\mathbb{Z}}$...theory of integers.

First Step: Purification

- Iteratively **replace** each sub-term t by a fresh constant $c \in C$ and add the literal $c = t$.
- **Separate** into two sets of literals φ_1, φ_2 , so that φ_i is on the signature Σ^i .

Example

- \mathcal{T}_E ...theory of the equality (EUF), \mathcal{T}_Z ...theory of integers.
- $1 \leq x \wedge x \leq 2 \wedge f(x) \neq f(1) \wedge f(x) \neq f(2)$

First Step: Purification

- Iteratively **replace** each sub-term t by a fresh constant $c \in C$ and add the literal $c = t$.
- **Separate** into two sets of literals φ_1, φ_2 , so that φ_i is on the signature Σ^i .

Example

- \mathcal{T}_E ...theory of the equality (EUF), \mathcal{T}_Z ...theory of integers.
- $1 \leq x \wedge x \leq 2 \wedge f(x) \neq f(1) \wedge f(x) \neq f(2)$
- $w_1 \leq x \wedge x \leq w_2 \wedge f(x) \neq f(w_1) \wedge f(x) \neq f(w_2) \wedge w_1 = 1 \wedge w_2 = 2$

First Step: Purification

- Iteratively **replace** each sub-term t by a fresh constant $c \in C$ and add the literal $c = t$.
- Separate** into two sets of literals φ_1, φ_2 , so that φ_i is on the signature Σ^i .

Example

- $\mathcal{T}_{\mathbb{E}}$...theory of the equality (EUF), $\mathcal{T}_{\mathbb{Z}}$...theory of integers.
- $1 \leq x \wedge x \leq 2 \wedge f(x) \neq f(1) \wedge f(x) \neq f(2)$
- $w_1 \leq x \wedge x \leq w_2 \wedge f(x) \neq f(w_1) \wedge f(x) \neq f(w_2) \wedge w_1 = 1 \wedge w_2 = 2$
- $w_1 \leq x \wedge x \leq w_2 \wedge w_1 = 1 \wedge w_2 = 2$ and
 $f(x) \neq f(w_1) \wedge f(x) \neq f(w_2)$

First Step: Purification

- Iteratively **replace** each sub-term t by a fresh constant $c \in C$ and add the literal $c = t$.
- Separate** into two sets of literals φ_1, φ_2 , so that φ_i is on the signature Σ^i .

Example

- $\mathcal{T}_{\mathbb{E}}$...theory of the equality (EUF), $\mathcal{T}_{\mathbb{Z}}$...theory of integers.
- $1 \leq x \wedge x \leq 2 \wedge f(x) \neq f(1) \wedge f(x) \neq f(2)$
- $w_1 \leq x \wedge x \leq w_2 \wedge f(x) \neq f(w_1) \wedge f(x) \neq f(w_2) \wedge w_1 = 1 \wedge w_2 = 2$
- $w_1 \leq x \wedge x \leq w_2 \wedge w_1 = 1 \wedge w_2 = 2$ and
 $f(x) \neq f(w_1) \wedge f(x) \neq f(w_2)$
- Simplify $1 \leq x \wedge x \leq 2 \wedge w_1 = 1 \wedge w_2 = 2$ and
 $f(x) \neq f(w_1) \wedge f(x) \neq f(w_2)$

Second Step: Guess and Check

- For the separation φ_1, φ_2 consider all the shared constants X .

Second Step: Guess and Check

- For the separation φ_1, φ_2 consider all the shared constants X .
- **Guess** some equivalence relation R on X .

Second Step: Guess and Check

- For the separation φ_1, φ_2 consider all the shared constants X .
- **Guess** some equivalence relation R on X .
- Build the **arrangement formula**:

$$\alpha(X, R) = \bigwedge_{u, v \in X, uRv} u = v \wedge \bigwedge_{u, v \in X, \neg uRv} u \neq v$$

Second Step: Guess and Check

- For the separation φ_1, φ_2 consider all the shared constants X .
- **Guess** some equivalence relation R on X .
- Build the **arrangement formula**:

$$\alpha(X, R) = \bigwedge_{u, v \in X, uRv} u = v \wedge \bigwedge_{u, v \in X, \neg uRv} u \neq v$$

- **Check** satisfiability of $\varphi_1 \wedge \alpha(X, R)$ and $\varphi_2 \wedge \alpha(X, R)$.

Second Step: Guess and Check

- For the separation φ_1, φ_2 consider all the shared constants X .
- **Guess** some equivalence relation R on X .
- Build the **arrangement formula**:

$$\alpha(X, R) = \bigwedge_{u, v \in X, uRv} u = v \wedge \bigwedge_{u, v \in X, \neg uRv} u \neq v$$

- **Check** satisfiability of $\varphi_1 \wedge \alpha(X, R)$ and $\varphi_2 \wedge \alpha(X, R)$.
- $\varphi_1 \wedge \varphi_2$ is satisfiable iff the we get satisfiability for some R .

Guess and Check: Unsatisfiable Example

- $1 \leq x \wedge x \leq 2 \wedge w_1 = 1 \wedge w_2 = 2$ and
 $f(x) \neq f(w_1) \wedge f(x) \neq f(w_2)$

Guess and Check: Unsatisfiable Example

- $1 \leq x \wedge x \leq 2 \wedge w_1 = 1 \wedge w_2 = 2$ and
 $f(x) \neq f(w_1) \wedge f(x) \neq f(w_2)$
- Possible partitions: $\{\{x, w_1, w_2\}\}$ $\{\{x, w_1\}, \{w_2\}\}$
 $\{\{x, w_2\}, \{w_1\}\}$ $\{\{x\}, \{w_1, w_2\}\}$ $\{\{x\}, \{w_1\}, \{w_2\}\}$

Guess and Check: Unsatisfiable Example

- $1 \leq x \wedge x \leq 2 \wedge w_1 = 1 \wedge w_2 = 2$ and
 $f(x) \neq f(w_1) \wedge f(x) \neq f(w_2)$
- Possible partitions: $\{\{x, w_1, w_2\}\} \{\{x, w_1\}, \{w_2\}\}$
 $\{\{x, w_2\}, \{w_1\}\} \{\{x\}, \{w_1, w_2\}\} \{\{x\}, \{w_1\}, \{w_2\}\}$
- $x = w_1 = w_2$ UNSAT in $\mathcal{T}_{\mathbb{Z}}$ because $w_1 = 1 \wedge w_2 = 2$.

Guess and Check: Unsatisfiable Example

- $1 \leq x \wedge x \leq 2 \wedge w_1 = 1 \wedge w_2 = 2$ and
 $f(x) \neq f(w_1) \wedge f(x) \neq f(w_2)$
- Possible partitions: $\{\{x, w_1, w_2\}\} \{\{x, w_1\}, \{w_2\}\}$
 $\{\{x, w_2\}, \{w_1\}\} \{\{x\}, \{w_1, w_2\}\} \{\{x\}, \{w_1\}, \{w_2\}\}$
- $x = w_1 = w_2$ UNSAT in $\mathcal{T}_{\mathbb{Z}}$ because $w_1 = 1 \wedge w_2 = 2$.
- $x = w_1 \wedge x \neq w_2$ UNSAT in $\mathcal{T}_{\mathbb{E}}$ because $f(x) \neq f(w_1)$.

Guess and Check: Unsatisfiable Example

- $1 \leq x \wedge x \leq 2 \wedge w_1 = 1 \wedge w_2 = 2$ and $f(x) \neq f(w_1) \wedge f(x) \neq f(w_2)$
- Possible partitions: $\{\{x, w_1, w_2\}\} \{\{x, w_1\}, \{w_2\}\} \{\{x, w_2\}, \{w_1\}\} \{\{x\}, \{w_1, w_2\}\} \{\{x\}, \{w_1\}, \{w_2\}\}$
- $x = w_1 = w_2$ UNSAT in $\mathcal{T}_{\mathbb{Z}}$ because $w_1 = 1 \wedge w_2 = 2$.
- $x = w_1 \wedge x \neq w_2$ UNSAT in $\mathcal{T}_{\mathbb{E}}$ because $f(x) \neq f(w_1)$.
- $x = w_2 \wedge x \neq w_1$ UNSAT in $\mathcal{T}_{\mathbb{E}}$ because $f(x) \neq f(w_2)$.

Guess and Check: Unsatisfiable Example

- $1 \leq x \wedge x \leq 2 \wedge w_1 = 1 \wedge w_2 = 2$ and $f(x) \neq f(w_1) \wedge f(x) \neq f(w_2)$
- Possible partitions: $\{\{x, w_1, w_2\}\} \{\{x, w_1\}, \{w_2\}\} \{\{x, w_2\}, \{w_1\}\} \{\{x\}, \{w_1, w_2\}\} \{\{x\}, \{w_1\}, \{w_2\}\}$
- $x = w_1 = w_2$ UNSAT in $\mathcal{T}_{\mathbb{Z}}$ because $w_1 = 1 \wedge w_2 = 2$.
- $x = w_1 \wedge x \neq w_2$ UNSAT in $\mathcal{T}_{\mathbb{E}}$ because $f(x) \neq f(w_1)$.
- $x = w_2 \wedge x \neq w_1$ UNSAT in $\mathcal{T}_{\mathbb{E}}$ because $f(x) \neq f(w_2)$.
- $x \neq w_1 \wedge w_1 = w_2$ UNSAT in $\mathcal{T}_{\mathbb{Z}}$ because $w_1 = 1 \wedge w_2 = 2$.

Guess and Check: Unsatisfiable Example

- $1 \leq x \wedge x \leq 2 \wedge w_1 = 1 \wedge w_2 = 2$ and $f(x) \neq f(w_1) \wedge f(x) \neq f(w_2)$
- Possible partitions: $\{\{x, w_1, w_2\}\} \{\{x, w_1\}, \{w_2\}\} \{\{x, w_2\}, \{w_1\}\} \{\{x\}, \{w_1, w_2\}\} \{\{x\}, \{w_1\}, \{w_2\}\}$
- $x = w_1 = w_2$ UNSAT in $\mathcal{T}_{\mathbb{Z}}$ because $w_1 = 1 \wedge w_2 = 2$.
- $x = w_1 \wedge x \neq w_2$ UNSAT in $\mathcal{T}_{\mathbb{E}}$ because $f(x) \neq f(w_1)$.
- $x = w_2 \wedge x \neq w_1$ UNSAT in $\mathcal{T}_{\mathbb{E}}$ because $f(x) \neq f(w_2)$.
- $x \neq w_1 \wedge w_1 = w_2$ UNSAT in $\mathcal{T}_{\mathbb{Z}}$ because $w_1 = 1 \wedge w_2 = 2$.
- $x \neq w_1 \wedge x \neq w_2 \wedge w_1 \neq w_2$ UNSAT in $\mathcal{T}_{\mathbb{Z}}$ because it cannot be that $x \neq 1 \wedge x \neq 2 \wedge 1 \leq x \leq 2$.

Guess and Check: Satisfiable Example

- $f(x) = x + y \wedge x \leq y + z \wedge x + z \leq y \wedge y = 1 \wedge f(x) \neq f(2)$.

Guess and Check: Satisfiable Example

- $f(x) = x + y \wedge x \leq y + z \wedge x + z \leq y \wedge y = 1 \wedge f(x) \neq f(2)$.
- $w_1 = x + y \wedge x \leq y + z \wedge x + z \leq y \wedge y = 1 \wedge w_2 = 2$ and
 $w_1 = f(x) \wedge f(x) \neq f(w_2)$

Guess and Check: Satisfiable Example

- $f(x) = x + y \wedge x \leq y + z \wedge x + z \leq y \wedge y = 1 \wedge f(x) \neq f(2)$.
- $w_1 = x + y \wedge x \leq y + z \wedge x + z \leq y \wedge y = 1 \wedge w_2 = 2$ and
 $w_1 = f(x) \wedge f(x) \neq f(w_2)$
- Shared: x, w_1, w_2 .

Guess and Check: Satisfiable Example

- $f(x) = x + y \wedge x \leq y + z \wedge x + z \leq y \wedge y = 1 \wedge f(x) \neq f(2)$.
- $w_1 = x + y \wedge x \leq y + z \wedge x + z \leq y \wedge y = 1 \wedge w_2 = 2$ and
 $w_1 = f(x) \wedge f(x) \neq f(w_2)$
- Shared: x, w_1, w_2 .
- Simplify as $w_1 = x + 1 \wedge x \leq 1 + z \wedge x + z \leq 1 \wedge y = 1 \wedge w_2 = 2$
and $w_1 = f(x) \wedge f(x) \neq f(w_2)$

Guess and Check: Satisfiable Example

- $f(x) = x + y \wedge x \leq y + z \wedge x + z \leq y \wedge y = 1 \wedge f(x) \neq f(2)$.
- $w_1 = x + y \wedge x \leq y + z \wedge x + z \leq y \wedge y = 1 \wedge w_2 = 2$ and $w_1 = f(x) \wedge f(x) \neq f(w_2)$
- Shared: x, w_1, w_2 .
- Simplify as $w_1 = x + 1 \wedge x \leq 1 + z \wedge x + z \leq 1 \wedge y = 1 \wedge w_2 = 2$ and $w_1 = f(x) \wedge f(x) \neq f(w_2)$
- $x = w_1$ gives contradiction with $w_1 = x + 1$.

Guess and Check: Satisfiable Example

- $f(x) = x + y \wedge x \leq y + z \wedge x + z \leq y \wedge y = 1 \wedge f(x) \neq f(2)$.
- $w_1 = x + y \wedge x \leq y + z \wedge x + z \leq y \wedge y = 1 \wedge w_2 = 2$ and $w_1 = f(x) \wedge f(x) \neq f(w_2)$
- Shared: x, w_1, w_2 .
- Simplify as $w_1 = x + 1 \wedge x \leq 1 + z \wedge x + z \leq 1 \wedge y = 1 \wedge w_2 = 2$ and $w_1 = f(x) \wedge f(x) \neq f(w_2)$
- $x = w_1$ gives contradiction with $w_1 = x + 1$.
- $x = w_2$ gives contradiction with $f(x) \neq f(w_2)$.

Guess and Check: Satisfiable Example

- $f(x) = x + y \wedge x \leq y + z \wedge x + z \leq y \wedge y = 1 \wedge f(x) \neq f(2)$.
- $w_1 = x + y \wedge x \leq y + z \wedge x + z \leq y \wedge y = 1 \wedge w_2 = 2$ and $w_1 = f(x) \wedge f(x) \neq f(w_2)$
- Shared: x, w_1, w_2 .
- Simplify as $w_1 = x + 1 \wedge x \leq 1 + z \wedge x + z \leq 1 \wedge y = 1 \wedge w_2 = 2$ and $w_1 = f(x) \wedge f(x) \neq f(w_2)$
- $x = w_1$ gives contradiction with $w_1 = x + 1$.
- $x = w_2$ gives contradiction with $f(x) \neq f(w_2)$.
- x has to be in its own equivalence class, i.e. $x \neq w_1, x \neq w_2$.

Guess and Check: Satisfiable Example

- $f(x) = x + y \wedge x \leq y + z \wedge x + z \leq y \wedge y = 1 \wedge f(x) \neq f(2)$.
- $w_1 = x + y \wedge x \leq y + z \wedge x + z \leq y \wedge y = 1 \wedge w_2 = 2$ and $w_1 = f(x) \wedge f(x) \neq f(w_2)$
- Shared: x, w_1, w_2 .
- Simplify as $w_1 = x + 1 \wedge x \leq 1 + z \wedge x + z \leq 1 \wedge y = 1 \wedge w_2 = 2$ and $w_1 = f(x) \wedge f(x) \neq f(w_2)$
- $x = w_1$ gives contradiction with $w_1 = x + 1$.
- $x = w_2$ gives contradiction with $f(x) \neq f(w_2)$.
- x has to be in its own equivalence class, i.e. $x \neq w_1, x \neq w_2$.
- Adding $w_1 = w_2$ permits models:
 $\{x = 1, z = 0, y = 1, w_1 = w_2 = 2\}$
 $\{w_1 = w_2 = *_1, x = *_2, f = \{*_1 \mapsto *_2, *_2 \mapsto *_1\}\}$

Guess and Check: Satisfiable Example

- $f(x) = x + y \wedge x \leq y + z \wedge x + z \leq y \wedge y = 1 \wedge f(x) \neq f(2)$.
- $w_1 = x + y \wedge x \leq y + z \wedge x + z \leq y \wedge y = 1 \wedge w_2 = 2$ and $w_1 = f(x) \wedge f(x) \neq f(w_2)$
- Shared: x, w_1, w_2 .
- Simplify as $w_1 = x + 1 \wedge x \leq 1 + z \wedge x + z \leq 1 \wedge y = 1 \wedge w_2 = 2$ and $w_1 = f(x) \wedge f(x) \neq f(w_2)$
- $x = w_1$ gives contradiction with $w_1 = x + 1$.
- $x = w_2$ gives contradiction with $f(x) \neq f(w_2)$.
- x has to be in its own equivalence class, i.e. $x \neq w_1, x \neq w_2$.
- Adding $w_1 = w_2$ permits models:
 $\{x = 1, z = 0, y = 1, w_1 = w_2 = 2\}$
 $\{w_1 = w_2 = *_1, x = *_2, f = \{*_1 \mapsto *_2, *_2 \mapsto *_1\}\}$
- Merged model
 $\{x = y = 1, z = 0, w_1 = w_2 = 2, f = \{2 \mapsto 1, 1 \mapsto 2, \text{rest arbitrary}\}\}$

Merging

- For the models (A_1, \mathcal{I}_1) and (A_2, \mathcal{I}_2) build a bijection h between A_2 and A_1 so that $h(\mathcal{I}_2(c)) = \mathcal{I}_1(c)$ for all shared constants c .

Merging

- For the models (A_1, \mathcal{I}_1) and (A_2, \mathcal{I}_2) build a bijection h between A_2 and A_1 so that $h(\mathcal{I}_2(c)) = \mathcal{I}_1(c)$ for all shared constants c .
- Such h exists because both models must respect the guessed equivalence relation.

Merging

- For the models (A_1, \mathcal{I}_1) and (A_2, \mathcal{I}_2) build a bijection h between A_2 and A_1 so that $h(\mathcal{I}_2(c)) = \mathcal{I}_1(c)$ for all shared constants c .
- Such h exists because both models must respect the guessed equivalence relation.

Example

Merging

- For the models (A_1, \mathcal{I}_1) and (A_2, \mathcal{I}_2) build a bijection h between A_2 and A_1 so that $h(\mathcal{I}_2(c)) = \mathcal{I}_1(c)$ for all shared constants c .
- Such h exists because both models must respect the guessed equivalence relation.

Example

- Models: $\mathcal{I}_1 = \{x = 1, z = 0, y = 1, w_1 = w_2 = 2\}$
 $\mathcal{I}_2 = \{w_1 = w_2 = *_1, x = *_2, f = \{*_1 \mapsto *_2, *_2 \mapsto *_1\}\}$

Merging

- For the models (A_1, \mathcal{I}_1) and (A_2, \mathcal{I}_2) build a bijection h between A_2 and A_1 so that $h(\mathcal{I}_2(c)) = \mathcal{I}_1(c)$ for all shared constants c .
- Such h exists because both models must respect the guessed equivalence relation.

Example

- Models: $\mathcal{I}_1 = \{x = 1, z = 0, y = 1, w_1 = w_2 = 2\}$
 $\mathcal{I}_2 = \{w_1 = w_2 = *_1, x = *_2, f = \{*_1 \mapsto *_2, *_2 \mapsto *_1\}\}$
- $h(*_1) = 2, h(*_2) = 1$, rest some arbitrary bijection.

Nelson-Oppen: When does it work?

- Nelson-Oppen **does not** to work for arbitrary theories.

Nelson-Oppen: When does it work?

- Nelson-Oppen **does not** to work for arbitrary theories.
- Theories needs to be **stably infinite**

Nelson-Oppen: When does it work?

- Nelson-Oppen **does not** to work for arbitrary theories.
- Theories needs to be **stably infinite**
 - A theory \mathcal{T} is stably infinite iff for every satisfiable $\Sigma_{\mathcal{T}}$ -formula ϕ , there is a \mathcal{T} -model that satisfies ϕ and that has a universe of infinite cardinality

Nelson-Oppen: When does it work?

- Nelson-Oppen **does not** to work for arbitrary theories.
- Theories needs to be **stably infinite**
 - A theory \mathcal{T} is stably infinite iff for every satisfiable $\Sigma_{\mathcal{T}}$ -formula ϕ , there is a \mathcal{T} -model that satisfies ϕ and that has a universe of infinite cardinality
 - **Examples:** EUF, LIA, LRA, etc.

Nelson-Oppen: When does it work?

- Nelson-Oppen **does not** to work for arbitrary theories.
- Theories needs to be **stably infinite**
 - A theory \mathcal{T} is stably infinite iff for every satisfiable $\Sigma_{\mathcal{T}}$ -formula ϕ , there is a \mathcal{T} -model that satisfies ϕ and that has a universe of infinite cardinality
 - **Examples:** EUF, LIA, LRA, etc.
- Combination of more than two theories is done analogously.

Nelson-Oppen: When does it work?

- Nelson-Oppen **does not** to work for arbitrary theories.
- Theories needs to be **stably infinite**
 - A theory \mathcal{T} is stably infinite iff for every satisfiable $\Sigma_{\mathcal{T}}$ -formula ϕ , there is a \mathcal{T} -model that satisfies ϕ and that has a universe of infinite cardinality
 - **Examples:** EUF, LIA, LRA, etc.
- Combination of more than two theories is done analogously.
- For certain theories there are more efficient methods than considering all partitions (convex theories).

Summary

- Theory combinations:

- Theory combinations:
 - Nelson-Oppen method enables combining theory solvers (under certain conditions).

- Theory combinations:
 - Nelson-Oppen method enables combining theory solvers (under certain conditions).
 - Theory solvers communicate through equality.