

Test case 2: Two-dimensional spinal cord

Problem statement

Dirichlet boundary:

- $u = 0$ on Γ_{pia} , $u_y = 0$ on $\Gamma_{\text{top,bot}}$
- $p = p_{\text{applied}}$ on Γ

Parameters:

	E Young's modulus	ν Poisson's ratio	Λ	Fluid mobility	b Fluid storage coefficient	τ Coupling strength
Pia mater	$2 \cdot 10^4$	0.35	$4 \cdot 10^{-6}$	$1 \cdot 10^{-5}$	4.5	
White matter	$5 \cdot 10^1$	0.35	$2 \cdot 10^{-5}$	$1 \cdot 10^{-2}$	1.8	
Grey matter	$5 \cdot 10^1$	0.35	$2 \cdot 10^{-6}$	$1 \cdot 10^{-2}$	1.8	
Central canal	$5 \cdot 10^1$	0.35	$1 \cdot 10^{-3}$	$1 \cdot 10^{-2}$	1.8	




Figure 1: 2D spinal cord problem. The parameters are as shown in the figure, except that $b = 0$ everywhere (giving $\tau \rightarrow \infty$), and $\Delta t = 10^{-3}$.

Jan, Joachim has run a bunch of simulations and we have looked through them in order to find a reasonable story. Joachim, I already lost the sketch due to moving between offices etc. In planned to write this in google docs, but was not able to upload the pdfs as images.

Anyway, here goes the story.

Pictures 2–3 shows that Krylov methods are superior to Richardson. This is not a big surprise, but provides a clear link to the splitting schemes.

The figures also show both the error (in energy norm) and the residual for BiCGStab and LGMRES. Here dotted lines are residual and the solid line is energy. Here, it is clear that it is a close correspondance between error and residual for LGMRES and less clear for BiCGStab. We therefore continue the discussion with only LGMRES and error.

JBH: It is not immediately obvious that the Krylov methods are very superior. Given that BiCGStab is about twice as expensive per iteration, and LGMRES is 30+ times as expensive per iteration, they may something like 30% faster. What is clear is: (a) They are reasonable preconditioners, compared to the bog-standard Block Jacobi method; (b) Even the unstable splittings converge, at least in LGMRES; and (c) Error is much better controlled in LGMRES than in either Richardson or BiCGStab.

Figure 4 shows that the same pattern holds when the splitting is performed using inexact single-block solvers.

Next, lets consider the eigenvalues of the individual components.




Figure 2: Richardson iterations (exact subsolves) is equivalent to the traditional splitting schemes. The residual is shown as solid lines; energy-norm error as dotted lines. Coupling strength $\tau \rightarrow \infty$ when $b \rightarrow 0$, so the unstable splittings diverge quickly. (Undrained split is not included in the comparison, because it is undefined for $b = 0$.)




Figure 3: BiCGStab and LGMRES (exact subsolves). The residuals is shown as solid lines; energy-norm error as dotted lines. For LGMRES, the energy-norm error is (nearly?) invisible because it overlaps with the residual lines.




Figure 4: Richardson and LGMRES with inexact subsolves (1 cycle of ML).




Figure 5: Eigenvalues of the preconditioned A block at $N = 32$ (left) and $N = 64$ (right).




Figure 6: Eigenvalues of the preconditioned $-C$ block at $N = 32$ (left) and $N = 64$ (right).




Figure 7: Eigenvalues of the preconditioned $-S_C$ block at $N = 32$ (left) and $N = 64$ (right).

Figures 5–7 shows the preconditioned eigenvalues of the preconditioned single blocks. Clearly, the eigenvalues here are nicely bounded. We may here investigate how the number of ML sweeps affect the eigenvalues.

This more or less concludes the discussion of the 2D spinal cord example.

The contrast example is shown in figure 8. Figures 9 and 10 show the convergence of Richardson and LGMRES on this example, and figures 11–11 show the component eigenvalues.

Only the (Schur-based) fixed stress preconditioner is effective in this example. This is expected from earlier works: preconditioning based on the unmodified C block does not work with high contrast. Furthermore, for the Fixed Stress method, the convergence is similar to the 2D spinal cord example for both Richardson and LGMRES.

The methods do NOT show any significant grid dependence. I did see grid dependence earlier, but that was probably (mostly) caused by the ILU preconditioner on the C/S_C block.

JBH: Can we drop everything except fixed stress in the rest of the examples?

Test case 4: Pore seal


Problem statement

Dirichlet boundary:

- $u = 0$ on Γ_{bottom}

Parameters:

	E Young's modulus	ν Poisson's ratio	Λ	Fluid mobility	b Fluid storage coefficient	τ Coupling strength
Top half	$3 \cdot 10^3$	0.45	$1 \cdot 10^{-8}$	$1 \cdot 10^{-6}$	91	
Bottom half	$3 \cdot 10^3$	0.45	1	$1 \cdot 10^{-6}$	91	



Designed to be hard!



Figure 8: Contrast problem. There are differences in the parameters. Again $b = 0$ everywhere (giving $\tau \rightarrow \infty$), and $\Delta t = 10^{-3}$. Also, $p = 0$ at the top, and E is about 1/10 of that shown here. These changes are arbitrary: I can create new figures with the same parameters if necessary.




Figure 9: Contrast problem ($\delta = 10^{-8}$), Richardson, $N = 40$ (left) and $N = 160$ (right).




Figure 10: Contrast problem ($\delta = 10^{-8}$), LGMRES, $N = 40$ (left) and $N = 160$ (right).




Figure 11: Contrast problem ($\delta = 10^{-8}$). Eigenvalues, $N = 40$ (left) and $N = 160$ (right).




Figure 12: Contrast problem ($\delta = 10^{-8}$). Eigenvalues, $N = 40$ (left) and $N = 160$ (right).

When the contrast is increased to 10^{-12} , something unexpected and bad happens with LGMRES: It fails to converge at high resolution, but the BiCGStab/Richardson methods don't! See figures 13–15. My only guess is that this is some kind of numerical sensitivity in the LGMRES method that may even be implementation dependent.

JBH: This is a nasty surprise. I don't know, maybe we should just forget about LGMRES and focus on BiCGStab after all. We can show decent and grid-independent results on BiCGStab, but we haven't got any "killer" examples.




Figure 13: Contrast problem ($\delta = 10^{-12}$), Richardson, $N = 40$ (left) and $N = 640$ (right). (NOTE: cycles=2)




Figure 14: Contrast problem ($\delta = 10^{-12}$), LGMRES, $N = 40$ (left) and $N = 640$ (right). (NOTE: cycles=2)




Figure 15: Contrast problem ($\delta = 10^{-12}$), LGMRES, $N = 40$ (left) and $N = 640$ (right).




Figure 16: Contrast problem ($\delta = 10^{-12}$). Eigenvalues, $N = 40$ (left) and $N = 640$ (right).




Figure 17: Contrast problem ($\delta = 10^{-12}$). Eigenvalues, $N = 40$ (left) and $N = 640$ (right).

The end, for now.