

AST 1430

Cosmology

Week Index	Dates	Topics
Keith	Week 1	logistics, introduction, basic observations
	Week 2	Basic GR, RW metric, Distances, coordinates, Friedmann equations, cosmological models
	Week 3	Consistency with observations, Early Hot Universe, BBN
	Week 4	Inflation, Perturbations & Structure pre-recombination
	Week 5	CMB: basics, polarization, secondaries
	Week 6	Early-Universe Presentations
Reading Week – No Class		
Jo	Week 7	Post-recombination growth of structure, formation of dark matter halos, halo mass function
	Week 8	The relation between dark matter halos and galaxies
	Week 9	Probing the cosmic density field / clustering
	Week 10	Late Universe Presentations; Icosmological observations: BAO, supernovae, weak lensing, etc.
	Week 11	H0 controversy: how fast exactly is the Universe expanding today?
	Week 12	Review

asst1

asst2


asst3

Late-time cosmological observations


Why late-time cosmological observations?

- CMB measured in exquisite detail, with highly precise and accurate constraints on cosmological parameters
- So what do we need late-time constraints for?
 - Without assuming flatness, CMB alone cannot determine cosmological parameters well (i.e., need to assume Ω_k , or H_0 , or similar)
 - Extensions to the basic flat LCDM model:
 - Dark-energy with non-trivial equation of state (CMB not really sensitive to DE at all)
 - Effective number of neutrino species
 - Neutrino masses
 - Non-standard dark matter


The CMB




Planck 2018 VI (2020)




Planck 2018 VI (2020)



Planck 2018 VI (2020)




Planck 2018 VI (2020)



Physical density parameters
Optical depth reionization
Acoustic scale

Initial fluctuation spectrum

CMB degeneracy with flatness / H_0



Planck 2018 VI (2020)


CMB constraints

- Geometry vs. Growth-of-structure
- Geometry:
 - Physical baryon and dark-matter density: $\Omega_b h^2$ and $\Omega_c h^2$
 - Angular acoustic scale
 - Angular diameter distance to last scattering: $D_M(z=1090)/r_d$
- Growth-of-structure:
 - Amplitude and shape of the initial curvature power spectrum: n_s , A_s
 - Optical depth to reionization

Supernovae

Supernova constraints

- Supernovae allow measurements of absolute distance —> constrain distance-redshift relation
- Distance derived from standardizable luminosity —> measure luminosity distance $d_L(z)$
$$d_L(z) = (1 + z)c \int_0^z \frac{dz'}{H(z')},$$
- Two uses:
 - Measure H_0 (requires absolute calibration)
 - Measure redshift derivatives of $H(z)$ (does not require absolute calibration)



Riess et al. (1996)

Supernovae and the accelerating Universe

- Expansion of the luminosity distance at low redshift

$$d_L(z) \approx \frac{c}{H_0} \left[z + z^2 \left(\frac{1 - q_0}{2} \right) + \mathcal{O}(z^3) \right]$$


- q_0 is the *deceleration parameter*, given by

$$q_0 = -\frac{\ddot{a}/a}{\dot{a}^2} = \frac{1}{2} \sum_i \Omega_i (1 + 3w_i)$$


where (Ω_i, w_i) are the density equation-of-state parameter of component i (e.g., $w_i = 0$ for matter, $w_i = -1$ for a cosmological constant)

- E.g., LCDM

$$q_0 = \frac{1}{2} \Omega_m - \Omega_\Lambda$$



Riess et al. (1996)




Brout et al. (2022: Pantheon+)

Cosmological constraints from supernovae

$$q_0 = \frac{1}{2} \Omega_m - \Omega_\Lambda$$

- Measurement $q_0 \approx -0.55$
- Contours tilted with respect to the $q_0 = \text{constant}$ line, because measurements are not at $z=0$




Kowalski et al. (2008)

Cosmological constraints from supernovae

$$q_0 = \frac{1}{2} \Omega_m - \Omega_\Lambda$$

- Measurement $q_0 \approx -0.55$
- Contours tilted with respect to the $q_0 = \text{constant}$ line, because measurements are not at $z=0$
- Combining with CMB shows that the Universe is flat
- Supernovae constraints are pure geometry, no growth of structure




Kowalski et al. (2008)

Baryon acoustic oscillations

BAO constraints


- BAO measurements determine Hubble function $H(z)/r_d$ and comoving angular diameter distance $D_M(z)/r_d$ at the median redshift of a survey
- CMB measurements of $\Omega_b h^2$ and $\Omega_c h^2$ calibrate the BAO ruler r_d , so turn BAO measurements into absolute distance measurements



Aubourg et al. (2015)

BAO constraints


- BAO measurements determine Hubble function $H(z)/r_d$ and comoving angular diameter distance $D_M(z)/r_d$ at the median redshift of a survey
- CMB measurements of $\Omega_b h^2$ and $\Omega_c h^2$ calibrate the BAO ruler r_d , so turn BAO measurements into absolute distance measurements



Aubourg et al. (2015)

BAO constraints


- BAO measurements determine Hubble function $H(z)/r_d$ and co-moving angular diameter distance $D_M(z)/r_d$ at the median redshift of a survey
- CMB measurements of $\Omega_b h^2$ and $\Omega_c h^2$ calibrate the BAO ruler r_d , so turn BAO measurements into absolute distance measurements



Aubourg et al. (2015)

BAO constraints


- BAO measurements determine Hubble function $H(z)/r_d$ and co-moving angular diameter distance $D_M(z)/r_d$ at the median redshift of a survey
- CMB measurements of $\Omega_b h^2$ and $\Omega_c h^2$ calibrate the BAO ruler r_d , so turn BAO measurements into absolute distance measurements



Kowalski et al. (2008)

BAO constraints

- BAO measurements determine Hubble function $H(z)/r_d$ and comoving angular diameter distance $D_M(z)/r_d$ at the median redshift of a survey
- CMB measurements of $\Omega_b h^2$ and $\Omega_c h^2$ calibrate the BAO ruler r_d , so turn BAO measurements into absolute distance measurements




Brout et al. (2022: Pantheon+)

Combined constraints

Flatness


- Combination of CMB + SNe or BAO strongly constrains curvature parameter Ω_k
- Planck 2018 + BAO:
 $\Omega_k = 0.0007^{+0.0037}_{-0.0037}$



Planck 2018 VI (2020)

Flatness


- Combination of CMB + SNe or BAO strongly constrains curvature parameter Ω_k
- Planck 2018 + BAO:
 $\Omega_k = 0.0007^{+0.0037}_{-0.0037}$
- Supernovae don't add much



Aubourg et al. (2015)

Dark energy that is not a cosmological constant

- Simplest model has DE equation of state: $p = w \rho$ with constant w ($w=-1 \rightarrow$ cosmological constant)
- Little constraint from Planck alone




Planck 2018 VI (2020)

Dark energy that is not a cosmological constant

- Simplest model has DE equation of state: $p = w \rho$ with constant w ($w=-1 \rightarrow$ cosmological constant)
- Little constraint from Planck alone
- SNe + Planck: tight constraint

$$w = -0.982^{+0.022}_{-0.038}$$





Brout et al. (2022: Pantheon+)

Dark energy that is not a cosmological constant

- Simplest model has DE equation of state: $p = w \rho$ with constant w ($w=-1 \rightarrow$ cosmological constant)
- Little constraint from Planck alone
- SNe + Planck: tight constraint

$$w = -0.982^{+0.022}_{-0.038}$$


- BAO helps a bit, especially when considering non-flat models



Aubourg et al. (2015)

Dark energy that is not a cosmological constant

- Second simplest model has DE equation of state: $p = w \rho$ with $w = w_0 + w_a (1-a)$
- Little constraint from Planck alone




Planck 2018 VI (2020)

Dark energy that is not a cosmological constant

- Second simplest model has DE equation of state: $p = w \rho$ with $w = w_0 + w_a(1-a)$
- Little constraint from Planck alone
- SNe + Planck: tight constraint

$$w_0 = -0.851^{+0.092}_{-0.099} \quad , \quad w_a = -0.70^{+0.49}_{-0.51}$$

- BAO helps a bit, especially when considering non-flat models





Brout et al. (2022: Pantheon+)

Dark energy that is not a cosmological constant

- Second simplest model has DE equation of state: $p = w \rho$ with $w = w_0 + w_a(1-a)$
- Little constraint from Planck alone
- SNe + Planck: tight constraint

$$w_0 = -0.851^{+0.092}_{-0.099} \quad , \quad w_a = -0.70^{+0.49}_{-0.51}$$

- BAO helps a bit, especially when considering non-flat models



Aubourg et al. (2015)

Other extensions

Spoiler: it's LCDM everywhere!

Parameter	TT+lowE	TT, TE, EE+lowE	TT, TE, EE+lowE+lensing	TT, TE, EE+lowE+lensing+BAO
Ω_K	$-0.056^{+0.044}_{-0.050}$	$-0.044^{+0.033}_{-0.034}$	$-0.011^{+0.013}_{-0.012}$	$0.0007^{+0.0037}_{-0.0037}$
Σm_ν [eV]	< 0.537	< 0.257	< 0.241	< 0.120
N_{eff}	$3.00^{+0.57}_{-0.53}$	$2.92^{+0.36}_{-0.37}$	$2.89^{+0.36}_{-0.38}$	$2.99^{+0.34}_{-0.33}$
Y_P	$0.246^{+0.039}_{-0.041}$	$0.240^{+0.024}_{-0.025}$	$0.239^{+0.024}_{-0.025}$	$0.242^{+0.023}_{-0.024}$
$dn_s/d\ln k$	$-0.004^{+0.015}_{-0.015}$	$-0.006^{+0.013}_{-0.013}$	$-0.005^{+0.013}_{-0.013}$	$-0.004^{+0.013}_{-0.013}$
$r_{0.002}$	< 0.102	< 0.107	< 0.101	< 0.106
w_0	$-1.56^{+0.60}_{-0.48}$	$-1.58^{+0.52}_{-0.41}$	$-1.57^{+0.50}_{-0.40}$	$-1.04^{+0.10}_{-0.10}$

Growth of structure constraints

Growth of structure constraints

- Constraints shown so far largely only use geometric constraints on the expansion history (+ CMB)
- Growth-of-structure constraints can add valuable additional info, generally on combination of Ω_m and σ_8 :
 - RSD: $f \times \sigma_8 \sim (\Omega_m)^{0.55} \times \sigma_8$ as well as $b \times \sigma_8$
 - Lensing: $(\Omega_m)^{0.5} \times \sigma_8$, possibly b and σ_8 separately
 - Abundance of galaxy clusters: $(\Omega_m)^a \times \sigma_8$ with $a \sim 0.2$, exact form set by how cluster masses are obtained (X-ray, SZ, ...)

Weak lensing

Extremely brief overview of cosmological weak lensing

Cosmic shear

- Galaxies have a range of intrinsic shapes projected randomly on the sky
- Averaged over enough galaxies, average shape should be constant across the sky
- Lensing of background galaxies by the foreground matter distribution systematically distorts the shape —> cosmic shear ($\sim 1\%$ effect)
- Lensing is determined by the gravitational potential

$$\delta(\mathbf{k}, a) = -\frac{2}{3} \frac{k^2 a}{\Omega_{0,m} H_0^2} \Phi(\mathbf{k}, a),$$

Extremely brief overview of cosmological weak lensing


Cosmic shear

- Lensing is determined by the gravitational potential


$$\delta(\mathbf{k}, a) = -\frac{2}{3} \frac{k^2 a}{\Omega_{0,m} H_0^2} \Phi(\mathbf{k}, a),$$

- Like galaxies, we can auto- or cross-correlate shears:
 - Auto-correlation gives measurement of the power spectrum of the potential $\sim (\Omega_m)^2 P_{\delta\delta}$ —> measure $\Omega_m \times \sigma_8$
 - Cross-correlation with galaxies (galaxy-galaxy lensing): measure $b\Omega_m(\sigma_8)^2$
 - Combination with galaxy clustering ($\sim b^2 \times [\sigma_8]^2$) —> measure $\Omega_m \times \sigma_8$
- In practice, non-linear effects change these scalings, often constrained combination

$$S_8 = \sigma_8 \sqrt{\Omega_m / 0.3}$$




Heymans et al. (2020; KiDS)



h Heymans et al. (2020; KiDS)

S_8 tension


- Lensing gives lower value of S_8 than Planck



Heymans et al. (2020; KiDS)

S_8 tension

- Lensing gives lower value of S_8 than Planck



Planck 2018 VI (2020)

Conclusion

- Early+late Universe constraints complementary and highly consistent with simple flat LCDM model (BBN also consistent)
- Late-Universe measurements crucial for obtaining constraints on dark energy and its equation of state
- Some tensions:
 - H_0 tension: H_0 derived from Planck + BAO + uncalibrated SNe 5σ off from locally measured value —> see next lecture
 - S_8 tension
 - Sometimes a data point here or there is a bit off, but nothing that screams an obvious new model