

Ab initio methods in solid state physics Density-functional perturbation theory (DFPT)

Przemek Piekarz and Paweł T. Jochym

Computational Materials Science
Institute of Nuclear Physics
Polish Academy of Sciences
Kraków

May 13, 2021

Linear response

S. Baroni, S. de Gironcoli, A. D. Corso, and P. Giannozzi,
Rev. Mod. Phys. **73**, 515 (2001)

We assume that the external potential acting on the electrons is a differentiable function of a set of parameters, $\lambda = \{\lambda_i\}$ ($\lambda_i = \mathbf{R}_I$ in the case of atomic displacements), and the total energy is given by

$$E = T + V_{\text{el}} + \int d\mathbf{r} V_{\lambda}(\mathbf{r}) n(\mathbf{r}). \quad (1)$$

According to the Hellmann-Feynman theorem, the first and second derivatives of the ground-state energy read

$$\frac{\partial E}{\partial \lambda_i} = \int d\mathbf{r} \frac{\partial V_{\lambda}(\mathbf{r})}{\partial \lambda_i} n_{\lambda}(\mathbf{r}), \quad (2)$$

$$\frac{\partial^2 E}{\partial \lambda_i \partial \lambda_j} = \int d\mathbf{r} \frac{\partial^2 V_{\lambda}(\mathbf{r})}{\partial \lambda_i \partial \lambda_j} n_{\lambda}(\mathbf{r}) + \int d\mathbf{r} \frac{\partial V_{\lambda}(\mathbf{r})}{\partial \lambda_i} \frac{\partial n_{\lambda}(\mathbf{r})}{\partial \lambda_j}. \quad (3)$$

Linear response

The variation of the Kohn-Sham orbitals $\Delta\psi_m(\mathbf{r})$ is obtained by standard first-order perturbation theory

$$(H - \varepsilon_m)|\Delta\psi_m(\mathbf{r})\rangle = -(\Delta V(\mathbf{r}) - \Delta\varepsilon_m)|\psi_m(\mathbf{r})\rangle, \quad (4)$$

where the first-order correction to the self-consistent potential is

$$\Delta V(\mathbf{r}) = \Delta V_{ion}(\mathbf{r}) + \int d\mathbf{r}' \frac{\Delta n(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} + \frac{dv_{xc}}{dn} \Delta n(\mathbf{r}), \quad (5)$$

the linear change of electron density $n(\mathbf{r}) = \sum_m |\psi_m(\mathbf{r})|^2$ is given by

$$\Delta n(\mathbf{r}) = 4 \sum_m \psi_m^*(\mathbf{r}) \Delta\psi_m(\mathbf{r}) = 4 \sum_m \psi_m^*(\mathbf{r}) \sum_i \frac{\partial \psi_m(\mathbf{r})}{\partial \lambda_i} \Delta\lambda_i, \quad (6)$$

and the first-order variation of the Kohn-Sham eigenvalue ε_m

$$\Delta\varepsilon_m = \langle \psi_m | \Delta V | \psi_m \rangle. \quad (7)$$

Linear response – reciprocal space

One of the greatest advantages of DFPT is that the responses to perturbations of different wavelengths (wave vectors \mathbf{k}) are decoupled.

For the Bloch functions

$$\psi_{\mathbf{k}}(\mathbf{r}) = e^{i\mathbf{k}\mathbf{r}} u_{\mathbf{k}}(\mathbf{r}), \quad (8)$$

the Fourier component of the charge-density response at wave vector \mathbf{q} is obtained from

$$\Delta n_{\mathbf{q}}(\mathbf{r}) = \sum_{\mathbf{k}} u_{\mathbf{k}}^*(\mathbf{r}) \Delta u_{\mathbf{k}+\mathbf{q}}(\mathbf{r}). \quad (9)$$

The Fourier component of the self-consistent potential response reads

$$\Delta v_{\mathbf{q}}(\mathbf{r}) = \Delta v_{\mathbf{q}}^{ion}(\mathbf{r}) + \int d\mathbf{r}' \frac{\Delta n_{\mathbf{q}}(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} + \frac{dv_{xc}}{dn} \Delta n_{\mathbf{q}}(\mathbf{r}). \quad (10)$$

Linear response – phonon calculations

The position of the l th atom is given by

$$\mathbf{R}_l = \mathbf{R}_n + \boldsymbol{\tau}_i + \mathbf{u}_i(n) = \mathbf{R}_n + \boldsymbol{\tau}_i + \mathbf{u}_i(\mathbf{q})e^{i\mathbf{q}\mathbf{R}_n}, \quad (11)$$

where \mathbf{R}_n is the position of the n th unit cell in the Bravais lattice, $\boldsymbol{\tau}_i$ is the equilibrium position of the atom in the unit cell, and $\mathbf{u}_i(n)$ indicates the deviation from equilibrium of the nuclear position. The force constants matrix depends on the distance between two unit cells, $\mathbf{R} = \mathbf{R}_n - \mathbf{R}_m$

$$\Phi_{ij}^{\alpha\beta}(\mathbf{R}_n - \mathbf{R}_m) = \frac{\partial^2 V}{\partial u_i^\alpha(n) \partial u_j^\beta(m)}. \quad (12)$$

The Fourier transform of the force constants matrix gives the dynamical matrix

$$\sqrt{m_i m_j} D_{ij}^{\alpha\beta}(\mathbf{q}) = \sum_{\mathbf{R}} \Phi_{st}^{\alpha\beta}(\mathbf{R}) e^{-i\mathbf{q}\mathbf{R}} = \frac{1}{N} \frac{\partial^2 V}{\partial u_i^\alpha(\mathbf{q}) \partial u_j^\beta(\mathbf{q})}. \quad (13)$$

Dynamical matrix

Dynamical matrix consists of the electronic and ionic part

$$D_{ij}^{\alpha\beta}(\mathbf{q}) = D_{ij}^{\alpha\beta}(\mathbf{q})_{\text{el}} + D_{ij}^{\alpha\beta}(\mathbf{q})_{\text{ion}} \quad (14)$$

The electronic part includes changes in charge density

$$D_{ij}^{\alpha\beta}(\mathbf{q})_{\text{el}} = \frac{1}{N\sqrt{m_i m_j}} \int d\mathbf{r} \left[\frac{\partial n(\mathbf{r})}{\partial u_i^\alpha(\mathbf{q})} \frac{\partial V_{\text{ion}}(\mathbf{r})}{\partial u_j^\beta(\mathbf{q})} + n(\mathbf{r}) \frac{\partial^2 V_{\text{ion}}(\mathbf{r})}{\partial u_i^\alpha(\mathbf{q}) \partial u_j^\beta(\mathbf{q})} \right]. \quad (15)$$

The phonon frequencies $\omega(\mathbf{q})$ and polarization vectors $e_{i\alpha}(\mathbf{q})$ are obtained from the eigenequation

$$\sum_{j\beta} D_{ij}^{\alpha\beta}(\mathbf{q}) e_{j\beta}(\mathbf{q}) = \omega^2(\mathbf{q}) e_{i\alpha}(\mathbf{q}). \quad (16)$$

Phonons in polar materials

In the long-wavelength limit ($\mathbf{q} \rightarrow 0$), the dynamical matrix can be split into the sum of an analytic and a nonanalytic contribution (Born and Huang, 1954)

$$D_{ij}^{\alpha\beta} = D_{ij,\text{an}}^{\alpha\beta} + D_{ij,\text{na}}^{\alpha\beta}, \quad (17)$$

where analytic part is the matrix obtained from the response to a zone-center phonon ($\mathbf{q} = 0$), calculated at zero macroscopic electric field. The nonanalytic part describes the longitudinal optic/transverse optic (LO/TO) splitting of the infrared modes, and has the general form

$$D_{ij,\text{na}}^{\alpha\beta} = \frac{4\pi e^2}{\Omega} \frac{\sum_{\gamma} Z_i^{*\gamma\alpha} q_{\gamma} \sum_{\nu} Z_j^{*\nu\beta} q_{\nu}}{\sum_{\gamma\nu} q_{\gamma} \epsilon_{\infty}^{\gamma\nu} q_{\nu}}, \quad (18)$$

where $\epsilon_{\infty}^{\gamma\nu}$ is the electronic dielectric-constant tensor and $Z_i^{*\gamma\alpha}$ is the Born effective charge tensor

$$Z_i^{*\gamma\alpha} = \frac{\Omega}{e} \frac{\partial P_{\alpha}}{\partial u_i^{\beta}(\mathbf{q} = 0)}. \quad (19)$$

Electron-phonon interaction

F. Giustino, Rev. Mod. Phys. **89**, 015003 (2017)

Electron-phonon interaction is important for transport properties – electric and heat conductivity. Interaction of phonons with electrons leads to reduction of phonon lifetime $\tau_{i\mathbf{q}}$

$$\frac{1}{\tau_{i\mathbf{q}}} = \frac{4\pi}{\hbar} \sum_{mn} \int \frac{d\mathbf{k}}{\Omega_{\text{BZ}}} |g_{mni}(\mathbf{k}, \mathbf{q})|^2 (f_{n\mathbf{k}} - f_{m\mathbf{k}+\mathbf{q}}) \delta(\varepsilon_{n\mathbf{k}} - \varepsilon_{m\mathbf{k}+\mathbf{q}} - \hbar\omega_{i\mathbf{q}}). \quad (20)$$

Electron-phonon matrix elements are given by

$$g_{mni}(\mathbf{k}, \mathbf{q}) = \langle u_{m\mathbf{k}+\mathbf{q}} | \Delta_{i\mathbf{q}} V | u_{n\mathbf{k}} \rangle = \int d\mathbf{r} u_{m\mathbf{k}+\mathbf{q}}^* \Delta_{i\mathbf{q}} V u_{n\mathbf{k}}, \quad (21)$$


where

$$\Delta_{i\mathbf{q}} V = \sum_{j\alpha} e_{j\alpha,i}(\mathbf{q}) \sum_{\mathbf{R}} e^{-i\mathbf{q}(\mathbf{r}-\mathbf{R})} \frac{\partial V}{\partial u_j^\alpha}. \quad (22)$$

Phonon mediated superconductivity

In the Bardeen-Cooper-Schrieffer (BCS) theory, the superconducting current is carried by the Cooper pairs, created by to attraction between electrons due to electron-phonon coupling. The strength of electron-phonon interaction is described by the coupling constant

$$\lambda = 2 \int d\omega \frac{\alpha^2 F(\omega)}{\omega}. \quad (23)$$



$\alpha^2 F(\omega)$ is the Eliashberg function

$$\alpha^2 F(\omega) = \frac{1}{N_F} \int \frac{d\mathbf{k} d\mathbf{q}}{\Omega_{\text{BZ}}^2} \sum_{mni} |g_{mni}(\mathbf{k}, \mathbf{q})|^2 \delta(\varepsilon_{n\mathbf{k}} - \varepsilon_F) \delta(\varepsilon_{m\mathbf{k}+\mathbf{q}} - \varepsilon_F) \delta(\omega - \omega_{i\mathbf{q}}). \quad (24)$$

Figure made by Andrzej Ptok

Phonon mediated superconductivity

The superconducting critical temperature can be estimated using on a semiempirical expression first introduced by McMillan and then refined by Allen and Dynes

$$T_c = \frac{\hbar\omega_{\log}}{1.2} \exp \left[- \frac{1.04(1 + \lambda)}{\lambda - \mu^*(1 + 0.62\lambda)} \right], \quad (25)$$

where ω_{\log} is obtained using the Eliashberg function

$$\omega_{\log} = \exp \left[\frac{2}{\lambda} \int d\omega \frac{\alpha^2 F(\omega)}{\omega} \log \omega \right]. \quad (26)$$


The parameter μ^* is obtained as

$$\frac{1}{\mu^*} = \frac{1}{\mu} - \log \frac{\omega_p}{\omega_{ph}}, \quad (27)$$

where μ is the average electron-electron Coulomb repulsion, ω_p is the characteristic plasma frequency, and ω_{ph} is the largest phonon frequency.

Superconductivity of LaH_{10}

A. P. Drozdov *et al.*, Nature **569**, 528 (2019)



Superconductivity of LaH_{10}

I. A. Kruglov *et al.*, Phys. Rev. B **101**, 024508 (2020)

The calculations of the critical temperature and electron-phonon coupling (EPC) parameters were carried out using the QUANTUM ESPRESSO package within density functional perturbation theory, employing the plane-wave pseudopotential method and PBE exchange-correlation functional.

