.

Description of the Agent-Based Model

The simulations for this work were performed by our agent-based simulation framework **QHG** ("Quantitative Humane Geophylogeny").

Environment

The agents in **QHG**, live and move on the nodes of a grid (for the simulations described here, we used a spherical grid based onthe subdivision of an icosahedron). Each node of the grid is assigned a set of environmental data pertainiing to its coordinates: geography (e.g. altitude, sealevels (REF:SEALEVEL), ice (REF:ICE), rivers), climate (temperature, precipitation), and vegetation (NPP (REF:NPP)). The carrying capacity of the cell is calulated from its NPP by means of a ramp function, given by the parameters $K_{min}, K_{max}, NPP_{min}, NPP_{max}$ as follows: If $NPP_{cur} < NPP_{min}$ then $K_{cur} = K_{min}$, if $NPP_{cur} > NPP_{max}$ then $K_{cur} = K_{max}$. Otherwise,

$$K_{cur} = K_{min} + (K_{max} - K_{min})(NPP_{cur} - NPP_{min})/(NPP_{max} - NPP_{min})$$

The environment data is updated every 1'000 steps (years) to reflect changes in climate, ice and sea-level

Seafaring, e.g. island-hopping to Australia, is realized by adding low-probability connections between two nodes separated by water

For a full description of the environmental variables cf. table 1.

Agents

At every time step an agent can perform different actions such as move, mate and bear offspring, and/or die. Movement is strongly influenced by the environment: the direction in which to move is a random choice weighted by altitude and carrying capacity of the neighboring nodes, and agents can not move to nodes under water or covered with ice.

Birth and death probabilities are calculated with linear functions leading to logistic growth:

$$p_{birth} = b_0 + (\theta - b_0)\frac{N}{K}, \qquad p_{death} = d_0 + (d_0 - \theta)\frac{N}{K},$$

where N is the number of agents, K is the local carrying capacity and θ is the turnover.

A description of the agents' configurable attributes can be found in table 2.

Table 1: Environmental Variables

Geography	
`Latitude`	latitude of node (radians)
`Longitude`	longitude of node (radians)
`Altitude`	altitude of node (meters)
`Ice`	ice (1) or no ice (0)
`Water`	presence of water (e.g. rivers)
`Coastal`	close to coast (1) or not (0)
`Distances`	distances to neighboring nodes
`Area`	area of region associated wit cell
Climate	
`AnnualMeanTemp`	current annual mean temperature in node
`AnnualRainfall`	current annual total rainfall in node
Vegetation	
`NPP`	Net Primary Production in node

Table 2: Agent Attributes

Birth and Death Probabilies	
`Verhulst_b0`	birth probability for $N = 0$
`Verhulst_d0`	death probability for $N = 0$
`Verhulst_theta`	turn-over rate
Fertility	
`Fertility_interbirth`	interbirth interval (years)
`Fertility_max_age`	maximum fertile age (years)
`Fertility_min_age`	minimum fertile age (years)
Old Age Cut-Off	
`OAD_max_age`	average age cut-off (years)
`OAD_uncertainty`	variance
NPP-Capacity Conversion	
`NPPCap_K_max`	K_{max} parameter for ramp function
`NPPCap_K_min`	K_{min} parameter for ramp function
`NPPCap_NPP_max`	NPP_{max} parameter for ramp function
`NPPCap_NPP_min`	NPP_{min} parameter for ramp function
`NPPCap_water_factor`	Capacity increase for presence of water
`NPPCap_coastal_factor`	Capacity increase for vicinity of coast
Move Probability	
`WeightedMove_prob`	move probability
Navigation	
`Navigate_dist0`	reference distance for crossing
`Navigate_prob0`	probability of crossing at reference distance
`Navigate_decay`	factor for exponential decay of probability with distance