

Neutron Update

Jyotirmai Singh

Neutron-Nucleus Cross Section

- Contributions from nuclear, electromagnetic (electrostatic, magnetic moment, polarisation), new interactions
- Collect into coherent scattering length b_c , electromagnetic contribution χ_{em} and new contribution χ_{new} .
- $\chi_{em} = Z(b_F + b_l)/b_c$, Foldy scattering length and intrinsic neutron-electron scattering length.
- $f(q)$ is the form factor of the atom, modeled to 10^{-4} accuracy by $f \approx [1 + 3(q/q_0)^2]^{-0.5}$, $q_0 = 6.86 \text{ \AA}^{-1}$
- b_s Schwinger scat. length, b_i incoherent scat. length, both assumed negligible.

$$V_{new}(r) = -\frac{1}{4\pi} g^2 Q_1 Q_2 \frac{e^{-\mu r}}{r},$$

$$\frac{d\sigma}{d\Omega}(q) = b_c^2 \left\{ 1 + \chi_{em}[1 - f(q)] + \chi_{new} \frac{\mu^2}{q^2 + \mu^2} \right\}^2 + b_s^2(q) + b_i^2 + O(b_F^2)$$


$$\simeq b_c^2 \left\{ 1 + 2\chi_{em}[1 - f(q)] + 2\chi_{new} \frac{\mu^2}{q^2 + \mu^2} \right\},$$

Table 1


Typical values of the various contributions δb to the total neutron scattering length b of a heavy atom

Class	Interaction	δb (fm)
I	Strong interaction	10.0
	Atomic magnetic dipole moment*	10.0
II	Spin-orbit (Schwinger)	0.1
	Foldy	0.1
	Neutron electric polarizability	0.05
	Intrinsic electrostatic	0.01
	Nuclear magnetic dipole moment*	0.005
III	Neutron electric dipole moment*	$\leq 10^{-8}$
	Neutron electric charge*	$\leq 10^{-10}$
	Weak interaction	$\sim 10^{-34}$


* If any.




Cross Section with and without new force
 $\lambda = 1 \text{ \AA}, a = 100 \text{ \AA}$




Cross Section with and without new force zoom in



$$\lambda = 1 \text{ \AA}$$



$$\lambda = 10 \text{ \AA}$$



$$\lambda = 0.1 \text{ \AA}$$

Lower energy
neutrons seem
easier to use

Scattering off Lattice

- Key feature of lattice - only allows certain momentum transfers.
- For a crystal, output intensity is given by $I(\theta) \sim F_{hkl}^2$, h,k,l Miller Indices related to lattice FT and indicates where diffractive peaks occur.
- Use Bragg Law to convert h,k,l into an angular position. For a cubic lattice:


Bragg's Law

$$\lambda = 2d \sin(\theta)$$

For a cubic system


$$\frac{1}{d^2} = \frac{h^2 + k^2 + l^2}{a^2}$$

$$d_{hkl} = \frac{a}{\sqrt{h^2 + k^2 + l^2}} \implies \lambda = \frac{2a \sin(\theta)}{\sqrt{h^2 + k^2 + l^2}} \implies \sin^2(\theta) = \frac{\lambda^2}{4a^2} (h^2 + k^2 + l^2)$$




$$F_{hkl} = \sum_{j=1}^N f_j e^{[-2\pi i(hx_j + ky_j + lz_j)]},$$

Sum over all atoms of unit cell. I think f_j is the single atom neutron cross section for the j^{th} atom.

Lattice Scattering Intensity $\mu = 2.0\text{e}+02$ 

Neutrons off Cubic Xenon Lattice

$$a = 10 \text{ \AA}, \mu = 200 \text{ eV}, \lambda = 1 \text{ \AA}$$


Lattice Scattering Intensity $\mu = 2.0\text{e}+02$ 

Neutrons off Cubic Xenon Lattice


$$a = 100 \text{ \AA}, \mu = 200 \text{ eV}, \lambda = 1 \text{ \AA}$$

5 Miller Planes


All plots with
 $a = 100 \text{ \AA}$, $\mu = 200 \text{ eV}$, $\lambda = 1 \text{ \AA}$




4 Miller Planes



3 Miller Planes




Question:
 How many of
 these peaks
 do we care
 about?




$$\lambda = 1 \text{ \AA}, a = 100 \text{ \AA}$$

$$\lambda/a = 0.01$$



$$\lambda = 10 \text{ \AA}, a = 100 \text{ \AA}$$

$$\lambda/a = 0.1$$



$$\lambda = 0.1 \text{ \AA}, a = 100 \text{ \AA}$$

$$\lambda/a = 0.001$$

**For lattice λ/a is important.
Lower λ/a gives sharper differentiation in a narrower angle range.**

Summary/Next Steps

- Neutron cross section dominated by strong force and electromagnetic effects, but can see new force via scattering.
- For simple atomic scattering higher λ gives better signal.
- For (primitive cubic) lattice scattering, important parameter is λ/a . Lower λ/a gives sharper contrast in smaller angular range.
- Next step is to introduce irregular lattice, i.e. unit cell axes with different lengths a, b, c and then put different atoms in the unit cell.