

Literature Review

Jomar Alcantara

2018-04-26

Contents

1	Introduction to Dementia and the Literature Review	3
2	Description of Dementia	3
2.1	Alzheimer's Disease	3
2.2	Vascular Dementia	3
2.3	Dementia with Lewy Bodies	3
2.4	Parkinson's disease dementia	3
2.5	Fronto-temporal dementia	3
2.6	Primary Progressive Aphasia	3
2.7	Mild Cognitive Impairment	3
3	Assessment of Language function in Dementia	3
4	Current ways of diagnosing Dementia using Language	4
5	Existing studies that have used Natural Language Processing to help in diagnosing psychological difficulties, particularly language difficulties	5
5.1	Nun Study - Snowden (1996)	5
5.2	Comparing Presidents Speeches - Berisha et al (2015)	5
6	State of literature into Natural Language processing techniques	5
7	Conclusion	5

1 Introduction to Dementia and the Literature Review

According to a recent report commissioned by the Alzheimer's Society, they estimate the prevalence of Dementia in the UK at approximately 815,000 people. This represents 1 in 14 of the population aged 65 or over. This report also estimates an annual healthcare spend on 4.3 billion of which approximately 85 million is spend solely on diagnosis. They also estimate that the overall impact of dementia (excluding the costs associated with early onset dementia) is 26.3 billion annually. Globally, this picture is a lot bleaker. A recent report suggests that in 2015 there were 46 million people with a diagnosis of dementia and that number is expected to hit 131.5 million by 2050(CITE). The report also states that the worldwide cost of dementia in 2018 is estimated to be in the region of one trillion US dollars.

The purpose of this review is to seek to understand how dementia impacts speech and language production from a psychological and linguistic perspective. With a view to potentially using natural language processing and technology to aid in the early diagnosis of dementia and/or Mild Cognitive Impairment.

A search of the literature was conducted mainly from a psychological perspective as this would inform the theoretical side of things. A search of the literature from a Computer Science perspective was also made with a view to exploring the how perspectives of machine learning, more specifically natural language processing can be applied to

2 Description of Dementia

2.1 Alzheimer's Disease

2.2 Vascular Dementia

2.3 Dementia with Lewy Bodies

2.4 Parkinson's disease dementia

2.5 Fronto-temporal dementia

2.6 Primary Progressive Aphasia

2.7 Mild Cognitive Impairment

3 Assessment of Language function in Dementia

Language can be defined as the ability to encode ideas into words and/or symbols for the means of communication. Difficulty in language, both spoken and written, are often described as symptoms of various types of Alzheimers' disease (AD). One of the challenges when using speech as a predictor is recognising whether there is a problem with language production or with the motor skills. Impairments in speech that arise from any process that disrupts the neuraxis

from the cortex to muscle and encompass dysarthria (disturbance in articulation) and dysphonia (disturbance in the production in vocal sounds). However it is important to use language as a tool to aid the diagnosis of Dementia, as it provides vital clues to aid a clinician in differentiating in the different types of Dementia, which will in turn aid a clinician in an attempt to manage an individual case of Dementia. Due to the subtle nature of the languages changes that are experienced in those who suffer with different types of Dementia, it is often misdiagnosed.

A common way for clinicians' to diagnose dementia is through use of the Mini-Mental State Examination. The Mini-Mental State Examination (MMSE) has shown to be particularly effective at differentiating between Dementia and other psychological disorders such as clinical depression, schizophrenia and personality disorder and is currently the most widely used measure of diagnosing Dementia within clinical psychology. However, it is not without it's faults...

4 Current ways of diagnosing Dementia using Language

Battery of tests to differentiate between Controls, those with MCI, those with AD.

- FCSRDT - As most effective
- Mini Mental State Examination (MMSE)
- Addenbrooke's Cognitive Examination (ACE-III)
- Dementia Rating Scale (DRS)
- Western Aphasia Battery (WAB)
- Boston Diagnostic Aphasia Examination (BDAE)
- Boston Naming Test (BNT)

5 Existing studies that have used Natural Language Processing to help in diagnosing psychological difficulties, particularly language difficulties

5.1 Nun Study - Snowden (1996)

5.2 Comparing Presidents Speeches - Berisha et al (2015)

6 State of literature into Natural Language processing techniques

7 Conclusion

Identification of language impairment is important in Dementia because it aids diagnosis of specific types of dementia, which in turn can alter the prognosis and change the management of the degenerative disorder. As these differences in language are quite subtle, the varying subtypes of dementia are frequently misdiagnosed.

Given the burden on the diagnosis of dementia on clinicians, it appears to be useful to find some non-invasive protocols for the early diagnosis of dementia. It has already been shown that analysis of speech and language has shown markers that pre-date the official diagnosis of dementia (Snowdon et al, 1996; Berisha et al, 2015).