

# Circuit Specific Modulation by Corticotrophin Releasing Factor in Ventral Tegmental Area Neurons


Driscoll, J. R.<sup>1,3</sup> Fields, H. L.<sup>1,2,3</sup> Margolis, E. B.<sup>1,2</sup>

1. Dept. of Neurology, 2. Wheeler Center for the Neurobiology of Addiction, University of California San Francisco, San Francisco, CA 94143

3. Helen Wills Neuroscience Institute, University of California Berkeley, Berkeley CA, 94709


## Introduction

Several rodent models of stress (e.g. footshock, foot pinch, restraint stress, and social defeat stress) increase the activity of VTA neurons<sup>1,8,13-14</sup> and result in an increase in dopamine in the terminal regions of VTA neurons<sup>2,7,9,11,12,17</sup>. Corticotrophin releasing factor (CRF) is released in the VTA during stress and is reported to increase the firing rate of dopamine neurons<sup>1,13-14</sup>, yet decrease dopamine release in the nucleus accumbens<sup>15</sup>. These results suggest that CRF differentially modulates VTA neurons with different projection targets. However, the synaptic action of CRF on specific subsets of VTA neurons and their role in influencing behavior has yet to be characterized.




## Methods

We used whole cell patch clamp recordings in VTA neurons *ex vivo* to examine the synaptic actions of CRF. Current clamp was used to assess changes in membrane potential or firing rate. Voltage clamp was used to determine effect on glutamate EPSCs.



To analyze circuit specific responses of CRF the retrograde fluorescent marker Dil was injected into midbrain terminal regions including nucleus accumbens (NAc), medial prefrontal cortex (mPFC), and amygdala (AMY). A schematic shows Dil injection sites in NAc, mPFC, and AMY, with VTA projection lines.




### Acknowledgements:

We thank Catriona Miller, Jocelyn Breton, Allison Coker, and Peter Fong for technical assistance.

This work was supported by funding from the State of California for medical research on alcohol and substance abuse through the University of California, San Francisco, National Science Foundation Graduate Research Fellowship under grant no. DGE 1106400, and NIDA grant R01DA016782.

## Electrophysiology


### I. CRF increases the firing rate in neurons that are spontaneously active




Neurons were filled with biocytin during the recording and later tested for tyrosine hydroxylase (TH) content with immunocytochemistry.

Both TH positive and TH negative spontaneously active neurons showed a consistent increase in firing rate to bath application of CRF.

### II. Neurons that are quiescent at baseline respond to CRF with either depolarizations or hyperpolarizations.




### III. Are CRF's excitatory actions mediated by activation of hyperpolarization-activated cyclic nucleotide-gated (HCN) channels?




## CRF has mixed effects on Glutamatergic EPSCs

Using microdialysis, Wang et al. 2005 showed that CRF increases both glutamate and dopamine concentrations in the VTA; further, these effects are blocked with a local glutamate antagonist<sup>16</sup>. To explore CRF's modulation of glutamatergic inputs, we investigated CRF's ability to modulate glutamatergic excitatory postsynaptic currents.




## Do CRF Effects Differ by neuronal projection target?


### I. NAc-Projecting Neurons



### II. mPFC-Projecting Neurons



### III. Amygdala-Projecting Neurons



## Summary

### Synaptic actions of CRF in VTA neurons:

Are excitatory in neurons that are already spontaneously active, regardless of projection target

Are heterogeneous in quiescent neurons causing both excitations and inhibitions in both dopaminergic and non-dopaminergic neurons projecting to NAc and mPFC

Universally excite neurons that project to the amygdala

Future work is needed to understand how modulation of these different projections contribute to CRF-induced behavioral effects

### References:

- Anstrom, K.K., Woodward, D.J., 2005. Restraint Increases Dopaminergic Burst Firing in Awake Rats. *Neuropharmacology* 49, 1832–1840. doi:10.1038/sj.nph.1307230
- Deutch, A.Y., Lee, M.C., Gilman, M.H., Cameron, D.A., Goldstein, M., Iadarola, M.J., 1991. Stress Selectively Increases Fos Protein in Dopamine Neurons Innervating the Prefrontal Cortex. *Cereb. Cortex* 1, 273–292. doi:10.1093/cercor/1.2.273
- Deutch, A.Y., Lee, M.C., Gilman, M.H., Cameron, D.A., Goldstein, M., Iadarola, M.J., 1991. Stress selectively increases fos protein in dopamine neurons innervating the prefrontal cortex. *Cereb. Cortex* 1, 273–292.
- Inglis, F.M., Moghaddam, B., 1999. Dopamineergic innervation of the amygdala is highly responsive to stress. *Journal of Neuroscience* 19, 1088–1094.
- Jedema, P.F., Deutch, A.J., 1994. Glutamatergic control of dopamine release during stress in the rat prefrontal cortex. *Journal of Neuroscience* 14, 765–788.
- Kalivas, P.W., Abbott, R., 1987. Endocrinological regulation of monoaminergic terminals in the shell of the nucleus accumbens by stress. *Brain Res.* 414, 339–348. doi:10.1016/0006-8950(87)90015-1
- Kalivas, P.W., Duffy, P., 1995. Selective activation of dopamine transmission in the shell of the nucleus accumbens by stress. *European Journal of Neuroscience* 12, 2677–2685. doi:10.1111/j.1469-6568.2008.04792.x
- Lavicky, J., Dunn, A.J., 1998. Corticotrophin-Releasing Factor Stimulates Catecholamine Release in Hypothalamus and Prefrontal Cortex in Freely Moving Rats as Measured by Microdialysis. *Endocrinology* 139, 165–171. doi:10.1210/en.139.1.165
- Piazza, P.V., Le Moal, M., 1998. The role of stress in drug self-administration. *Trends in Pharmacological Sciences* 19, 67–74. doi:10.1016/S1563-624X(98)80015-2
- Roth, R.H., Tam, S.-Y., Iida, Y., Yang, J.-X., Deutch, A.Y., 1988. Stress and the Mesocorticolimbic Dopamine System. *Annals of the New York Academy of Sciences* 510, 325–328. doi:10.1111/j.1749-6632.1988.tb32426.x
- Thierry, A.M., Tassin, J.P., Blanc, G., Glowinski, J., 1976. Selective activation of dopamine transmission in the shell of the nucleus accumbens by stress. *Nature* 263, 242–244. doi:10.1038/263242b0
- Ungless, M.A., Smith, D.L., Fowler, T.L., Yake, R., Ron, D., Benet, A., 2003. Corticotropin-Releasing Factor Requires CRF Binding Protein to Potentiate NMDA Receptor via CRF Receptor 2 in Dopamine Neurons. *Neuron* 39, 401–407. doi:10.1016/S0896-6273(03)00461-6
- Wantat, M.J., Hwang, F.W., Phillips, P.E., Bonci, A., 2013. CRF acts in the midbrain to attenuate dopamine release to rewards but not their predictors. *Nature* 500, 383–386. doi:10.1038/nature12500
- Wang, B., Shaham, Y., Zitman, D., Azari, S., Wise, R.A., You, Z.-B., 2005. Cocaine Experience Establishes Control of Midbrain Glutamate and Dopamine by Conditioned Stress. *The Journal of Physiology* 586, 2157–2170. doi:10.1113/jphysiol.2007.150978
- Wise, R.A., Morales, M., 2010. A ventral tegmental CRF-glutamate-dopamine interaction in addiction. *Brain Research* 1314, 38–43. doi:10.1016/j.brainres.2009.09.101

