

Circuit Specific Modulation of Corticotrophin Releasing Factor in Ventral Tegmental Neurons


Driscoll, J. R.^{1,3} Fields, H. L.^{1,2,3} Margolis, E. B.^{1,2}

1. Dept. of Neurology, 2. Wheeler Center for the Neurobiology of Addiction, University of California San Francisco, San Francisco, CA 94143

3. Helen Wills Neuroscience Institute, University of California Berkeley, Berkeley CA, 94709


Introduction

Several rodent models of stress (e.g. footshock, foot pinch, restraint stress, and social defeat stress) increase the activity of VTA neurons and result in an increase in dopamine in the terminal regions of VTA neurons. Corticotrophin releasing factor (CRF) is released in the VTA during stress and is reported to increase the firing rate of dopamine neurons, yet decrease dopamine release in the Nucleus Accumbens. These results suggest that CRF differentially modulates VTA neurons with different projection targets. However, the synaptic action of CRF on specific subsets of VTA neurons and their role in influencing behavior has yet to be characterized.




Methods

We used whole cell patch clamp recordings in VTA neurons *ex vivo* to examine the synaptic actions of CRF. Current clamp was used to assess changes in membrane potential or firing rate. Voltage clamp was used to determine effect on glutamate EPSCs




To analyze circuit specific responses of CRF the retrograde fluorescent marker Dil was injected into midbrain terminal regions including, Nucleus Accumbens (NAc), medial prefrontal cortex (mPFC), and amygdala (AMY)




Electrophysiology

I. CRF increases the firing rate in neurons that are spontaneously active




Both TH positive and TH negative neurons showed a consistent increase in firing rate to bath application of CRF in spontaneously active neurons.

II. Neurons that are quiescent at baseline respond to CRF with either depolarizations or hyperpolarizations.




III. Are CRF's excitatory actions mediated by activation of hyperpolarization-activated cyclic nucleotide-gated (HCN) channels?




CRF has mixed effects on Glutamatergic EPSCs

Using Microdialysis, Wang et al. 2005 showed that CRF increases both glutamate and dopamine concentrations in the VTA; further these effects are blocked with a local glutamate antagonist. To explore CRF's relationship with glutamatergic inputs, we investigated CRF's ability to modulate glutamatergic excitatory postsynaptic currents.




Do CRF Neuronal Effects Differ by their Projecting Target?


I. NAc Projecting Neurons



II. mPFC Projecting Neurons



III. Amygdala Projecting Neurons



AMY-Projecting neurons are consistently excited by CRF, demonstrated by an increase in firing rate, depolarization, or facilitation of evoked glutamatergic EPSCs.

Summary

Synaptic Actions of CRF in VTA neurons:

Are heterogeneous, causing both excitations and inhibitions

Targets both dopaminergic and non-dopaminergic neurons

Are excitatory in neurons that are already spontaneously active, regardless of projection target

Excites neurons that project to the Amygdala

Further work is needed to understand how CRF effects behavior based on differences in projection

References:

- Beckstead, M. J. et al. CRF Enhancement of GABA Channel-Mediated Transmission in Dopamine Neurons. 34, 1926-1935 (2009).
- Blacktop, J. M. et al. Augmented Cocaine Seeking in Response to Stress or CRF Delivered into the Ventral Tegmental Area Following Long-Access Self-Administration Mediated by CRF Receptor Type 1 But Not CRF Receptor Type 2. Journal of Neuroscience 31, 11399-11403 (2011).
- Erb, S., Shaham, Y. & Stewart, J. The role of corticotropin-releasing factor and corticosterone in stress- and cocaine-induced relapse to cocaine seeking in rats. J. Neurosci. 18, 5529-5538 (1998).
- Erb, S., Shaham, Y. & Stewart, J. Stress reinstates cocaine-seeking behavior after prolonged extinction and a drug-free period. Psychopharmacology (Berl.) 138, 408-412 (1998).
- Hahn, J., Hopt, F. W. & Bonci, A. Chronic Cocaine Enhances Corticotropin-Releasing Factor-Dependent Potentiation of Excitatory Transmission in Ventral Tegmental Area Dopamine Neurons. Journal of Neuroscience 29, 6535-6544 (2009).
- Kalivas, P. W., Duffy, P. & Latimer, L. G. Neurochemical and behavioral effects of corticotropin-releasing factor in the ventral tegmental area of the rat. Journal of Pharmacology and Experimental Therapeutics 242, 757-763 (1987).
- Lavicky, J. & Dunn, A. J. Corticotropin-releasing factor stimulates catecholamine release in hypothalamus and prefrontal cortex in freely moving rats as assessed by microdialysis. J. Neurochem. 60, 602-612 (1993).
- Piazza, P. V. & Le Moal, M. The role of stress in drug self-administration. Trends in Pharmacological Sciences 19, 67-74 (1998).
- Riegel, A. C. & Williams, J. T. CRF Facilitates Calcium Release from Intracellular Stores in Midbrain Dopamine Neurons. Neuron 57, 559-570 (2008).
- Shaham, Y., Erb, S., Leung, S., Buzsaki, Y. & Stewart, J. CRF-15A-226, a selective, non-peptide antagonist of the corticotropin-releasing factor1 receptor attenuates stress-induced relapse to drug seeking in cocaine- and heroin-trained rats. Psychopharmacology (Berl.) 137, 184-190 (1998).
- Shaham, Y. et al. Corticotropin-releasing factor, but not corticosterone, is involved in stress-induced relapse to heroin-seeking in rats. J. Neurosci. 17, 2605-2614 (1997).
- Tagliaferro, P. & Morales, M. Synapses between corticotropin-releasing factor-containing axon terminals and dopaminergic neurons in the ventral tegmental area are predominantly glutamatergic. J. Comp. Neurol. 508, 616-626 (2007).
- Ungless, M. A. et al. Corticotropin-releasing factor requires CRF binding protein to potentiate NMDA receptors via CRF receptor 2 in dopamine neurons. Neuron 39, 401-407 (2003).
- Wan, M. J., Hopt, F. W., Stuber, G. D., Phillips, P. E. M. & Bonci, A. Corticotropin-releasing factor increases mouse ventral tegmental area dopamine neuron firing through a protein kinase C-dependent enhancement of Ih. J. Physiol. (Lond.) 586, 2157-2170 (2008).
- Wan, M. J., Bonci, A. & Phillips, P. E. M. CRF acts in the midbrain to attenuate accumbens dopamine release to rewards but not their predictors. Nature Publishing Group 16, 383-386 (2015).
- Wang, B., You, Z.-B., Rice, K. C. & Wise, R. A. Stress-induced relapse to cocaine seeking: roles for the CRF2 receptor and CRF-binding protein in the ventral tegmental area of the rat. Psychopharmacology (Berl.) 193, 283-294 (2007).
- Wang, B. et al. Cocaine experience establishes control of midbrain glutamate and dopamine by corticotropin-releasing factor: a role in stress-induced relapse to drug seeking. Journal of Neuroscience 25, 5389-5396 (2005).
- Williams, C. L., Buchta, W. C. & Riegel, A. C. CRF-R2 and the Heterosynaptic Regulation of VTA Glutamate during Reinstatement of Cocaine Seeking. Journal of Neuroscience 34, 10402-10414 (2014).

Acknowledgements:

We thank Catriona Miller, Jocelyn Breton, Allison Coker, and Peter Fong for technical assistance.

This work was supported by funding from the State of California for medical research on alcohol and substance abuse through the University of California, San Francisco, National Science Foundation Graduate Research Fellowship under grant no. DGE 1106400, and NIDA grant R01DA016782.

