

# Corticotrophin Releasing Factor Alters Kappa Opioid Receptor Function in the Ventral Tegmental Area.

<sup>1</sup> Dept. of Neurology, <sup>2</sup> Wheeler Center for the Neurobiology of Addiction, University of California San Francisco, San Francisco, CA 94143  
<sup>3</sup> Helen Wills Neuroscience Institute, University of California Berkeley, Berkeley CA, 94709

## Background

Depending on the degree or type, stress can be associated with either positive (rewarding) or negative (aversive) behavioral outcomes.

During stress, corticotrophin releasing factor (CRF) is released into the ventral tegmental area (VTA), which increases cocaine (Coc) signaling in the ventral tegmental regions including the nucleus accumbens (NAc) (Wang et al., 2009). DA in the NAc is typically associated with reward.


CRF release also leads to kappa opioid receptor (KOR) activation in brain areas that are involved in stress, resulting in conditioned place aversion (Lund et al., 2008; Bruchas et al., 2009). We previously showed that the KOR agonist U69593 selectively hyperpolarizes VTA DA neurons that project to the amygdala or prefrontal cortex, but not DA neurons projecting to the NAc (Margolis et al., 2006; 2008).

Understanding the interaction of CRF and the kappa opioid system may help to dissociate the negative deleterious aspects of stress from those that allow for positive physiological adaptation.

## Methods



Horizontal rat brain slice

Whole cell ex vivo electrophysiology was used to examine how CRF modulates responses to U69593 in VTA neurons



In current clamp (0-0 pA), VTA neurons were exposed to CRF (100 nM) and then to both applications of U69593, a KOR agonist, and CRF.


TH immunocytochemistry



U69593 and CRF were applied by bath perfusion

CRF was soaked first for 5-6 min and then washed out for at least 10 min before a second U69593 application


I. Biased Sampling



Baseline KOR responses were evaluated with U69593 (1  $\mu$ M) before application of CRF to confirm KOR responses in a subset of experiments

Comparison of multiple applications of U69593 to CRF then U69593 application in confirmed KOR responding neurons


II. Unbiased Sampling




In a separate set of experiments, cells were briefly sampled from the VTA and tested with CRF followed by U69593 without prior knowledge of kappa responses

## Electrophysiology


I. Current Clamp



II. CRF effect does not predict U69593 response




III. Voltage Clamp



IV. Summary: How often are excitations observed?


Biased Sampling:

Unbiased Sampling:




## Excitations to U69593


I. Comparison of U69593 effects in confirmed KOR responding cells



Histograms of response to second application of U69593



II. Timing of excitatory responses



## Summary

1) CRF exposure robustly altered KOR induced signaling in VTA neurons, causing a switch from inhibition to excitation in a subset of VTA dopaminergic neurons.

2) Excitators did not correlate with magnitude of the baseline U69593 response or the response to CRF

3) Excitators were seen in ~45% of cells that were previously identified with KOR hyperpolarizations and ~25% of neurons that were blindly sampled from the VTA

These results provide insight into how stress may profoundly alter neuronal responses to KOR activation in mesolimbic neurons.

## Future Directions

I. Intracellular vs Synaptic Mechanism

Future studies evaluate how CRF stimulates changes within the cell to alter response properties to KOR activation


1. G-protein activation

2. Receptor phosphorylation

3. Arrestin recruitment

4. MAPK signaling

5. Recycling



Credit to: Aghajanian and Bruchas, 2011

II. Behavioral Relevance

Conditioned Place Preference to U69593 in stressed and CRF pretreated animals

## References

Aghajanian, A., & Bruchas, M. (2011). Mechanisms of opioid behavior: Opposite effects of kappa and mu opioid receptors on dopamine release and behavior. *Neuroscience*, 171, 1087-1097.

Bruchas, M., & Aghajanian, A. (2011). The kappa opioid receptor: A key player in the brain's reward system. *Neuroscience*, 171, 1087-1097.

Bruchas, M., & Aghajanian, A. (2011). The kappa opioid receptor: A key player in the brain's reward system. *Neuroscience*, 171, 1087-1097.

Bruchas, M., & Aghajanian, A. (2011). The kappa opioid receptor: A key player in the brain's reward system. *Neuroscience*, 171, 1087-1097.

Bruchas, M., & Aghajanian, A. (2011). The kappa opioid receptor: A key player in the brain's reward system. *Neuroscience*, 171, 1087-1097.

Bruchas, M., & Aghajanian, A. (2011). The kappa opioid receptor: A key player in the brain's reward system. *Neuroscience*, 171, 1087-1097.

Bruchas, M., & Aghajanian, A. (2011). The kappa opioid receptor: A key player in the brain's reward system. *Neuroscience*, 171, 1087-1097.

Bruchas, M., & Aghajanian, A. (2011). The kappa opioid receptor: A key player in the brain's reward system. *Neuroscience*, 171, 1087-1097.

Bruchas, M., & Aghajanian, A. (2011). The kappa opioid receptor: A key player in the brain's reward system. *Neuroscience*, 171, 1087-1097.

Bruchas, M., & Aghajanian, A. (2011). The kappa opioid receptor: A key player in the brain's reward system. *Neuroscience*, 171, 1087-1097.

Acknowledgements:

We thank Catherine Allen, Joseph Brown, Allison Coker, and Peter Fung for technical assistance. This work was supported by funding from the National Institutes of Health (NIH) and the University of California, San Francisco (UCSF) Center for the Neurobiology of Addiction. We also thank Dr. David M. Holtzman for his helpful comments on this manuscript.