

Analysis of Equations of State for Neutron Star Modeling

Joseph Nyhan

College of the Holy Cross

April 27, 2022

Outline

- What is a neutron star?

Outline

- What is a neutron star?
- What is an equation of state (EoS)?

Outline

- What is a neutron star?
- What is an equation of state (EoS)? How do they fit into our model of a neutron star?

- What is a neutron star?
- What is an equation of state (EoS)? How do they fit into our model of a neutron star?
- How can we use an EoS to make macroscopic predictions about neutron stars?

Outline

- What is a neutron star?
- What is an equation of state (EoS)? How do they fit into our model of a neutron star?
- How can we use an EoS to make macroscopic predictions about neutron stars?
- An example and its predictions:

- What is a neutron star?
- What is an equation of state (EoS)? How do they fit into our model of a neutron star?
- How can we use an EoS to make macroscopic predictions about neutron stars?
- An example and its predictions:
 - ▶ QHD-I

Neutron Stars

Neutron Stars

- The collapsed core of a supergiant star

Neutron Stars

- The collapsed core of a supergiant star
- After a supernovae explosion, the dense core is left over

Neutron Stars

- The collapsed core of a supergiant star
- After a supernovae explosion, the dense core is left over
- Radius: ~ 10 km

Neutron Stars

- The collapsed core of a supergiant star
- After a supernovae explosion, the dense core is left over
- Radius: ~ 10 km; Mass: $\sim 1 \odot$.

Neutron Stars

- The collapsed core of a supergiant star
- After a supernovae explosion, the dense core is left over
- Radius: ~ 10 km; Mass: $\sim 1 \odot$.
- Made mostly of neutrons, protons, and electrons

Neutron Stars

- The collapsed core of a supergiant star
- After a supernovae explosion, the dense core is left over
- Radius: ~ 10 km; Mass: $\sim 1 \odot$.
- Made mostly of neutrons, protons, and electrons; overall, is neutral

Neutron Stars

- The collapsed core of a supergiant star
- After a supernovae explosion, the dense core is left over
- Radius: ~ 10 km; Mass: $\sim 1 \odot$.
- Made mostly of neutrons, protons, and electrons; overall, is neutral
- Core held together by intense gravitational attraction

Neutron Stars

- The collapsed core of a supergiant star
- After a supernovae explosion, the dense core is left over
- Radius: ~ 10 km; Mass: $\sim 1 \odot$.
- Made mostly of neutrons, protons, and electrons; overall, is neutral
- Core held together by intense gravitational attraction
 - ▶ Gravitational acceleration on Earth's Surface: $\approx 10 \text{ m/s}^2$

Neutron Stars

- The collapsed core of a supergiant star
- After a supernovae explosion, the dense core is left over
- Radius: ~ 10 km; Mass: $\sim 1 \odot$.
- Made mostly of neutrons, protons, and electrons; overall, is neutral
- Core held together by intense gravitational attraction
 - ▶ Gravitational acceleration on Earth's Surface: $\approx 10 \text{ m/s}^2$
 - ▶ Neutron star: $\approx 10^{12} \text{ m/s}^2$

Neutron Stars

- The collapsed core of a supergiant star
- After a supernovae explosion, the dense core is left over
- Radius: ~ 10 km; Mass: $\sim 1 \odot$.
- Made mostly of neutrons, protons, and electrons; overall, is neutral
- Core held together by intense gravitational attraction
 - ▶ Gravitational acceleration on Earth's Surface: $\approx 10 \text{ m/s}^2$
 - ▶ Neutron star: $\approx 10^{12} \text{ m/s}^2$ (escape velocity $\sim 100\,000 \text{ km/s}$ =

Neutron Stars

- The collapsed core of a supergiant star
- After a supernovae explosion, the dense core is left over
- Radius: ~ 10 km; Mass: $\sim 1 \odot$.
- Made mostly of neutrons, protons, and electrons; overall, is neutral
- Core held together by intense gravitational attraction
 - ▶ Gravitational acceleration on Earth's Surface: $\approx 10 \text{ m/s}^2$
 - ▶ Neutron star: $\approx 10^{12} \text{ m/s}^2$ (escape velocity $\sim 100\,000 \text{ km/s} = c/3$)

Neutron Stars

- The collapsed core of a supergiant star
- After a supernovae explosion, the dense core is left over
- Radius: ~ 10 km; Mass: $\sim 1 \odot$.
- Made mostly of neutrons, protons, and electrons; overall, is neutral
- Core held together by intense gravitational attraction
 - ▶ Gravitational acceleration on Earth's Surface: $\approx 10 \text{ m/s}^2$
 - ▶ Neutron star: $\approx 10^{12} \text{ m/s}^2$ (escape velocity $\sim 100\,000 \text{ km/s} = c/3$)
- Why are they interesting?

Neutron Stars

- The collapsed core of a supergiant star
- After a supernovae explosion, the dense core is left over
- Radius: ~ 10 km; Mass: $\sim 1 \odot$.
- Made mostly of neutrons, protons, and electrons; overall, is neutral
- Core held together by intense gravitational attraction
 - ▶ Gravitational acceleration on Earth's Surface: $\approx 10 \text{ m/s}^2$
 - ▶ Neutron star: $\approx 10^{12} \text{ m/s}^2$ (escape velocity $\sim 100\,000 \text{ km/s} = c/3$)
- Why are they interesting?
 - ▶ Smallest, densest observed stellar objects

Neutron Stars

- The collapsed core of a supergiant star
- After a supernovae explosion, the dense core is left over
- Radius: ~ 10 km; Mass: $\sim 1 \odot$.
- Made mostly of neutrons, protons, and electrons; overall, is neutral
- Core held together by intense gravitational attraction
 - ▶ Gravitational acceleration on Earth's Surface: $\approx 10 \text{ m/s}^2$
 - ▶ Neutron star: $\approx 10^{12} \text{ m/s}^2$ (escape velocity $\sim 100\,000 \text{ km/s} = c/3$)
- Why are they interesting?
 - ▶ Smallest, densest observed stellar objects
 - ▶ Exotic physics

Equation of State (EoS)

What is an equation of state?

Equation of State (EoS)

What is an equation of state?

- A relationship between *energy density* (denoted ϵ)

Equation of State (EoS)

What is an equation of state?

- A relationship between *energy density* (denoted ϵ) and pressure (denoted P)

Equation of State (EoS)

What is an equation of state?

- A relationship between *energy density* (denoted ϵ) and pressure (denoted P)
 - ▶ $\epsilon = \epsilon(P)$

Equation of State (EoS)

What is an equation of state?

- A relationship between *energy density* (denoted ϵ) and pressure (denoted P)
 - ▶ $\epsilon = \epsilon(P) \Leftrightarrow P = P(\epsilon)$

Equation of State (EoS)

What is an equation of state?

- A relationship between *energy density* (denoted ϵ) and pressure (denoted P)
 - ▶ $\epsilon = \epsilon(P) \Leftrightarrow P = P(\epsilon)$
- Encodes the fundamental interparticle interactions within a neutron star

Equation of State (EoS)

What is an equation of state?

- A relationship between *energy density* (denoted ϵ) and pressure (denoted P)
 - ▶ $\epsilon = \epsilon(P) \Leftrightarrow P = P(\epsilon)$
- Encodes the fundamental interparticle interactions within a neutron star
- True EoS within a neutron star is unknown;

Equation of State (EoS)

What is an equation of state?

- A relationship between *energy density* (denoted ϵ) and pressure (denoted P)
 - ▶ $\epsilon = \epsilon(P) \Leftrightarrow P = P(\epsilon)$
- Encodes the fundamental interparticle interactions within a neutron star
- True EoS within a neutron star is unknown; multitude of candidates, each based on a slightly different model

Equation of State (EoS)

What is an equation of state?

- A relationship between *energy density* (denoted ϵ) and pressure (denoted P)
 - ▶ $\epsilon = \epsilon(P) \Leftrightarrow P = P(\epsilon)$
- Encodes the fundamental interparticle interactions within a neutron star
- True EoS within a neutron star is unknown; multitude of candidates, each based on a slightly different model
 - ▶ Models can be very complicated

Equation of State (EoS)

What is an equation of state?

- A relationship between *energy density* (denoted ϵ) and pressure (denoted P)
 - ▶ $\epsilon = \epsilon(P) \Leftrightarrow P = P(\epsilon)$
- Encodes the fundamental interparticle interactions within a neutron star
- True EoS within a neutron star is unknown; multitude of candidates, each based on a slightly different model
 - ▶ Models can be very complicated; often simplifications must be made

Equation of State (EoS)

What is an equation of state?

- A relationship between *energy density* (denoted ϵ) and pressure (denoted P)
 - ▶ $\epsilon = \epsilon(P) \Leftrightarrow P = P(\epsilon)$
- Encodes the fundamental interparticle interactions within a neutron star
- True EoS within a neutron star is unknown; multitude of candidates, each based on a slightly different model
 - ▶ Models can be very complicated; often simplifications must be made

Static Solutions

Static Solutions

- Solve a system of equations to produce an “image” of a neutron star

Static Solutions

- Solve a system of equations to produce an “image” of a neutron star
 - ▶ Called the Tolman-Oppenheimer-Volkoff (TOV) equations

Static Solutions

- Solve a system of equations to produce an “image” of a neutron star
 - ▶ Called the Tolman-Oppenheimer-Volkoff (TOV) equations
- Specify an EoS and a *central pressure* P_0 and get curves describing ϵ , P , and m .

Static Solutions

- Solve a system of equations to produce an “image” of a neutron star
 - ▶ Called the Tolman-Oppenheimer-Volkoff (TOV) equations
- Specify an EoS and a *central pressure* P_0 and get curves describing ϵ , P , and m .
- A solution is known as a *static solution*

Static Solutions

- Solve a system of equations to produce an “image” of a neutron star
 - ▶ Called the Tolman-Oppenheimer-Volkoff (TOV) equations
- Specify an EoS and a *central pressure* P_0 and get curves describing ϵ , P , and m .
- A solution is known as a *static solution*; spherically symmetric

Static Solutions

- Solve a system of equations to produce an “image” of a neutron star
 - ▶ Called the Tolman-Oppenheimer-Volkoff (TOV) equations
- Specify an EoS and a *central pressure* P_0 and get curves describing ϵ , P , and m .
- A solution is known as a *static solution*; spherically symmetric





Figure: Example static solution for $P_0 = 10^{-4}$ for an EoS called “SLy.”

Static Solutions: $M(R)$ and $M(P_0)$ diagrams

Static Solutions: $M(R)$ and $M(P_0)$ diagrams

Use static solutions to make predictions using an EoS:

Static Solutions: $M(R)$ and $M(P_0)$ diagrams

Use static solutions to make predictions using an EoS:

- 1 Create static solutions for a range of P_0 values:

Static Solutions: $M(R)$ and $M(P_0)$ diagrams

Use static solutions to make predictions using an EoS:

- 1 Create static solutions for a range of P_0 values: $P_0 \in [10^{-6}, 10^{-1}]$.

Static Solutions: $M(R)$ and $M(P_0)$ diagrams

Use static solutions to make predictions using an EoS:

- 1 Create static solutions for a range of P_0 values: $P_0 \in [10^{-6}, 10^{-1}]$.
- 2 Find and store the mass M and radius R for each value of P_0 .

Static Solutions: $M(R)$ and $M(P_0)$ diagrams

Use static solutions to make predictions using an EoS:

- 1 Create static solutions for a range of P_0 values: $P_0 \in [10^{-6}, 10^{-1}]$.
- 2 Find and store the mass M and radius R for each value of P_0 .
- 3 Create $M(R)$ and $M(P_0)$ curves

Static Solutions: $M(R)$ and $M(P_0)$ diagrams

Use static solutions to make predictions using an EoS:

- 1 Create static solutions for a range of P_0 values: $P_0 \in [10^{-6}, 10^{-1}]$.
- 2 Find and store the mass M and radius R for each value of P_0 .
- 3 Create $M(R)$ and $M(P_0)$ curves







Figure: Example curves for EoS “SLy.” $1 \odot = 1.989 \times 10^{30}$ kg (solar mass)

Critical Values of P , R , and M




Critical Values of P , R , and M

$M(R) (\odot)$, SLy




$M(P_0) (\odot)$, SLy




Three important values:

Critical Values of P , R , and M

$M(R)$ (\odot), SLy




$M(P_0)$ (\odot), SLy




Three important values: *critical pressure*, *critical mass*, and *critical radius*.

Critical Values of P , R , and M

$M(R) (\odot)$, SLy



$M(P_0) (\odot)$, SLy




Three important values: *critical pressure*, *critical mass*, and *critical radius*.


- Determined by “peaks” of graph

Critical Values of P , R , and M

$M(R)$ (\odot), SLy



$M(P_0)$ (\odot), SLy




Three important values: *critical pressure*, *critical mass*, and *critical radius*.


- Determined by “peaks” of graph
- Maximum mass and radius predicted by EoS

Critical Values of P , R , and M

$M(R) (\odot)$, SLy




$M(P_0) (\odot)$, SLy



Three important values: *critical pressure*, *critical mass*, and *critical radius*.

- Determined by “peaks” of graph
- Maximum mass and radius predicted by EoS
- Largest “stable” pressure

Critical Values of P , R , and M



Three important values: *critical pressure*, *critical mass*, and *critical radius*.

- Determined by “peaks” of graph
- Maximum mass and radius predicted by EoS
- Largest “stable” pressure
- For SLy, $M_{\text{max}} = 2.05 \odot$, $R_{\text{max}} = 9.93 \text{ km}$, and $P_{\text{crit}} = 6.59 \times 10^{-3}$.

Quantum Hadrodynamics

Quantum Hadrodynamics

A theory of the interactions between the subatomic particles in the core of a neutron star

Quantum Hadrodynamics

A theory of the interactions between the subatomic particles in the core of a neutron star

- The protons and neutrons influence one another (e.g. apply forces);

Quantum Hadrodynamics

A theory of the interactions between the subatomic particles in the core of a neutron star

- The protons and neutrons influence one another (e.g. apply forces); mediated by exchanging a particle

Quantum Hadrodynamics

A theory of the interactions between the subatomic particles in the core of a neutron star

- The protons and neutrons influence one another (e.g. apply forces); mediated by exchanging a particle (called a *meson*)

Quantum Hadrodynamics

A theory of the interactions between the subatomic particles in the core of a neutron star

- The protons and neutrons influence one another (e.g. apply forces); mediated by exchanging a particle (called a *meson*)
- Describe these interactions mathematically;

Quantum Hadrodynamics

A theory of the interactions between the subatomic particles in the core of a neutron star

- The protons and neutrons influence one another (e.g. apply forces); mediated by exchanging a particle (called a *meson*)
- Describe these interactions mathematically; the Lagrangian \mathcal{L}

Quantum Hadrodynamics

A theory of the interactions between the subatomic particles in the core of a neutron star

- The protons and neutrons influence one another (e.g. apply forces); mediated by exchanging a particle (called a *meson*)
- Describe these interactions mathematically; the Lagrangian \mathcal{L}

Model requires input from actual experiments

Quantum Hadrodynamics

A theory of the interactions between the subatomic particles in the core of a neutron star

- The protons and neutrons influence one another (e.g. apply forces); mediated by exchanging a particle (called a *meson*)
- Describe these interactions mathematically; the Lagrangian \mathcal{L}

Model requires input from actual experiments

- Extreme conditions in core of a neutron star are not reproducible

Quantum Hadrodynamics

A theory of the interactions between the subatomic particles in the core of a neutron star

- The protons and neutrons influence one another (e.g. apply forces); mediated by exchanging a particle (called a *meson*)
- Describe these interactions mathematically; the Lagrangian \mathcal{L}

Model requires input from actual experiments

- Extreme conditions in core of a neutron star are not reproducible (e.g. pressures are too high, etc.)

Quantum Hadrodynamics

A theory of the interactions between the subatomic particles in the core of a neutron star

- The protons and neutrons influence one another (e.g. apply forces); mediated by exchanging a particle (called a *meson*)
- Describe these interactions mathematically; the Lagrangian \mathcal{L}

Model requires input from actual experiments

- Extreme conditions in core of a neutron star are not reproducible (e.g. pressures are too high, etc.)

Equations of Quantum Hadrodynamics are considered very difficult to solve;

Quantum Hadrodynamics

A theory of the interactions between the subatomic particles in the core of a neutron star

- The protons and neutrons influence one another (e.g. apply forces); mediated by exchanging a particle (called a *meson*)
- Describe these interactions mathematically; the Lagrangian \mathcal{L}

Model requires input from actual experiments

- Extreme conditions in core of a neutron star are not reproducible (e.g. pressures are too high, etc.)

Equations of Quantum Hadrodynamics are considered very difficult to solve; introduce simplifications in the QHD-I model

Quantum Hadrodynamics I (QHD-I)

We form the *Lagrangian* (encodes information about the energy in the system):

Quantum Hadrodynamics I (QHD-I)

We form the *Lagrangian* (encodes information about the energy in the system):

$$\begin{aligned}\mathcal{L} = & \bar{\psi}[\gamma_{\mu}(i\partial^{\mu} - g_v V^{\mu}) - (M - g_{\phi}\phi)]\psi \\ & + \frac{1}{2}(\partial_{\mu}\phi\partial^{\mu}\phi - m_{\phi}^2\phi^2) - \frac{1}{4}V_{\mu\nu}V^{\mu\nu} + \frac{1}{2}m_v^2 V_{\mu}V^{\mu},\end{aligned}$$

Quantum Hadrodynamics I (QHD-I)

We form the *Lagrangian* (encodes information about the energy in the system):

$$\mathcal{L} = \bar{\psi}[\gamma_{\mu}(i\partial^{\mu} - g_v V^{\mu}) - (M - g_{\phi}\phi)]\psi \\ + \frac{1}{2}(\partial_{\mu}\phi\partial^{\mu}\phi - m_{\phi}^2\phi^2) - \frac{1}{4}V_{\mu\nu}V^{\mu\nu} + \frac{1}{2}m_v^2 V_{\mu}V^{\mu},$$

where

- Nucleons (protons and neutrons) ψ

Quantum Hadrodynamics I (QHD-I)

We form the *Lagrangian* (encodes information about the energy in the system):

$$\mathcal{L} = \bar{\psi}[\gamma_{\mu}(i\partial^{\mu} - g_v V^{\mu}) - (M - g_{\phi}\phi)]\psi \\ + \frac{1}{2}(\partial_{\mu}\phi\partial^{\mu}\phi - m_{\phi}^2\phi^2) - \frac{1}{4}V_{\mu\nu}V^{\mu\nu} + \frac{1}{2}m_v^2 V_{\mu}V^{\mu},$$

where

- Nucleons (protons and neutrons) ψ
- Mesons: ϕ and V

Quantum Hadrodynamics I (QHD-I)

We form the *Lagrangian* (encodes information about the energy in the system):

$$\mathcal{L} = \bar{\psi}[\gamma_{\mu}(i\partial^{\mu} - g_v V^{\mu}) - (M - g_{\phi}\phi)]\psi \\ + \frac{1}{2}(\partial_{\mu}\phi\partial^{\mu}\phi - m_{\phi}^2\phi^2) - \frac{1}{4}V_{\mu\nu}V^{\mu\nu} + \frac{1}{2}m_v^2 V_{\mu}V^{\mu},$$

where

- Nucleons (protons and neutrons) ψ
- Mesons: ϕ and V
- Constants from experiment: g_v and g_{ϕ}

Quantum Hadrodynamics I (QHD-I)

We form the *Lagrangian* (encodes information about the energy in the system):

$$\mathcal{L} = \bar{\psi}[\gamma_{\mu}(i\partial^{\mu} - g_v V^{\mu}) - (M - g_{\phi}\phi)]\psi \\ + \frac{1}{2}(\partial_{\mu}\phi\partial^{\mu}\phi - m_{\phi}^2\phi^2) - \frac{1}{4}V_{\mu\nu}V^{\mu\nu} + \frac{1}{2}m_v^2 V_{\mu}V^{\mu},$$

where

- Nucleons (protons and neutrons) ψ
- Mesons: ϕ and V
- Constants from experiment: g_v and g_{ϕ}

From \mathcal{L} , we can determine ϵ and P , the EoS we desire.

RMF Simplifications

We introduce the *Relativistic Mean Field* (RMF) simplifications.

RMF Simplifications

We introduce the *Relativistic Mean Field* (RMF) simplifications. We treat the interactions (exchange of mesons) as their average values:

RMF Simplifications

We introduce the *Relativistic Mean Field* (RMF) simplifications. We treat the interactions (exchange of mesons) as their average values:

$$\phi \rightarrow \langle \phi \rangle = \phi_0, \quad V_\mu \rightarrow \langle V_\mu \rangle = V_0,$$

where ϕ_0 and V_0 are constants.

RMF Simplifications

We introduce the *Relativistic Mean Field* (RMF) simplifications. We treat the interactions (exchange of mesons) as their average values:

$$\phi \rightarrow \langle \phi \rangle = \phi_0, \quad V_\mu \rightarrow \langle V_\mu \rangle = V_0,$$

where ϕ_0 and V_0 are constants. This allows us to simplify \mathcal{L} considerably:

RMF Simplifications

We introduce the *Relativistic Mean Field* (RMF) simplifications. We treat the interactions (exchange of mesons) as their average values:

$$\phi \rightarrow \langle \phi \rangle = \phi_0, \quad V_\mu \rightarrow \langle V_\mu \rangle = V_0,$$

where ϕ_0 and V_0 are constants. This allows us to simplify \mathcal{L} considerably:

$$\mathcal{L}_{\text{RMF}} = \bar{\psi}[i\gamma_\mu\partial^\mu - g_v\gamma_0 V_0 - (M - g_s\phi_0)]\psi - \frac{1}{2}m_s^2\phi_0^2 + \frac{1}{2}m_\omega^2 V_0^2,$$

RMF Simplifications

We introduce the *Relativistic Mean Field* (RMF) simplifications. We treat the interactions (exchange of mesons) as their average values:

$$\phi \rightarrow \langle \phi \rangle = \phi_0, \quad V_\mu \rightarrow \langle V_\mu \rangle = V_0,$$

where ϕ_0 and V_0 are constants. This allows us to simplify \mathcal{L} considerably:

$$\mathcal{L}_{\text{RMF}} = \bar{\psi}[i\gamma_\mu\partial^\mu - g_v\gamma_0 V_0 - (M - g_s\phi_0)]\psi - \frac{1}{2}m_s^2\phi_0^2 + \frac{1}{2}m_\omega^2 V_0^2,$$

Determining ϕ_0 , V_0 , ϵ , and P :

RMF Simplifications

We introduce the *Relativistic Mean Field* (RMF) simplifications. We treat the interactions (exchange of mesons) as their average values:

$$\phi \rightarrow \langle \phi \rangle = \phi_0, \quad V_\mu \rightarrow \langle V_\mu \rangle = V_0,$$

where ϕ_0 and V_0 are constants. This allows us to simplify \mathcal{L} considerably:

$$\mathcal{L}_{\text{RMF}} = \bar{\psi}[i\gamma_\mu \partial^\mu - g_v \gamma_0 V_0 - (M - g_s \phi_0)]\psi - \frac{1}{2}m_s^2 \phi_0^2 + \frac{1}{2}m_\omega^2 V_0^2,$$

Determining ϕ_0 , V_0 , ϵ , and P : $\varphi_\alpha \in \{\phi_0, V_0, \psi\}$

$$\partial_\nu \left(\frac{\partial \mathcal{L}}{\partial(\partial_\nu \varphi_\alpha)} \right) - \frac{\partial \mathcal{L}}{\partial \varphi_\alpha} = 0, \quad T^{\mu\nu} = \frac{\partial \mathcal{L}}{\partial(\partial_\mu \varphi_\alpha)} \partial^\nu \varphi_\alpha - \mathcal{L} \eta^{\mu\nu}.$$

RMF Simplifications

We introduce the *Relativistic Mean Field* (RMF) simplifications. We treat the interactions (exchange of mesons) as their average values:

$$\phi \rightarrow \langle \phi \rangle = \phi_0, \quad V_\mu \rightarrow \langle V_\mu \rangle = V_0,$$

where ϕ_0 and V_0 are constants. This allows us to simplify \mathcal{L} considerably:

$$\mathcal{L}_{\text{RMF}} = \bar{\psi}[i\gamma_\mu \partial^\mu - g_v \gamma_0 V_0 - (M - g_s \phi_0)]\psi - \frac{1}{2}m_s^2 \phi_0^2 + \frac{1}{2}m_\omega^2 V_0^2,$$

Determining ϕ_0 , V_0 , ϵ , and P : $\varphi_\alpha \in \{\phi_0, V_0, \psi\}$

$$\partial_\nu \left(\frac{\partial \mathcal{L}}{\partial (\partial_\nu \varphi_\alpha)} \right) - \frac{\partial \mathcal{L}}{\partial \varphi_\alpha} = 0, \quad T^{\mu\nu} = \frac{\partial \mathcal{L}}{\partial (\partial_\mu \varphi_\alpha)} \partial^\nu \varphi_\alpha - \mathcal{L} \eta^{\mu\nu}.$$

$$\epsilon = \langle T^{00} \rangle, \quad P = \langle T^{ii} \rangle$$

Resulting Equations

From above, we obtain the following equations:

Resulting Equations

From above, we obtain the following equations:

$$\phi_0 = \frac{g_\phi}{m_\phi^2} \frac{1}{\pi^2} \int_0^{k_f} dk \frac{(M - g_\phi \phi_0) k^2}{\sqrt{k^2 + (M - g_\phi \phi_0)}},$$

$$V_0 = \frac{g_v}{m_v^2} \frac{k_f^3}{3\pi^2},$$

$$\epsilon = \frac{1}{2} m_\phi^2 \phi_0^2 + \frac{1}{2} m_v^2 V_0^2 + \frac{1}{\pi^2} \int_0^{k_f} dk k^2 \sqrt{k^2 + m^{*2}},$$

$$P = -\frac{1}{2} m_\phi^2 \phi_0^2 + \frac{1}{2} m_v^2 V_0^2 + \frac{1}{3} \left(\frac{1}{\pi^2} \int_0^{k_f} dk \frac{k^4}{\sqrt{k^2 + m^{*2}}} \right).$$

where $m^* = (M - g_\phi \phi)$, the *reduced mass*.

Resulting Equations

From above, we obtain the following equations:

$$\begin{aligned}\phi_0 &= \phi_0(k_f, \phi_0), & V_0 &= V_0(k_f), \\ \epsilon &= \epsilon(k_f, \phi_0, V_0), & P &= P(k_f, \phi_0, V_0).\end{aligned}$$

Resulting Equations

From above, we obtain the following equations:

$$\begin{aligned}\phi_0 &= \phi_0(k_f, \phi_0), & V_0 &= V_0(k_f), \\ \epsilon &= \epsilon(k_f, \phi_0, V_0), & P &= P(k_f, \phi_0, V_0).\end{aligned}$$

Goal:

Resulting Equations

From above, we obtain the following equations:

$$\begin{aligned}\phi_0 &= \phi_0(k_f, \phi_0), & V_0 &= V_0(k_f), \\ \epsilon &= \epsilon(k_f, \phi_0, V_0), & P &= P(k_f, \phi_0, V_0).\end{aligned}$$

Goal: create a list of values that show us $\epsilon(P)$;

Resulting Equations

From above, we obtain the following equations:

$$\begin{aligned}\phi_0 &= \phi_0(k_f, \phi_0), & V_0 &= V_0(k_f), \\ \epsilon &= \epsilon(k_f, \phi_0, V_0), & P &= P(k_f, \phi_0, V_0).\end{aligned}$$

Goal: create a list of values that show us $\epsilon(P)$; each value of k_f gives us a different ϵ and P .

Resulting Equations

From above, we obtain the following equations:

$$\begin{aligned}\phi_0 &= \phi_0(k_f, \phi_0), & V_0 &= V_0(k_f), \\ \epsilon &= \epsilon(k_f, \phi_0, V_0), & P &= P(k_f, \phi_0, V_0).\end{aligned}$$

Goal: create a list of values that show us $\epsilon(P)$; each value of k_f gives us a different ϵ and P .

To produce the EoS:

Resulting Equations

From above, we obtain the following equations:

$$\begin{aligned}\phi_0 &= \phi_0(k_f, \phi_0), & V_0 &= V_0(k_f), \\ \epsilon &= \epsilon(k_f, \phi_0, V_0), & P &= P(k_f, \phi_0, V_0).\end{aligned}$$

Goal: create a list of values that show us $\epsilon(P)$; each value of k_f gives us a different ϵ and P .

To produce the EoS: (repeat the following)

Resulting Equations

From above, we obtain the following equations:

$$\begin{aligned}\phi_0 &= \phi_0(k_f, \phi_0), & V_0 &= V_0(k_f), \\ \epsilon &= \epsilon(k_f, \phi_0, V_0), & P &= P(k_f, \phi_0, V_0).\end{aligned}$$

Goal: create a list of values that show us $\epsilon(P)$; each value of k_f gives us a different ϵ and P .

To produce the EoS: (repeat the following)

- Choose a k_f value

Resulting Equations

From above, we obtain the following equations:

$$\begin{aligned}\phi_0 &= \phi_0(k_f, \phi_0), & V_0 &= V_0(k_f), \\ \epsilon &= \epsilon(k_f, \phi_0, V_0), & P &= P(k_f, \phi_0, V_0).\end{aligned}$$

Goal: create a list of values that show us $\epsilon(P)$; each value of k_f gives us a different ϵ and P .

To produce the EoS: (repeat the following)

- Choose a k_f value
- calculate ϕ_0 and V_0

Resulting Equations

From above, we obtain the following equations:

$$\begin{aligned}\phi_0 &= \phi_0(k_f, \phi_0), & V_0 &= V_0(k_f), \\ \epsilon &= \epsilon(k_f, \phi_0, V_0), & P &= P(k_f, \phi_0, V_0).\end{aligned}$$

Goal: create a list of values that show us $\epsilon(P)$; each value of k_f gives us a different ϵ and P .

To produce the EoS: (repeat the following)

- Choose a k_f value
- calculate ϕ_0 and V_0 ; use *rootfinding* for ϕ_0

Resulting Equations

From above, we obtain the following equations:

$$\begin{aligned}\phi_0 &= \phi_0(k_f, \phi_0), & V_0 &= V_0(k_f), \\ \epsilon &= \epsilon(k_f, \phi_0, V_0), & P &= P(k_f, \phi_0, V_0).\end{aligned}$$

Goal: create a list of values that show us $\epsilon(P)$; each value of k_f gives us a different ϵ and P .

To produce the EoS: (repeat the following)

- Choose a k_f value
- calculate ϕ_0 and V_0 ; use *rootfinding* for ϕ_0
- Using those values, calculate P and ϵ and store in a table

$M(R)$ and $M(P_0)$ Curves for QHD-I

We use the tabulated values of P and ϵ to solve the TOV equations:

$M(R)$ and $M(P_0)$ Curves for QHD-I

We use the tabulated values of P and ϵ to solve the TOV equations:




Figure: $M(R)$ and $M(P_0)$ curves for QHD-I EoS.

$M(R)$ and $M(P_0)$ Curves for QHD-I

We use the tabulated values of P and ϵ to solve the TOV equations:




Figure: $M(R)$ and $M(P_0)$ curves for QHD-I EoS.

These curves give

$$M_{\max} = 6.1 \odot, \quad R_{\max} = 42.1 \text{ km}, \quad P_{\text{crit}} = 1.98 \times 10^{-4}.$$

Conclusion

- An equation of state is a relationship between energy density and pressure within a neutron star

Conclusion

- An equation of state is a relationship between energy density and pressure within a neutron star
- We use the TOV equations to predict the maximum mass and radius that a given EoS will produce

Conclusion

- An equation of state is a relationship between energy density and pressure within a neutron star
- We use the TOV equations to predict the maximum mass and radius that a given EoS will produce
- We use the QHD-I parameter set and RMF simplifications to solve a system of equations and generate an equation of state

Thanks

- Prof. Ben Kain

Thanks

- Prof. Ben Kain
- Holy Cross Physics Department

Thanks

- Prof. Ben Kain
- Holy Cross Physics Department
- My family

Thanks

- Prof. Ben Kain
- Holy Cross Physics Department
- My family
- Special thanks: Prof. Perry and the AV team