

Sequences

12 February 2020 10:12

We expect a sequence (a_n) to converge to a limit l if its terms eventually get close to l . To make this precise, a number ϵ , say, is used as a measure of "closeness to l "

Definition of a convergent sequence. A sequence (a_n) of real numbers is said to *converge* to a limit $l \in \mathbb{R}$ if for every $\epsilon > 0$ there is an integer N (which depends on ϵ) with $|a_n - l| < \epsilon$ for all $n > N$. When (a_n) converges to l we write

$$\lim_{n \rightarrow \infty} a_n = l \quad \text{or} \quad a_n \rightarrow l.$$

Note that $|a_n - l|$ is the distance between the points a_n and l on the real line. The definition says that *no matter how small* a positive number ϵ we take, the distance between a_n and l will eventually be smaller than ϵ , i.e., the numbers a_n will eventually lie between $l - \epsilon$ and $l + \epsilon$.

A useful tool to help identify whether a sequence converges is to break it down into a series of simpler sequences and then use the rules below:

Combination Rules for Convergent Sequences

If $(a_n), (b_n), (c_n)$ are convergent sequences with $a_n \rightarrow \alpha, b_n \rightarrow \beta, c_n \rightarrow \gamma$, then:

sum rule	$a_n + b_n \rightarrow \alpha + \beta$	[CR(1)]
scalar multiple rule	$\lambda a_n \rightarrow \lambda \alpha \quad (\text{for } \lambda \in \mathbb{R})$	[CR(2)]
product rule	$a_n b_n \rightarrow \alpha \beta$	[CR(3)]
reciprocal rule	$1/a_n \rightarrow 1/\alpha \quad (\text{provided } \alpha \neq 0)$	[CR(4)]
quotient rule	$b_n/a_n \rightarrow \beta/\alpha \quad (\text{provided } \alpha \neq 0)$	[CR(5)]
'hybrid' rule	$b_n c_n/a_n \rightarrow \beta \gamma/\alpha \quad (\text{provided } \alpha \neq 0)$	[CR(6)]

Problem. Show that the sequence (a_n) defined by

$$a_n = \frac{(n+2)(2n-1)}{3n^2+1}$$

converges and find its limit.

We cannot use the quotient rule directly since the sequences in the numerator and denominator do not individually converge. Examples like this can be tackled by dividing throughout by the term which increases fastest.

Solution. Dividing numerator and denominator by n^2 gives

$$a_n = \frac{(n+2)(2n-1)}{3n^2+1} = \frac{(1+2/n)(2-1/n)}{3+1/n^2}.$$

Now $(1/n)$ and $(1/n^2)$ converge to 0, so applying the combination rules CR(1,2,3 and 5) gives

$$a_n \rightarrow \frac{(1+0)(2-0)}{3+0} \quad \text{i.e.,} \quad a_n \rightarrow \frac{2}{3}.$$

Subsequences

A subsequence is a sequence with some terms deleted.

Bounds

A sequence is bounded above/below if there exists a U/L for which all $a_n < U$ or $a_n > L$.

A sequence is called bounded if it is both bounded below and above.

Increasing / Decreasing Sequences

Divergent Sequences

A sequence a_n is said to diverge to infinity if for every $k \in \mathbb{R}$ there is an N with $a_n > k$ whenever $n > N$. If a_n diverges to infinity we write $a_n \rightarrow \infty$.

E.g. fibonacci sequence diverges to infinity

$-n$ diverges to $-\infty$

$(-1)^n$ oscillates

Basic Convergent Sequences

Basic properties of convergent sequences

- (1) A convergent sequence has a unique limit.
- (2) If $a_n \rightarrow l$, then every subsequence of (a_n) also converges to l .
- (3) If $a_n \rightarrow l$ then $|a_n| \rightarrow |l|$.
- (4) The squeeze rule. If $a_n \rightarrow l$ and $b_n \rightarrow l$ and $a_n \leq c_n \leq b_n$ for all n , then $c_n \rightarrow l$.

$$c_n \rightarrow l.$$

- (5) A convergent sequence (a_n) is bounded, i.e., there is a $B > 0$ with $-B \leq a_n \leq B$ for all n .
- (6) Any increasing sequence which is bounded above converges. Any decreasing sequence which is bounded below converges.

e.g. to prove that -1^n does not converge, we observe that the even subsequence is $1, 1, 1, \dots$ which converges to 1, and the odd subsequence is $-1, -1, -1, \dots$ which converges to -1 . However, as per boxed rule 2, if a sequence converges then all of its subsequence's also converge to the same limit l . This is not the case and as such the sequence does not converge.

Basic Convergent Sequences

Basic Convergent Sequences

- (1) $\lim_{n \rightarrow \infty} \frac{1}{n^p} = 0$ for any $p > 0$
- (2) $\lim_{n \rightarrow \infty} c^n = 0$ for any c with $|c| < 1$
- (3) $\lim_{n \rightarrow \infty} c^{1/n} = 1$ for any $c > 0$
- (4) $\lim_{n \rightarrow \infty} n^p c^n = 0$ for $p > 0$ and $|c| < 1$
- (5) $\lim_{n \rightarrow \infty} \frac{c^n}{n!} = 0$ for any $c \in \mathbb{R}$
- (6) $\lim_{n \rightarrow \infty} \left(1 + \frac{c}{n}\right)^n = e^c$ for any $c \in \mathbb{R}$