

Advanced Mixed Integer Programming Formulation Techniques

Constructing MIP Formulations using Convex Hulls

Joey Huchette and Juan Pablo Vielma

Massachusetts Institute of Technology

Spring School of the International Symposium on
Combinatorial Optimization (ISCO 2018)

Marrakesh, Morocco, April, 2018


Course Material

- Code:
 - <https://github.com/joehuchette/ISCO-spring-school/tree/master/code>
- Slides:
 - <https://github.com/joehuchette/ISCO-spring-school/tree/master/slides>

Constructing Ideal Formulations For Pure Integer

- Pure Integer :

$$Q := \text{conv} \left(\{p^i\}_{i=1}^n \right)$$



Integral Formulations for Small Sets

- $S = \{x \in \{0,1\}^3 : x_3 = x_1 \times x_2\}$
- Constraints = $\text{conv}(S)$
 - $\text{conv}(W) := \left\{ \sum_{w \in W} w \lambda_w : \lambda \in \Delta^W \right\}$
 - $\Delta^W := \left\{ \lambda \in \mathbb{R}_+^W : \sum_{w \in W} \lambda_w = 1 \right\}$
- $S = \{(0,0,0), (1,0,0), (0,1,0), (1,1,1)\}$
- $\text{conv}(S) :$

Integral Formulations for Small Sets

```
julia> using Polyhedra, CDDLib, JuMP
julia> points = SimpleVRepresentation([0 0 0; 1 0 0; 0 1 0; 1 1 1]);
julia> poly = polyhedron(points, CDDLibrary(:exact))
julia> ineq = SimpleHRepresentation(poly)
H-representation
begin
  4 4 rational
  1//1 -1//1 -1//1 1//1
  0//1 1//1 0//1 -1//1
  0//1 0//1 1//1 -1//1
  0//1 0//1 0//1 1//1
end
julia> model = Model(); @variable(model,x[1:3]);
julia> @constraint(model,convert.(Int64,ineq.A)*x .<= convert.(Int64,ineq.b))
4-element
Array{JuMP.ConstraintRef{JuMP.Model,JuMP.GenericRangeConstraint{JuMP.GenericAffExpr{Float64,JuMP.Variable}}},1}:
 x[1] + x[2] - x[3] ≤ 1
 -x[1] + x[3] ≤ 0
 -x[2] + x[3] ≤ 0
 -x[3] ≤ 0
```

Integral Formulations for Small Sets

- $S = \{x \in \{0,1\}^3 : x_3 = x_1 \times x_2\}$
- Constraints = $\text{conv}(S)$
 - $\text{conv}(W) := \left\{ \sum_{w \in W} w \lambda_w : \lambda \in \Delta^W \right\}$
 - $\Delta^W := \left\{ \lambda \in \mathbb{R}_+^W : \sum_{w \in W} \lambda_w = 1 \right\}$
- $S = \{(0,0,0), (1,0,0), (0,1,0), (1,1,1)\}$
- $\text{conv}(S)$:
 - $x_1 + x_2 \leq 1 + x_3$
 - $x_3 \leq x_1$
 - $x_3 \leq x_2$
 - $0 \leq x_3$
- Always works for 0-1 problems

Careful With General Integer Problems

- All different : $S = \{x \in [n]^n : x_i \neq x_j \ \forall i \neq j\}$
- Permutahedron:
 - $\text{conv}(S) = \left\{ x \in \mathbb{R}^n : \begin{array}{l} \sum_{i=1}^n x_i = \sum_{i=1}^n i \\ \sum_{i \in I} x_i \geq \binom{|I|+1}{2} \ \forall I: \emptyset \neq I \subseteq [n] \end{array} \right\}$
- If n is odd then:
 - $\left(\frac{n+1}{2}, \dots, \frac{n+1}{2}\right) \in \text{conv}(S) \cap \mathbb{Z}^n$
 - $\left(\frac{n+1}{2}, \dots, \frac{n+1}{2}\right) \notin S$
- “Hole-Free”: $S \subseteq \mathbb{R}^n$ such that $\text{conv}(S) \cap \mathbb{Z}^n = S$


Solution: Extended Formulations

- (Finite?) Non-hole free subset of integers is always projection of hole-free subset of integers
- All different : $S = \{x \in [n]^n : x_i \neq x_j \ \forall i \neq j\}$
- $T := \left\{ (x, Y) \in \mathbb{R}^n \times \{0,1\}^{n \times n} : \begin{array}{l} \sum_{i=1}^n Y_{i,j} = 1 \ \forall j \in [n] \\ \sum_{j=1}^n Y_{i,j} = 1 \ \forall i \in [n] \\ \sum_{j=1}^n j Y_{i,j} = x_i \ \forall i \in [n] \end{array} \right\}$
- $S = \text{proj}_x(T)$

What About Mixed-Integer Sets?


- Pure Integer $S \subseteq \mathbb{Z}^n$

$$P \cap \mathbb{Z}^n = S \quad (P \subseteq \mathbb{R}^n)$$



- Mixed-Integer $S = \bigcup_{i=1}^n P^i \subseteq \mathbb{R}^n$


$$Q := \text{conv}(S)?$$



What is a MIP Formulation?

Simple Formulation for Univariate Functions

$$z = f(x)$$



Size = $O(\# \text{ of segments})$

Non-Ideal: Fractional Extreme Points

$$\begin{pmatrix} x \\ z \end{pmatrix} = \sum_{j=1}^5 \begin{pmatrix} d_j \\ f(d_j) \end{pmatrix} \lambda_j$$

$$1 = \sum_{j=1}^5 \lambda_j, \quad \lambda_j \geq 0$$

$$y \in \{0, 1\}^4, \quad \sum_{i=1}^4 y_i = 1$$

$$0 \leq \lambda_1 \leq y_1$$

$$0 \leq \lambda_2 \leq y_1 + y_2$$


$$0 \leq \lambda_3 \leq y_2 + y_3$$

$$0 \leq \lambda_4 \leq y_3 + y_4$$

$$0 \leq \lambda_5 \leq y_4$$

Advanced Formulation for Univariate Functions

$$z = f(x)$$



Size = $O(\log_2 \# \text{ of segments})$

Ideal: Integral Extreme Points

$$\begin{pmatrix} x \\ z \end{pmatrix} = \sum_{j=1}^5 \begin{pmatrix} d_j \\ f(d_j) \end{pmatrix} \lambda_j$$

$$1 = \sum_{j=1}^5 \lambda_j, \quad \lambda_j \geq 0$$

$$y \in \{0, 1\}^2$$

$$0 \leq \lambda_1 + \lambda_5 \leq 1 - y_1$$



$$0 \leq \lambda_3 \leq y_1$$

$$0 \leq \lambda_4 + \lambda_5 \leq 1 - y_2$$


$$0 \leq \lambda_1 + \lambda_2 \leq y_2$$

Computational Performance

- Advanced formulations provide an computational advantage
- Advantage is significantly more important for free solvers
- State of the art commercial solvers can be significantly better than free solvers
- Still, free is free!




Formulation Improvements can be Significant



Abstracting Univariate Functions

$$P_i := \left\{ \lambda \in \Delta^5 : \lambda_j = 0 \quad \forall j \notin T_i \right\}$$

$$T_i := \{i, i+1\} \quad i \in \{1, \dots, 4\}$$




Δ^5

=

$$1 = \sum_{j=1}^5 \lambda_j, \quad \lambda_j \geq 0$$



Special Ordered Sets
of Type 2 (SOS2)



$$\lambda \in \bigcup_{i=1}^4 P_i \subseteq \Delta^5$$

Abstraction Works for Multivariate Functions

$$P_i := \{\lambda \in \Delta^m : \lambda_j = 0 \quad \forall v_j \notin T_i\}$$



$$\lambda \in \bigcup_{i=1}^n P_i \subseteq \Delta^m$$

Standard Formulation for SOS2 = Unary Encoding

$Q = \text{LP relaxation} \rightarrow$

$$y \in \{0, 1\}^4,$$

$$\begin{aligned} \sum_{i=1}^5 \lambda_i &= 1 \\ \sum_{i=1}^4 y_i &= 1 \end{aligned}$$

$$\begin{aligned} 0 &\leq \lambda_1 \leq y_1 \\ 0 &\leq \lambda_2 \leq y_1 + y_2 \\ 0 &\leq \lambda_3 \leq y_2 + y_3 \\ 0 &\leq \lambda_4 \leq y_3 + y_4 \\ 0 &\leq \lambda_5 \leq y_4 \end{aligned}$$

$$(\lambda, y) \in Q \cap (\mathbb{R}^5 \times \mathbb{Z}^4)$$

$$\Updownarrow$$

$$y = e^i \wedge \lambda \in P_i$$

$$P_i := \{\lambda \in \Lambda^5 : \lambda_j = 0 \quad j \notin \{i, i+1\}\}$$

Unary Encoding

Advanced = Binary Encoded Formulation

$Q = \text{LP relaxation} \rightarrow$

$$\sum_{i=1}^5 \lambda_i = 1$$

$$h^1 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, h^2 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, h^3 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, h^4 = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

- V. and Nemhauser '08.

$$0 \leq \lambda_1 + \lambda_5 \leq 1 - y_1$$

$$0 \leq \lambda_3 \leq y_1$$

$$0 \leq \lambda_4 + \lambda_5 \leq 1 - y_2$$

$$0 \leq \lambda_1 + \lambda_2 \leq y_2$$

$$(\lambda, y) \in Q \cap (\mathbb{R}^5 \times \mathbb{Z}^2)$$




$$\Leftrightarrow$$

$$y = h^i \wedge \lambda \in P_i$$

$$P_i := \{ \lambda \in \Lambda^5 : \lambda_j = 0 \quad j \notin \{i, i+1\} \}$$

Binary Encoding

Embedding Formulation = Ideal non-Extended




$$Q(H) := \text{conv} \left(\bigcup_{i=1}^n P_i \times \{h^i\} \right)$$

$$(x, y) \in Q \cap (\mathbb{R}^d \times \mathbb{Z}^k) \iff y = h^i \wedge x \in P_i$$

$$\text{ext}(Q) \subseteq \mathbb{R}^d \times \mathbb{Z}^k \quad H := \{h^i\}_{i=1}^n \subseteq \{0, 1\}^k, \quad h^i \neq h^j$$

Alternative Encodings

- Careful with general integers:




- Options for 0-1 encodings:
 - Traditional or **Unary** encoding

$$H = \left\{ y \in \{0, 1\}^n : \sum_{i=1}^n y_i = 1 \right\}$$

$$= \{e^i\}_{i=1}^n$$

$$e_j^i = \begin{cases} 1 & i = j \\ 0 & i \neq j \end{cases}$$

- Binary** encodings: $H \equiv \{0, 1\}^{\log_2 n}$
- Others (e.g. **incremental** encoding \equiv unary)



Embedding Formulations and Complexity


- **Embedding formulation** of $\lambda \in \bigcup_{i=1}^n P_i \subseteq \mathbb{R}^V$:
 - Encoding $H := \{h^i\}_{i=1}^n \subseteq \{0, 1\}^k$, $h^i \neq h^j$
 - $Q(H) := \text{conv} \left(\bigcup_{i=1}^n P_i \times \{h^i\} \right)$
- **Embedding complexity** = size smallest formulation
 - $\text{mc}(\mathcal{P}) := \min_H \{\text{size}(Q(H))\}$,
 $\text{size}(Q(H)) := \# \text{ facets}$

Special Ordered Sets = Simplex Faces = $\mathcal{P} := \{P_i\}_{i=1}^n$

- $\Delta^{d+1} := \left\{ x \in \mathbb{R}_+^{d+1} : \sum_{i=1}^{d+1} x_i = 1 \right\} = \text{conv} \left(\{e^i\}_{i=1}^{d+1} \right)$
 $P_i := \text{conv} \left(\{e^j\}_{j \in T_i} \right) = \left\{ x \in \Delta^{d+1} : \sum_{j \notin T_i} x_j \leq 0 \right\}$
 $T_i \subseteq \{1, \dots, d+1\}$
- $\text{mc}(\mathcal{P}) := \min_H \{ \text{size}(Q(H)) \},$
 $\text{size}(Q(H)) := \# \text{ facets}$
- $\text{mc}_G(\mathcal{P}) := \min_H \{ \text{size}_G(Q(H)) \},$
 $\text{size}_G(Q(H)) := \# \text{ non-bound facets}$

Special Ordered Sets of Type 2 (SOS2) = $\mathcal{P} := \{P_i\}_{i=1}^n$

- $P_i := \text{conv}(\{e^i, e^{i+1}\}) \subseteq \Delta^{n+1}, \quad i \in [n]$



,
 $2 \lceil \log_2 n \rceil$

$\{e^i\}$

$\{b \cdot h^{i+1}\}$

$$b \cdot \underbrace{(h^{i+1} - h^i)} = 0$$

Embedding Formulation for SOS2: Part 1

- From encodings (H) to hyperplanes:

$$\{h^i\}_{i=1}^n$$

$$h^1 = \begin{pmatrix} 1 \\ 1 \end{pmatrix}, h^2 = \begin{pmatrix} 0 \\ 0 \end{pmatrix}, h^3 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$$

$$c^i = h^{i+1} - h^i$$




non-bound facets = $2 \times$ # of hyperplanes

$$\bigcup_{i=1}^n$$

Hyperplanes spanned by

$$\{b^i \cdot y = 0\}_{j=1}^L$$



Embedding Formulation for SOS2: Part 2

$$Q(H) = \left\{ b^i \cdot y = 0 \right\}_{j=1}^L$$

$$L(H) := \text{aff}(H) - h^1$$

$$\begin{aligned} & (b^j \cdot h^1) x_1 + \sum_{i=2}^n \min \{ b^j \cdot h^i, b^j \cdot h^{i-1} \} x_i + (b^j \cdot h^n) x_{n+1} \leq b^j \cdot y \quad \forall j \\ & - (b^j \cdot h^1) x_1 - \sum_{i=2}^n \max \{ b^j \cdot h^i, b^j \cdot h^{i-1} \} x_i - (b^j \cdot h^n) x_{n+1} \leq -b^j \cdot y \quad \forall j \\ & \sum_{i=1}^{n+1} x_i = 1, \quad x \in \mathbb{R}_+^{n+1} \\ & y \in L(H) \end{aligned}$$

- # non-bound facets = $2 \times$ # of hyperplanes

Embedding Complexity for SOS2

- Lower Bound: $L(H) := \text{aff}(H) - h^1$

$$\text{mc}_G(\mathcal{P}) \geq 2 \times \min \# \text{ of hyperplanes}$$

$$\min \# \text{ of hyperplanes} \geq \dim(L(H))$$

$$\dim(L(H)) \geq \lceil \log_2 n \rceil$$

- Upper Bound: $H = \{0, 1\}^{\lceil \log_2 n \rceil}$

$$\text{-- Gray code: } \{h^i - h^{i+1}\}_{i=1}^{n-1} \equiv \{e^i\}_{i=1}^{\lceil \log_2 n \rceil}$$

$$\text{size}_G(Q(H)) = 2 \lceil \log_2 n \rceil$$

$$n + 1 \leq \text{mc}(\mathcal{P}) \leq n + 1 + 2 \lceil \log_2 n \rceil$$

Embedding Complexity for SOS2

- Unary encoding (Padberg / Lee and Wilson, early 00's):

$$\text{size}_G(Q(H)) = 2(n - 1), \quad \text{size}(Q(H)) = 2n$$

- Smallest Binary** encoding (V. and Nemhauser '08, Muldoon '12):

$$\text{size}_G(Q(H)) = 2 \lceil \log_2 n \rceil,$$

$$2 + 2 \lceil \log_2 n \rceil \leq \text{size}(Q(H)) \leq n + 1 + 2 \lceil \log_2 n \rceil$$

- Adding lower bounds (# hyperplanes \geq dimension):

$$\text{mc}_G(\mathcal{P}) = 2 \lceil \log_2 n \rceil,$$

$$n + 1 \leq \text{xc}(\mathcal{P}) \leq \text{mc}(\mathcal{P}) \leq n + 1 + 2 \lceil \log_2 n \rceil$$

Validity of Formulation May Not Be Evident

$$Q(H) = \bigcap_{j=1}^L \{b^j \cdot y = 0\} \quad L(H) := \text{aff}(H) - h^1$$

$$\begin{aligned} (b^j \cdot h^1) x_1 + \sum_{i=2}^n \min \{b^j \cdot h^i, b^j \cdot h^{i-1}\} x_i + (b^j \cdot h^n) x_{n+1} &\leq b^j \cdot y \quad \forall j \\ - (b^j \cdot h^1) x_1 - \sum_{i=2}^n \max \{b^j \cdot h^i, b^j \cdot h^{i-1}\} x_i - (b^j \cdot h^n) x_{n+1} &\leq -b^j \cdot y \quad \forall j \\ \sum_{i=1}^{n+1} x_i &= 1, \quad x \in \mathbb{R}_+^{n+1} \\ y &\in L(H) \end{aligned}$$

- # non-bound facets = $2 \times$ # of hyperplanes

Validity of Formulation May Not Be Evident

- $H = (0, 1, 1, 1)^T, (0, 1, 0, 0)^T, (0, 0, 0, 0)^T, (0, 1, 0, 1)^T, (0, 0, 0, 1)^T, (1, 0, 0, 0)^T, (1, 1, 0, 1)^T, (1, 0, 1, 1)^T, (1, 1, 1, 1)^T$
- $(c^i)_{i=1}^8 = (0, 0, -1, -1)^T, (0, -1, 0, 0)^T, (0, 1, 0, 1)^T, (0, -1, 0, 0)^T, (1, 0, 0, -1)^T, (0, 1, 0, 1)^T, (0, -1, 1, 0)^T, (0, 1, 0, 0)^T$
- $b^1 = (1, 0, 0, -1, 1)^T, b^2 = (1, 0, 0, 1)^T, b^3 = (1, -1, -1, 1)^T, b^4 = (1, 0, 0, 0)^T$ and $b^5 = (0, 0, 1, 0)^T$

Validity of Formulation May Not Be Evident

$$\sum_{j=1}^{10} \lambda_j = 1,$$

$$\lambda_5 + \lambda_6 + \lambda_7 + \lambda_8 + \lambda_9 + \lambda_{10} \leq y_1 - y_3 + y_4$$

$$\lambda_4 + \lambda_5 + \lambda_6 + 2\lambda_7 + 2\lambda_8 + \lambda_9 + \lambda_{10} \geq y_1 - y_3 + y_4$$

$$\lambda_1 + \lambda_5 + \lambda_6 + \lambda_7 + 2\lambda_8 + 2\lambda_9 + 2\lambda_{10} \leq y_1 + y_4$$

$$\lambda_1 + \lambda_2 + \lambda_4 + \lambda_5 + \lambda_6 + 2\lambda_7 + 2\lambda_8 + 2\lambda_9 + 2\lambda_{10} \geq y_1 + y_4$$

$$-\lambda_1 - \lambda_2 - \lambda_3 + \lambda_6 + \lambda_7 + \lambda_8 \leq y_1 - y_2 - y_3 + y_4$$

$$-\lambda_1 - \lambda_2 + \lambda_5 + \lambda_6 + \lambda_7 + \lambda_8 + \lambda_9 \geq y_1 - y_2 - y_3 + y_4$$

$$\lambda_7 + \lambda_8 + \lambda_9 + \lambda_{10} \leq y_1$$

$$\lambda_6 + \lambda_7 + \lambda_8 + \lambda_9 + \lambda_{10} \geq y_1$$



$$\lambda_1 + \lambda_9 + \lambda_{10} \leq y_3$$

$$\lambda_1 + \lambda_2 + \lambda_8 + \lambda_9 + \lambda_{10} \geq y_3$$

$$\lambda_j \geq 0$$

Abstraction Works for Multivariate Functions

$$P_i := \{\lambda \in \Delta^m : \lambda_j = 0 \quad \forall v_j \notin T_i\}$$



$$\lambda \in \bigcup_{i=1}^n P_i \subseteq \Delta^m$$

Formulations and Complexity for Triangulations

- Lower bound:

$$\left(\sqrt{n/2} + 1\right)^2 \leq \text{mc}(\mathcal{P})$$

- Size of unary formulation is:

(Lee and Wilson '01)



$$\text{mc}(\mathcal{P}) \leq \binom{2\sqrt{n/2}}{\sqrt{n/2}} + \left(\sqrt{n/2} + 1\right)^2$$

- Small binary formulation for **union jack triangulation** of size:

(V. and Nemhauser '08)

$$\text{mc}(\mathcal{P}) \leq 4\log_2 \sqrt{n/2} + 2 + \left(\sqrt{n/2} + 1\right)^2$$

$$n = 2m^2$$




Encoding Selection Matters

- Size of unary formulation is:
(Lee and Wilson '01)

$$\binom{2\sqrt{n/2}}{\sqrt{n/2}} + \left(\sqrt{n/2} + 1\right)^2$$

General Inequalities

Variable Bounds






- Size of one binary formulation:
(V. and Nemhauser '08)

$$4 \log_2 \sqrt{n/2} + 2 + \left(\sqrt{n/2} + 1 \right)^2$$

- Right embedding = significant computational advantage over alternatives (Extended, Big-M, etc.)


Unary Encoding, Minkowski Sum and Cayley Trick



$$Q \cap (\mathbb{R}^2 \times \{0.5\}) \equiv P_1 + P_2 =$$

$$H = \{e^i\}_{i=1}^n$$

$$Q(H) \cap (\mathbb{R}^d \times \{\frac{1}{n} \sum_{i=1}^n e^i\}) \equiv \sum_{i=1}^n P_i$$



Faces of Cayley Embedding


- Two types of facets (or faces):

- $P_1 \times \{0\} \equiv y_i \geq 0$


- $\text{conv}((F_1 \times 0) \cup (F_2 \times 1))$

F_i proper face of P_i

- Not all combinations of faces
 - Which ones are valid?





Valid Combinations = Common Normals



$$N(F_1) \cap N(F_2) \neq \emptyset$$



$\text{conv}((F_1 \times 0) \cup (F_2 \times 1))$
is face of $Q(H)$



Redundancy in Embedding Formulations

Formulations and Complexity for Triangulations

- Lower bound:

$$\left(\sqrt{n/2} + 1\right)^2 \leq \text{mc}(\mathcal{P})$$

- Size of unary formulation is:

(Lee and Wilson '01)



$$\text{mc}(\mathcal{P}) \leq \binom{2\sqrt{n/2}}{\sqrt{n/2}} + \left(\sqrt{n/2} + 1\right)^2$$

- Small binary formulation for **union jack triangulation** of size:

(V. and Nemhauser '08)


$$\text{mc}(\mathcal{P}) \leq 4\log_2 \sqrt{n/2} + 2 + \left(\sqrt{n/2} + 1\right)^2$$

$$n = 2m^2$$



What About Other Triangulations

- $\min_{H \in \mathcal{H}_3(8)} \text{size}_M(Q(\mathcal{T}, H)) \geq 9.$



$$\lambda_{(1,1)} + \lambda_{(3,3)} \leq 1 - y_1,$$

$$\lambda_{(1,3)} + \lambda_{(2,2)} + \lambda_{(3,1)} \leq y_1$$

$$\lambda_{(1,2)} + \lambda_{(2,1)} \leq 1 - y_2,$$

$$\lambda_{(2,3)} + \lambda_{(3,2)} \leq y_2$$

$$\lambda_{(1,1)} + \lambda_{(2,1)} + \lambda_{(3,1)} \leq 1 - y_3,$$


$$\lambda_{(1,3)} + \lambda_{(2,3)} + \lambda_{(3,3)} \leq y_3$$

$$\lambda_{(1,1)} + \lambda_{(1,2)} + \lambda_{(1,3)} \leq 1 - y_4,$$

$$\lambda_{(3,1)} + \lambda_{(3,2)} + \lambda_{(3,3)} \leq y_4$$

$$\sum_{v \in V} \lambda_v = 1, \quad \lambda \in \mathbb{R}_+^V, \quad y \in \{0, 1\}^4.$$

Redundant Embedding Formulation



- $\text{size}_M \left(Q \left(\mathcal{T}, \{h^i\}_{i=1}^8 \right) \right) = 19$
- v/s 8 inequalities