

Advanced Mixed Integer Programming Formulation Techniques

Nonlinear MIP Formulations

Joey Huchette and Juan Pablo Vielma

Massachusetts Institute of Technology

Spring School of the International Symposium on
Combinatorial Optimization (ISCO 2018)

Marrakesh, Morocco, April, 2018

(Nonlinear) Mixed Integer Programming (MIP)



$$\min f(x)$$

s.t.

$$x \in C$$

$$x_i \in \mathbb{Z} \quad i \in I$$

Mostly **convex** f and C .



“Convex” Nonlinear MIP Problem

$$\max \quad g(x) + h(y)$$

s.t.

$$f_i(x, y) \leq b \quad \forall i \in [k]$$

$$x \in \mathbb{Z}^n$$

$$y \in \mathbb{R}^m$$

g , h and f_i convex

- Only non-convexities are due to integrality

MI Second Order Cone Programming (MISOCP)

$$\max \quad c \cdot x + h \cdot y$$

s.t.

$$\|D^i x + E^i y\|_2 \leq f^i \cdot x + g^i \cdot y + c_0^i \quad \forall i \in \{1, \dots, k\}$$

$$Ax + By \leq b$$

Second Order Conic or Conic Quadratic Problems

- Problems using **Euclidean norm**:
 - e.g. Portfolio Optimization Problems

$$\max \quad \bar{a}x$$

s.t.

$$\|Q^{1/2}x\|_2 \leq \sigma$$

$$\sum_{j=1}^n x_j = 1, \quad x \in \mathbb{R}_+^n$$

$$x_j \leq z_j \quad \forall j \in [n]$$

$$\sum_{j=1}^n z_j \leq K, \quad z \in \{0, 1\}^n$$

- \bar{a} expected returns.
- $Q^{1/2}$ square root of covariance matrix.
- K maximum number of assets.
- σ maximum risk.




Conic Quadratic or Second Order Cone

$$\|D^i x + E^i y\|_2 \leq f^i \cdot x + g^i \cdot y + c_0^i \quad \forall i \in \{1, \dots, k\}$$

- Or linear inequalities +

$$- \sum_{i=1}^{m-1} y_i^2 \leq y_m^2, \quad y_m \geq 0$$

$$- y_1^2 \leq y_2 y_3, \quad y_2, y_3 \geq 0$$



Mixed Integer Conic Programming

- \mathcal{K}
 - Convex cone = closed under sum and scaling
 - Closed (usually pointed and maybe more)

$$P : \quad \min \quad \mathbf{c}'\mathbf{x}$$

s.t.

$$\mathbf{b} - \mathbf{A}\mathbf{x} \in \mathcal{K}^c \quad \mathbf{x} \in \mathcal{K}^v$$

$$x_i \in \mathbb{Z} \quad \forall x \in I$$

Conic Duality

- Cone dual to \mathcal{K}
 - $\mathcal{K}^* = \{\mathbf{y} \in \mathbb{R}^n : \mathbf{x}'\mathbf{y} \geq 0, \forall \mathbf{x} \in \mathcal{K}\}$
 - $\mathcal{K} = \mathcal{K}_1 \times \cdots \times \mathcal{K}_K \qquad \mathcal{K}^* = \mathcal{K}_1^* \times \cdots \times \mathcal{K}_K^*$

$$P : \quad \min \quad \mathbf{c}'\mathbf{x} \qquad D : \quad \max \quad -\mathbf{b}'\mathbf{y}$$

$$\text{s.t.} \qquad \qquad \qquad \text{s.t.}$$

$$\mathbf{b} - \mathbf{A}\mathbf{x} \in \mathcal{K}^c \qquad \mathbf{x} \in \mathcal{K}^v \qquad \mathbf{c} + \mathbf{A}'\mathbf{y} \in \mathcal{K}^{v*} \qquad \mathbf{y} \in \mathcal{K}^{c*}$$


Some Standard Cones

- Free and zero cones: $\text{:Free} = \mathbb{R}^n$, $\text{:Zero} = \{\mathbf{0}\}$
- Orthant cones: $\text{:NonNeg} = \mathbb{R}_+^n$, $\text{:NonPos} = \mathbb{R}_-^n$
- Second-order (Lorentz) cone :SOC
 $= \{(t, \mathbf{v}) \in \mathbb{R} \times \mathbb{R}^{n-1} : \|\mathbf{v}\|_2 \leq t\}$
- Rotated second-order cone :SOCRotated
 $= \{(w, t, \mathbf{v}) \in \mathbb{R} \times \mathbb{R} \times \mathbb{R}^{n-2} : \|\mathbf{v}\|_2^2 \leq 2tw, t \geq 0, w \geq 0\}$
- Positive semidefinite cone $\text{:SDP} = \{\mathbf{V} \in \mathbb{S}^n : \mathbf{V} \succeq \mathbf{0}\}$
- Exponential cone :ExpPrimal
 $= \text{cl}\{(r, s, t) \in \mathbb{R}^3 : s > 0, s \exp(r/s) \leq t\}$
- Dual exponential cone :ExpDual
 $= \text{cl}\{(u, v, w) \in \mathbb{R}^3 : u < 0, w \geq 0, v \geq -u \log(-u/w) + u\}$



Cones not supported by conic solvers include the power cone for $\alpha \in (0, 1)$: $\{(r, s, t) \in \mathbb{R}^3 : |t| \leq r^\alpha s^{1-\alpha}, r \geq 0, s \geq 0\}$

Also Nonlinear Mixed Integer Programming (MIP)

- Example: Find minimum volume ellipsoid that contains 90% of data points




Minimum Volume Ellipsoid = SDP



Nonlinear MIP B&B Algorithms


- NLP (QCP) Based B&B
- (Dynamic) LP Based B&B
 - Few cuts = high speed.
 - Possible slow convergence.
- Lifted LP B&B
 - Extended or Lifted relaxation.
 - Static relaxation
 - Mimic NLP B&B.
 - Dynamic relaxation
 - Standard LP B&B

$$\begin{aligned} \max \quad & \sum_{i=1}^n c_i x_i \\ \text{s.t.} \quad & Ax + Dz \leq b, \\ & g_i(x) \leq 0, \quad i \in I, \quad x \in \mathbb{Z}^n \\ & x \in \mathbb{Z}^{n_1} \times \mathbb{R}^{n_2} \end{aligned}$$



LP v/s NLP B&B for CPLEX v11 for $n = 20$ and 30

- Results from V., Ahmed and Nemhauser 2008.



Dynamic Lifted LP for Separable Problems


- Motivating example from Hijazi et al. '14

$$F^n := \left\{ x \in \mathbb{R}^n : \sum_{i=1}^n \left(x_i - \frac{1}{2} \right)^2 \leq \frac{n-1}{4} \right\}$$

Extended formulation of F^n requires 2^n cuts.

$$\left(x_i - \frac{1}{2} \right)^2 \leq z_i \quad \forall i \in [n]$$

$$\sum_{i=1}^n z_i \leq \frac{n-1}{4}$$




$B^n \cap \mathbb{Z}^n = \emptyset$ with only $2n$ cuts

on extended formulation.


Lifted or Extended Approximations

- Projection = multiply constraints.
- V., A. and N. 2008:
 - Extremely accurate, but static and complex approximation by Ben-Tal and Nemirovski
- V., Dunning, Huchette and Lubin 2016: Simple, dynamic and good approximation:




$$\|y\|_2 \leq y_0 \quad \longrightarrow \quad \begin{aligned} & y_i^2 \leq z_i \cdot y_0 \quad \forall i \in [n] \\ & \sum_{i=1}^n z_i \leq y_0 \end{aligned}$$

CPLEX v12.6 for $n = 20, 30, 40, 50$ and 60



Gurobi v5.6.3 for $n = 20, 30, 40, 50$ and 60



All Major Solvers Now Implement Lifted LP

- First Talks:
 - SIAM Optimization (SIOPT), May 2014 \approx two weeks coding.
 - IBM Thomas J. Watson Research Center, December 2014.
- Paper in arxiv, May 28, 2015. 
 -  **CPLEX** v12.6.2, June 12, 2015.
 -  **GUROBI** v6.5, October 2015.
 -  **FICO** v8.0, May 2016.
 -  **SCIP** v4.0, March 2017.

However... We Can Still Beat CPLEX!


-  /  **JuMP**-based solver Pajarito

- Lubin, Yamangil, Bent and V. '16 and Coey, Lubin and V. '17.



solver	termination status counts				time(s)
	conv	wrong	not conv	limit	
SCIP	78	1	0	41	43.36
CPLEX	96	3	5	16	14.30
Paj-iter	96	1	0	23	38.70
Paj-MSD	101	0	0	19	18.12

Flexible Architecture Thanks to Julia-Opt Stack




- Fastest Open Source MISOCP Solver!
- Pajarito can also solve MISOCPs and MI-“EXP”

Nonlinear Mixed 0-1 Integer Formulations

- Modeling Finite Alternatives = Unions of Convex Sets

$$x \in \bigcup_{i=1}^n C_i \subseteq \mathbb{R}^d$$



Extended and Non-Extended Formulations for $\bigcup_{i=1}^n C_i$

$$C_i = \{x \in \mathbb{R}^d : f_i(x) \leq 0\}$$

Extended

$$\begin{aligned} \tilde{f}_i(x^i, y_i) &\leq 0 & \forall i \in [n] \\ \sum_{i=1}^n x^i &= x \\ \sum_{i=1}^n y_i &= 1 \\ y &\in \{0, 1\}^n \\ x, x^i &\in \mathbb{R}^d & \forall i \in [n] \end{aligned}$$

Strong, but large

Non-Extended

$$\begin{aligned} f_i(x) &\leq M_i(1 - y_i) & \forall i \in [n] \\ \sum_{i=1}^n y_i &= 1 \\ y &\in \{0, 1\}^n \\ x &\in \mathbb{R}^d & \forall i \in [n] \end{aligned}$$

Small, but weak?

Extended Formulations: Perspective “v/s” Cones

- e.g. Ceria and Soares '99

- e.g. Ben-tal and Nemirovski '01, Helton and Nie '09

$$C_i = \{x \in \mathbb{R}^d : f_i(x) \leq 0\}$$

$$\tilde{f}(x, y) = \begin{cases} yf(x/y) & \text{if } y > 0 \\ \lim_{\alpha \downarrow 0} \alpha f(x' - x + x/\alpha) & \text{if } y = 0 \\ +\infty & \text{if } y < 0 \end{cases}$$

$$\tilde{f}_i(x^i, y_i) \leq 0 \quad \forall i \in [n]$$

$$\sum_{i=1}^n x^i = x$$

$$\sum_{i=1}^n y_i = 1$$

$$y \in \{0, 1\}^n$$

$$x, x^i \in \mathbb{R}^d \quad \forall i \in [n]$$

$$C_i = \left\{ x \in \mathbb{R}^d : \begin{array}{l} \exists u \in \mathbb{R}^{p_i} \text{ s.t.} \\ A^i x + D^i u - b \in K^i \end{array} \right\}$$

K^i closed convex cone

$$A^i x^i + D^i u^i - b y_i \in K^i \quad \forall i \in [n]$$

$$\sum_{i=1}^n x^i = x$$

$$\sum_{i=1}^n y_i = 1$$

$$y \in \{0, 1\}^n$$

$$x, x^i \in \mathbb{R}^d \quad \forall i \in [n]$$

$$u^i \in \mathbb{R}^{p_i} \quad \forall i \in [n]$$

- Both formulations are **ideal** (extreme points of continuous relaxation satisfy integrality constraints)

Cones Can Mitigate Unintended Numerical Issues

- Let $C_i = \{x \in \mathbb{R}^2 : f_i(x) \leq 0\}$
where $f_i(x) = x_1^2 - x_2 - 1$

$$\tilde{f}_i(x, y) = \begin{cases} y(x_1/y)^2 - x_2 - y & \text{if } y > 0 \\ -x_2 & \text{if } y = x_1 = 0 \\ +\infty & \text{if o.w.} \end{cases}$$

- Conic (SOCP) representation

$$C_i = \left\{ x \in \mathbb{R}^2 : \sqrt{x_2^2 + 4x_1^2} \leq 2 + x_2 \right\}$$

$$\sqrt{(x_2^i)^2 + 4(x_1^i)^2} \leq 2y_i + x_2$$

Very Stable and Fast Conic Solvers

- Matrices grow quadratically so you can easily run out of memory.
- If it fits in memory you can probably solve relaxation

mosek

- And then the MIP



Advanced Convex MINLP Formulations

A Classical Strong Formulation for $\bigcup_{i=1}^k C_i$

$$C_i = \{x \in \mathbb{R}^n : A^i x \preceq_i b^i\}, \quad C_i^\infty = C_j^\infty$$

$$A^i x^i \preceq_i b^i z_i, \quad \forall i \in [k]$$

$$\sum_{i=1}^k x^i = x,$$

$$\sum_{i=1}^k z_i = 1, \quad z \in \{0, 1\}^k$$

$$x, x^i \in \mathbb{R}^n, \quad \forall i \in [k]$$

- Auxiliary continuous variables are copies of original variables
 - $y = (x^i)_{i=1}^k$
- “Ideal” Formulation Strength:
 - Extreme points of continuous relaxation satisfy **integrality constraints** on z
 - **Variable copies** crucial here, but **slow down computations** (usually worse than Big-M)

Generic Geometric Formulation = Gauge Functions

- For C such that $\mathbf{0} \in \text{int}(C)$ let:

$$\gamma_C(x) := \inf\{\lambda > 0 : x \in \lambda C\}$$

$$\text{epi}(\gamma_C) =$$

- If $b^i \in \text{int}(C_i)$ then **ideal** formulation:


$$\gamma_{C^i - \{b^i\}}(x^i - z_i b^i) \leq z_i \quad \forall i \in [k]$$

$$\sum_{i=1}^k x^i = x$$

$$\sum_{i=1}^k z_i = 1$$

$$z \in \{0, 1\}^k$$

$$x, x^i \in \mathbb{R}^n \quad \forall i \in [k]$$



Non-Extended and Big-M Formulations

- Gauge Big-M = no copies x^i of original variables:



$$\begin{aligned} \gamma_{C^i - \{b^i\}} \left(x - \sum_{j=1}^n y_j \overline{M}^{i,j} \right) &\leq \sum_{j=1}^n y_j \underline{M}^{i,j} \quad \forall i \in [n] \\ \sum_{i=1}^n y_i &= 1 \\ y &\in \{0, 1\}^n \end{aligned}$$

- Can be stronger than standard big-M
(e.g. $C_i = \{x \in \mathbb{R}^n : \|x\|_2 \leq r_i\}$ or $C_i = \{x \in \mathbb{R}^n : \|x\|_2^2 \leq r_i^2\}$)
- But may not be ideal
- What about **ideal non-extended** formulations?

Simple Non-Extended Ideal Formulation




- Unions of (nearly) Homothetic Closed Convex Sets:

$$C_i = \lambda_i C + b^i + C_\infty$$



$$\gamma_C \left(x - \sum_{i=1}^n y_i b^i \right) \leq \sum_{i=1}^n \lambda_i y_i$$
$$\sum_{i=1}^n y_i = 1, \quad y \in \{0, 1\}^n$$


Unary Encoding, Minkowski Sum and Cayley Trick



$$Q \cap (\mathbb{R}^2 \times \{0.5\}) \equiv P_1 + P_2 =$$

$$H = \{e^i\}_{i=1}^n$$

$$Q(H) \cap (\mathbb{R}^d \times \{\frac{1}{n} \sum_{i=1}^n \mathbf{e}^i\}) \equiv \sum_{i=1}^n P_i$$



Faces of Cayley Embedding


- Two types of facets (or faces):

- $P_1 \times \{0\} \equiv y_i \geq 0$


- $\text{conv}((F_1 \times 0) \cup (F_2 \times 1))$

F_i proper face of P_i

- Not all combinations of faces
 - Which ones are valid?





Valid Combinations = Common Normals






$$N(F_1) \cap N(F_2) \neq \emptyset$$




$\text{conv}((F_1 \times 0) \cup (F_2 \times 1))$
is face of $Q(H)$



Characterization Extends to Closed Convex Sets





Sticking Homothetic Formulations Together



Sufficient Conditions For Ideal Formulation

$$\sigma_S(u) := \sup\{u \cdot x : x \in S\}$$





$$\forall u \in \mathbb{R}^n \quad \exists j$$

s.t.


$$\sigma_{C_i}(u) = \sigma_{C_i^j}(u)$$

$$\forall i \in \{1, 2\}$$



*Similar to “lifting” of e.g.
Tawarmalani et al. ‘10*

May Need to “Find” Homothetic Constraints



$$x_1^2 \leq x_2 \leq 1$$




$$[-1, 1] \times 0$$


$$C_1 + (\mathbb{R}_+ \times \{0\}) :$$

$$(\max\{x_1, 0\})^2 \leq x_2 \leq 1$$

$$C_i + (\mathbb{R}_+ \times \{0\})$$





$$C_i + (\mathbb{R}_- \times \{0\})$$



Existing Small Ideal Formulations (Isotone Sets)

- Studied by Hijazi et al. '12 and Bonami et al. '15 ($n=1, 2$):
 - $C_i = \{x \in \mathbb{R}^d : l^i \leq x \leq u^i, \quad f_i(x) \leq 0\}$
- $f_i(x)$ component-wise monotonous ($i=1,2$ opposite).



- Ideal Formulation

$$y_1 l^1 + y_2 l^2 \leq x \leq y_1 u^1 + y_2 u^2$$

$$f_J^i(x, y) \leq 0$$

$$\forall J \subseteq [d], i \in [2]$$

$$y_1 + y_2 = 1$$

$$y_i \in \{0, 1\}$$

$$i \in [2]$$

Generalization and Simplification

- More than 2 sets (with general “opposite condition”).
- Generalization of the monotone/isotone condition (beyond affine transformation)
- Significantly smaller formulation: One non-linear constraint per set.

$$y_1 l^1 + y_2 l^2 \leq x \leq y_1 u^1 + y_2 u^2$$

~~$$f_J^i(x, y) \leq 0 \quad \forall J \subseteq [d], i \in [2]$$~~

$$y_1 + y_2 = 1$$

$$y_i \in \{0, 1\} \quad i \in [2]$$

$$\hat{f}^i(x, y) \leq 0 \quad \forall i \in [2]$$

Details of Size Reduction

$$C_i = \{x \in \mathbb{R}^d : l^i \leq x \leq u^i, \quad f_i(x) \leq 0\}$$

$$G_i = \{x \in \mathbb{R}^d : f_i(x) \leq 0\}$$

- Original formulation:


$$\gamma_{G_i}([x]_J) \leq y_i, \quad \boxed{\forall J \subseteq [d]} \quad ([x]_J)_j := \begin{cases} x_j & j \in J \\ 0 & o.w. \end{cases}$$

- Smaller formulation:

$$\gamma_{G_i}([x]^+) \leq y_i \quad ([x]^+)_j := \max\{x_j, 0\}$$

– max can cause representability issues.


Algebraic Representation Issues



$$x_1^2 \leq x_2 \leq 1$$



$$[-1, 1] \times 0$$



$$C_1 + (\mathbb{R}_+ \times \{0\}) : (\max\{x_1, 0\})^2 \leq x_2 \leq 1$$

- Non-basic semi-algebraic set contained in formulation.
- Finite polynomial inequalities requires max or auxiliary vars.

