



墨学教育
—MELBSTUDY—

FIT5047 Week 3

Knowledge representation

授课老师: Joe



FIT5047平时班 – Week 3

Propositional Logic



FIT5047平时班 – Week 3

Literal: a proposition or its negation

- E.g., P , $\neg P$

Clause: a disjunction of literals

- E.g., $\neg P \vee Q \vee A$

Negation: If S is a sentence, $\neg S$ is a sentence

Conjunction:

If S_1 and S_2 are sentences, $S_1 \wedge S_2$ is a sentence

Disjunction:

If S_1 and S_2 are sentences, $S_1 \vee S_2$ is a sentence

Implication:

If S_1 and S_2 are sentences, $S_1 \Rightarrow S_2$ is a sentence

Biconditional:

If S_1 and S_2 are sentences, $S_1 \Leftrightarrow S_2$ is a sentence



FIT5047平时班 – Week 3

$\neg S$ is true iff S is false
 $S_1 \wedge S_2$ is true iff S_1 is true and S_2 is true
 $S_1 \vee S_2$ is true iff S_1 is true or S_2 is true
 $S_1 \Rightarrow S_2 \equiv \neg S_1 \vee S_2$ is true iff S_1 is false or S_2 is true
 $S_1 \Leftrightarrow S_2$ is true iff $S_1 \Rightarrow S_2$ is true and $S_2 \Rightarrow S_1$ is true

P	Q	$\neg P$	$P \wedge Q$	$P \vee Q$	$P \Rightarrow Q$	$P \Leftrightarrow Q$
false	false	true	false	false	true	true
false	true	true	false	true	true	false
true	false	false	false	true	false	false
true	true	false	true	true	true	true



FIT5047平时班 – Week 3

- Two sentences are **logically equivalent** iff they are both true in the same models: $\alpha \equiv \beta$ iff $\alpha \models \beta$ and $\beta \models \alpha$

$$(\alpha \wedge \beta) \equiv (\beta \wedge \alpha) \quad \text{commutativity of } \wedge$$

$$(\alpha \vee \beta) \equiv (\beta \vee \alpha) \quad \text{commutativity of } \vee$$

$$((\alpha \wedge \beta) \wedge \gamma) \equiv (\alpha \wedge (\beta \wedge \gamma)) \quad \text{associativity of } \wedge$$

$$((\alpha \vee \beta) \vee \gamma) \equiv (\alpha \vee (\beta \vee \gamma)) \quad \text{associativity of } \vee$$

$$\neg(\neg\alpha) \equiv \alpha \quad \text{double-negation elimination}$$

$$(\alpha \Rightarrow \beta) \equiv (\neg\beta \Rightarrow \neg\alpha) \quad \text{contraposition}$$

$$(\alpha \Rightarrow \beta) \equiv (\neg\alpha \vee \beta) \quad \text{implication elimination}$$

$$(\alpha \Leftrightarrow \beta) \equiv ((\alpha \Rightarrow \beta) \wedge (\beta \Rightarrow \alpha)) \quad \text{biconditional elimination}$$

$$\neg(\alpha \wedge \beta) \equiv (\neg\alpha \vee \neg\beta) \quad \text{de Morgan}$$

$$\neg(\alpha \vee \beta) \equiv (\neg\alpha \wedge \neg\beta) \quad \text{de Morgan}$$

$$(\alpha \wedge (\beta \vee \gamma)) \equiv ((\alpha \wedge \beta) \vee (\alpha \wedge \gamma)) \quad \text{distributivity of } \wedge \text{ over } \vee$$

$$(\alpha \vee (\beta \wedge \gamma)) \equiv ((\alpha \vee \beta) \wedge (\alpha \vee \gamma)) \quad \text{distributivity of } \vee \text{ over } \wedge$$



FIT5047平时班 – Week 3

- **A sentence is valid if it is true in all models**
 - E.g., *True*, $A \vee \neg A$, $A \Rightarrow A$
- **Validity is connected to inference via the Deduction Theorem**
 - $\alpha \models \beta$ iff $\alpha \Rightarrow \beta$ is valid
- **A sentence is satisfiable if it is true in some model**
 - E.g., $A \vee B$, C
- **A sentence is unsatisfiable if it is true in no model**
 - E.g., $A \wedge \neg A$
- **Satisfiability and validity are connected**
 - α is valid iff $\neg \alpha$ is unsatisfiable
 - α is satisfiable iff $\neg \alpha$ is not valid
 - $\alpha \models \beta$ iff $\alpha \wedge \neg \beta$ is unsatisfiable



FIT5047平时班 – Week 3

Prove that $(A \wedge (A \Rightarrow B)) \Rightarrow B$ is valid

- $(A \wedge (\neg A \vee B)) \Rightarrow B$
- $((A \wedge \neg A) \vee (A \wedge B)) \Rightarrow B$
- $(A \wedge B) \Rightarrow B$
- $\neg(A \wedge B) \vee B$
- $\neg A \vee \neg B \vee B$
- True



- **Prove that α is valid iff $\neg\alpha$ is unsatisfiable**
 - If α is valid, it is true in all models \rightarrow there does not exist a model for which $\neg\alpha$ is true $\rightarrow \neg\alpha$ is unsatisfiable
 - If $\neg\alpha$ is unsatisfiable \rightarrow there does not exist a model for which $\neg\alpha$ is true $\rightarrow \alpha$ is true in all models $\rightarrow \alpha$ is valid
- **Prove that α is satisfiable iff $\neg\alpha$ is not valid**
 - If α is satisfiable, it is true in some models $\rightarrow \neg\alpha$ is not true in these models $\rightarrow \neg\alpha$ is not valid
 - If $\neg\alpha$ is not valid \rightarrow there exist some models for which $\neg\alpha$ is false $\rightarrow \alpha$ is true in these models $\rightarrow \alpha$ is satisfiable



Resolution

1. $\text{Eat} \Rightarrow \text{HaveLessMoney}$

2. $\neg \text{Eat} \Rightarrow \text{Hungry}$

Resolvent: $\text{HaveLessMoney} \vee \text{Hungry}$



Conversion to conjunctive normal form

1. Eliminate \Leftrightarrow , replacing $\alpha \Leftrightarrow \beta$ with $(\alpha \Rightarrow \beta) \wedge (\beta \Rightarrow \alpha)$
2. Eliminate \Rightarrow , replacing $\alpha \Rightarrow \beta$ with $\neg\alpha \vee \beta$
3. Move \neg inwards by repeated application of the following equivalences:
 - > double-negation: $\neg(\neg\alpha) \equiv \alpha$
 - > de Morgan $\neg(\alpha \wedge \beta) \equiv \neg\alpha \vee \neg\beta$
 - > de Morgan $\neg(\alpha \vee \beta) \equiv \neg\alpha \wedge \neg\beta$
4. Apply distributivity law (\wedge over \vee) and flatten



FIT5047平时班 – Week 3

$A \Leftrightarrow (B \vee C)$

1. Eliminate \Leftrightarrow : $(A \Rightarrow (B \vee C)) \wedge ((B \vee C) \Rightarrow A)$
2. Eliminate \Rightarrow : $(\neg A \vee (B \vee C)) \wedge (\neg (B \vee C) \vee A)$
3. Move \neg inwards: $(\neg A \vee (B \vee C)) \wedge ((\neg B \wedge \neg C) \vee A)$
 $(\neg A \vee B \vee C) \wedge ((\neg B \wedge \neg C) \vee A)$
4. Apply distributivity law:
 $(\neg A \vee B \vee C) \wedge (\neg B \vee A) \wedge (\neg C \vee A)$



- **Proof by refutation**

1. Negate the goal and add the negation to the set of clauses
2. Apply resolution to the clauses in the set of clauses until a contradiction is reached


- **Answer extraction**

1. Build a tautology by appending the goal itself to the negation of the goal
2. When the negated goal is contradicted, the answer resides in the goal



FIT5047平时班 – Week 3

- $KB = (A \Leftrightarrow (B \vee C)) \wedge \neg A$
 $(\neg A \vee B \vee C) \wedge (\neg B \vee A) \wedge (\neg C \vee A) \wedge \neg A$
- Prove: $\alpha = \neg B$





- **KB:**

- R1: $\neg P \vee Q$
- R2: $\neg L \vee \neg M \vee P$
- R3: $\neg B \vee \neg L \vee M$
- R4: $\neg A \vee \neg P \vee L$
- R5: $\neg A \vee \neg B \vee L$
- A
- B

- Negate Q: $\neg Q$

$$\begin{array}{l} \underline{\neg Q} \quad R1: \neg P \vee \underline{Q} \\ \quad \underline{\neg P} \\ R2: \neg L \vee \neg M \vee \underline{P} \\ \quad \neg L \vee \underline{\neg M} \\ R3: \neg B \vee \neg L \vee \underline{M} \\ \quad \neg B \vee \underline{\neg L} \\ R5: \neg A \vee \neg B \vee \underline{L} \\ \quad \underline{\neg A} \vee \neg B \\ \underline{A} \\ \quad \underline{\neg B} \\ \underline{B} \\ \quad \text{nil} \end{array}$$



FIT5047平时班 – Week 3

Definite clause – a disjunction of literals of which exactly one is positive

$$\neg A \vee \neg B \vee C$$

Horn clause – a disjunction of literals of which at most one is positive




FIT5047平时班 – Week 3

- **KB:**

- R1: $P \Rightarrow Q$
- R2: $L \wedge M \Rightarrow P$
- R3: $B \wedge L \Rightarrow M$
- R4: $A \wedge P \Rightarrow L$
- R5: $A \wedge B \Rightarrow L$
- A
- B

- **Prove Q**



Agenda	Count					Inferred
	R1	R2	R3	R4	R5	
AB	1	2	2	2	2	


L
M



FIT5047平时班 – Week 3

- **KB:**

- R1: $P \Rightarrow Q$
- R2: $L \wedge M \Rightarrow P$
- R3: $B \wedge L \Rightarrow M$
- R4: $A \wedge P \Rightarrow L$
- R5: $A \wedge B \Rightarrow L$
- A
- B



- **Prove Q**



First order logic



FIT5047平时班 – Week 3

Notation

Predicate symbol – MARRIED
Constant – *John* or *A*
Function – *Mother* or *f*
Variable – *x*

Predicate – a function that only returns Boolean

Function can return any value



FIT5047平时班 – Week 3

- disjunction – $P(x) \vee Q(y) \vee W(x,y)$
- conjunction – $P(x) \wedge Q(y)$
- implication – $P(x) \Rightarrow W(x,y) \equiv \neg P(x) \vee W(x,y)$



FIT5047平时班 – Week 3

Quantification

- Universal (\forall) –
 $\forall x [\text{ELEPHANT}(x) \Rightarrow \text{COLOUR}(x, \text{Gray})]$
- Existential (\exists) –
 $\exists x \text{ WRITE}(x, \text{Computer-Chess})$

predicate

constant

variable



$$\exists x(\forall y[P(x, y) \wedge Q(x, y)] \Rightarrow R(x))$$

$$\neg f(A)$$

$$\neg P(A, g(A, B, A))$$

$$f(P(A))$$

$$\forall P P(A)$$



FIT5047平时班 – Week 3

- **Existential inside the scope of universal**
E.g., $\forall s \exists c \text{ Eats}(s, c)$
- **Universal inside the scope of existential**
E.g., $\exists c \forall s \text{ Eats}(s, c)$



FIT5047平时班 – Week 3

- $\neg(\exists x)P(x) \equiv (\forall x) [\neg P(x)]$
There does not exist an x such that $P(x)$ is true \equiv
For all x , $P(x)$ is false
- $\neg(\forall x)P(x) \equiv (\exists x) [\neg P(x)]$
It is not true that for all x $P(x)$ is true \equiv
There exists an x , such that $\neg P(x)$
- $(\forall x)[P(x) \wedge Q(x)] \equiv (\forall x)P(x) \wedge (\forall y)Q(y)$
For all x , $P(x)$ and $Q(x)$ are true \equiv
For all x , $P(x)$ is true, and for all y $Q(y)$ is true
- $(\exists x)[P(x) \vee Q(x)] \equiv (\exists x)P(x) \vee (\exists y)Q(y)$
There is an x , such that $P(x)$ is true or $Q(x)$ is true \equiv
There is an x , such that $P(x)$ is true, or
there is a y , such that $Q(y)$ is true

$$(\forall x)[P(x) \vee Q(x)]$$

$$(\forall x)P(x) \vee (\forall y)Q(y)$$



Substitution

Substitution is a set of ordered pairs

$s = \{v_1|t_1, v_2|t_2, \dots, v_n|t_n\}$

where $v_i|t_i$ means that term t_i substitutes variable v_i throughout


– Example:

$P(x, y) \{x|A, y|B\} \rightarrow P(A, B)$

**Semantics: all elements are applied
*simultaneously***

– Example:

$P(w, y, g(z), x) \{x|g(y), y|h(z), z|x\} \rightarrow P(w, h(z), g(x), g(y))$





Substitution

$s_i s_j$ – composition of two substitutions

- Apply s_j to the terms of s_i
- Add any pairs of s_j having variables not in s_i

Example:

$s_1 = \{z|g(x,y)\}$ $s_2 = \{x|A, y|B, w|C, z|D\}$

$s_1 s_2 = \{z|g(x,y)\}\{x|A, y|B, w|C, z|D\} = \{z|g(A,B), x|A, y|B, w|C\}$

$s_2 s_1 = \{x|A, y|B, w|C, z|D\} \{z|g(x,y)\} = \{x|A, y|B, w|C, z|D\}$

NOT commutative: $s_1 s_2 \neq s_2 s_1$



Unification

$s = \{x|A, y|B\}$ unifies $P(x, f(y))$ with $P(x, f(B))$

You can substitute anything into variable,
not the other way around



Converting wffs into clauses

1. Eliminate implication symbols
2. Reduce scopes of negation symbols
3. Standardize variables (for each quantifier)
4. Eliminate existential quantifiers (*skolemize*)
5. Move all universal quantifiers to the front
6. Put result in conjunctive normal form (CNF)
7. Eliminate universal quantifiers
8. Eliminate \wedge symbols
9. Rename variables (standardize variables apart for each clause)



FIT5047平时班 – Week 3

$\forall x [\text{CanRead}(x) \Rightarrow \text{Intelligent}(x)]$

1. Eliminate \Rightarrow : $\forall x [\neg \text{CanRead}(x) \vee \text{Intelligent}(x)]$

7. Eliminate \forall : $\neg \text{CanRead}(x) \vee \text{Intelligent}(x)$



FIT5047平时班 – Week 3

$\forall x [\neg (\forall y) [P(x,y) \Rightarrow Q(x,y)]]$

1. Eliminate \Rightarrow : $\forall x [\neg (\forall y) [\neg P(x,y) \vee Q(x,y)]]$
2. Reduce scope of \neg : $\forall x [\exists y \neg [\neg P(x,y) \vee Q(x,y)]]$
 $\forall x \exists y [P(x,y) \wedge \neg Q(x,y)]$
4. Eliminate \exists : $\forall x [P(x,g(x)) \wedge \neg Q(x,g(x))]$
7. Eliminate \forall : $P(x,g(x)) \wedge \neg Q(x,g(x))$
8. Eliminate \wedge symbols: $\{ P(x,g(x)), \neg Q(x,g(x)) \}$
9. Standardize variables apart: $\{ P(x_1,g(x_1)), \neg Q(x_2,g(x_2)) \}$



FIT5047平时班 – Week 3

- 1. If a unit is easy, there are some students who are enrolled in it who are happy**

$$\forall u [\text{Easy}(u) \Rightarrow \exists s [\text{Enrolled}(s,u) \wedge \text{Happy}(s)]]$$

- 2. If a unit has a final exam, no students that are enrolled in it are happy**

$$\forall u [\text{HasFinal}(u) \Rightarrow \neg \exists s [\text{Enrolled}(s,u) \wedge \text{Happy}(s)]]$$

- 3. Prove that if a unit has a final exam, the unit is not easy**

$$\forall u [\text{HasFinal}(u) \Rightarrow \neg \text{Easy}(u)]$$



FIT5047平时班 – Week 3

1. $\forall u [\text{Easy}(u) \Rightarrow \exists s [\text{Enrolled}(s,u) \wedge \text{Happy}(s)]]$

Eliminate \Rightarrow : $\forall u [\neg \text{Easy}(u) \vee \exists s [\text{Enrolled}(s,u) \wedge \text{Happy}(s)]]$

Eliminate \exists : $\forall u [\neg \text{Easy}(u) \vee [\text{Enrolled}(g(u),u) \wedge \text{Happy}(g(u))]]$

Convert to CNF: $\forall u [[\neg \text{Easy}(u) \vee \text{Enrolled}(g(u),u)] \wedge [\neg \text{Easy}(u) \vee \text{Happy}(g(u))]]$

Eliminate \forall : $[\neg \text{Easy}(u) \vee \text{Enrolled}(g(u),u)] \wedge [\neg \text{Easy}(u) \vee \text{Happy}(g(u))]$

Eliminate \wedge and standardize variables apart:

1.1 $\neg \text{Easy}(u_1) \vee \text{Enrolled}(g(u_1),u_1)$

1.2 $\neg \text{Easy}(u_2) \vee \text{Happy}(g(u_2))$



FIT5047平时班 – Week 3

2. $\forall u [\text{HasFinal}(u) \Rightarrow \neg \exists s [\text{Enrolled}(s,u) \wedge \text{Happy}(s)]]$

Eliminate \Rightarrow : $\forall u [\neg \text{HasFinal}(u) \vee$
 $\neg \exists s [\text{Enrolled}(s,u) \wedge \text{Happy}(s)]]$

Reduce scope of \neg :

$\forall u [\neg \text{HasFinal}(u) \vee \forall s \neg [\text{Enrolled}(s,u) \wedge \text{Happy}(s)]]$
 $\forall u [\neg \text{HasFinal}(u) \vee \forall s [\neg \text{Enrolled}(s,u) \vee \neg \text{Happy}(s)]]$

Move \forall to the front: $\forall u \forall s [\neg \text{HasFinal}(u) \vee$
 $\neg \text{Enrolled}(s,u) \vee \neg \text{Happy}(s)]$

Eliminate \forall and standardize variables apart:

$\neg \text{HasFinal}(u_3) \vee \neg \text{Enrolled}(s_3, u_3) \vee \neg \text{Happy}(s_3)$



Using resolution to prove statement 3

Negate the goal:

3'. $\neg \forall u [\text{HasFinal}(u) \Rightarrow \neg \text{Easy}(u)]$

Eliminate \Rightarrow : $\neg \forall u [\neg \text{HasFinal}(u) \vee \neg \text{Easy}(u)]$

Reduce scope of \neg : $\exists u [\text{HasFinal}(u) \wedge \text{Easy}(u)]$


Eliminate \exists : $\text{HasFinal}(A) \wedge \text{Easy}(A)$

Eliminate \wedge : 3'.1 $\text{HasFinal}(A)$

3'.2 $\text{Easy}(A)$



FIT5047平时班 – Week 3





FIT5047平时班 – Week 3

Question answering

1. MANAGER(Purchasing-dept., John-Jones)
2. WORKSIN(Purchasing-dept., Joe-Smith)
3. $\forall x \forall y \forall z [\text{WORKSIN}(x,y) \wedge \text{MANAGER}(x,z)] \Rightarrow \text{BOSSOF}(y,z)$
4. **Goal:** Who is the boss of Joe Smith?
 $\exists x \text{BOSSOF}(\text{Joe-Smith}, x)$



FIT5047平时班 – Week 3

