

Developing an Antimicrobial Strategy for Sepsis in Malawi

-

Thesis submitted in accordance with the requirements of the Liverpool School of Tropical Medicine for the degree of Doctor in Philosophy by Joseph Michael Lewis

August 2019

Contents

Preface	9
1 Introduction	11
1.1 Chapter Overview	13
1.2 Sepsis in sub-Saharan Africa	13
1.3 ESBL-E in sub-Saharan Africa	13
1.4 Conclusions	13
1.5 Thesis overview	13
1.6 Appendix	13
1.7 References	13
2 Methods	15
2.1 Chapter Overview	17
2.2 Study site	17
2.3 Clinical Study	17
2.4 Diagnostic Laboratory Procedures	17
2.5 Molecular methods	17
2.6 Bioinformatics	17
2.7 Statistical Analysis	17
2.8 Study Team	17
2.9 Data Collection and Storage	17
2.10 Ethical Approval, Consent and Participant Remuneration	17
3 <i>Mycobacterium tuberculosis</i> BSI: an IPD meta analysis	19
4 A clinical and microbiological description of sepsis in Blantyre, Malawi	21
4.1 Chapter overview	22
4.2 Introduction and chapter aims	22
4.3 Aims and Methods	22

4.4	Results	22
4.5	Discussion	22
4.6	Conclusions and further work	22
4.7	Appendix	22
5	Longitudinal ESBL-E carriage in Malawian adults in health and disease	23
5.1	Chapter Overview	23
5.2	Introduction and chapter aims	23
5.3	Methods	23
5.4	Results	23
5.5	Discussion	23
5.6	Conclusions and further work	23
6	Whole genome sequencing of ESBL <i>E. coli</i> carriage isolates	25
6.1	Chapter overview	27
6.2	Methods	27
6.3	Results	27
6.4	Discussion	27
6.5	Appendix	27
7	Gut mucosal carriage of ESBL-E in Blantyre, Malawi	29
	References	31

List of Tables

List of Figures

Preface

Placeholder

Chapter 1

Introduction

Placeholder

1.1 Chapter Overview

1.2 Sepsis in sub-Saharan Africa

1.2.1 Search strategy

1.2.2 Defining sepsis

1.2.3 Applicability of sepsis-3 definitions in sub-Saharan Africa

1.2.4 Sepsis epidemiology in sub-Saharan Africa

1.2.4.1 Incidence

1.2.4.2 Risk factors: the sepsis population in sub-Saharan Africa

1.2.4.3 Outcomes

1.2.5 Sepsis aetiology in sub-Saharan Africa

1.2.5.1 Bacterial zoonoses, Rickettsioses and arboviruses

1.2.5.2 HIV opportunistic infections: PCP, histoplasmosis and cryptococcal disease

1.2.6 Sepsis management

1.2.6.1 Early goal directed therapy

1.2.6.2 Evidence to guide antimicrobial therapy in sSA

1.2.6.3 Evidence to guide intravenous fluid therapy in sub-Saharan Africa

1.3 ESBL-E in sub-Saharan Africa

1.3.1 Search strategy

1.3.2 Introduction: definition and classification of ESBL-E

1.3.3 Global molecular epidemiology of ESBL-E: an overview

1.3.3.1 1980s-1990s: First identification of ESBL in nosocomial pathogens

1.3.3.2 1990s-2010s: Emergence and globalisation of CTX-M

Chapter 2

Methods

Placeholder

2.1 Chapter Overview

2.2 Study site

2.2.1 Malawi

2.2.2 Queen Elizabeth Central Hospital

2.2.3 Participating Laboratories

2.2.3.1 Malawi-Liverpool-Wellcome Clinical Research Programme

2.2.3.2 Malawi College of Medicine Tuberculosis Laboratory

2.2.3.3 Wellcome Trust Sanger Institute

2.3 Clinical Study

2.3.1 Entry Criteria

2.3.2 Study Visits and Patient Sampling

2.3.2.1 Enrollment assessment and first six hours

2.3.2.2 Subsequent visits

2.3.2.3 Blood, urine, and stool, sputum and CSF collection

2.3.2.4 Imaging: chest x-ray and ultrasound scanning

2.3.3 Outcomes and sample size calculations

2.4 Diagnostic Laboratory Procedures

2.4.1 Point of care diagnostics

2.4.2 Laboratory diagnostics

2.4.2.1 Haematology and biochemistry

2.4.2.2 Aerobic blood and CSF culture

2.4.2.3 Mycobacterial blood culture

2.4.2.4 Sputum Xpert

Chapter 3

Mycobacterium tuberculosis BSI: an IPD meta analysis

Chapter 4

A clinical and microbiological description of sepsis in Blantyre, Malawi

Placeholder

4.1 Chapter overview

4.2 Introduction and chapter aims

4.3 Aims and Methods

4.4 Results

4.4.1 Study population

4.4.2 Symptoms and health-seeking behaviour

4.4.3 Admission physiology and laboratory investigations

4.4.4 Aetiology

4.4.5 Treatment

4.4.6 Outcome

4.4.7 Determinants of mortality

4.5 Discussion

4.5.1 Demographics and outcome: significant longer-term mortality

4.5.2 Aetiology: TB dominates as a cause of sepsis

4.5.3 Determinants of 28-day mortality: an expanded role for TB therapy?

4.5.4 Limitations

4.6 Conclusions and further work

4.7 Appendix

Chapter 5

Longitudinal ESBL-E carriage in Malawian adults in health and disease

Placeholder

5.1 Chapter Overview

5.2 Introduction and chapter aims

5.3 Methods

5.4 Results

5.4.1 Study population

5.5 Discussion

5.5.1 Limitations

5.6 Conclusions and further work

Chapter 6

Whole genome sequencing of ESBL *E. coli* carriage isolates

Placeholder

6.1 Chapter overview

6.2 Methods

6.2.1 Bioinformatic pipeline

6.2.2 Global *E. coli* collection

6.2.3 Statistical analysis

6.3 Results

6.3.1 Samples and quality control

6.3.2 Phylogroup, MLST and core genome phylogeny of study isolates

6.3.3 Study isolates in a global context

6.3.4 Antimicrobial resistance determinants

6.3.4.1 β -lactam resistance

6.3.4.2 Quinolone resistance

6.3.4.3 Aminoglycoside resistance

6.3.4.4 Chloramphenicol, co-trimoxazole, tetracycline and other resistance determinants

6.3.4.5 Clustering and lineage association of AMR determinants

6.3.5 Plasmid replicons

6.3.6 Testing metadata associations: SNP distance, hierBAPS sequence clusters and ESBL-clusters

6.3.6.1 Hierarchical BAPS clustering of core gene pseudosequences

6.3.6.2 ESBL-clusters

6.3.6.3 Assessing for healthcare-associated lineages

6.3.6.4 Assessing for within-patient conservation of lineage or MGE

6.4 Discussion

Chapter 7

Gut mucosal carriage of ESBL-E in Blantyre, Malawi

References