

[illegible]

Pest Name	Symptoms	Organic	Inorganic	Homemade
Parlatoria zizyphus Lucus	Parlatoria zizyphus Lucus infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Phyllocnistis citrella Stainton	Phyllocnistis citrella Stainton infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Phyllocoptes oleiverus ashmead	Phyllocoptes oleiverus ashmead infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Pieris canidia	Pieris canidia infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Polyphagotars onemus latus	Polyphagotars onemus latus infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Potosiabre vitarsis	Potosiabre vitarsis infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Prodenia litura	Prodenia litura infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Pseudococcus comstocki Kuwana	Pseudococcus comstocki Kuwana infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Rhytidodera bowrinii white	Rhytidodera bowrinii white infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Salurnis marginella Guerr	Salurnis marginella Guerr infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.