

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

Found 135 items with missing information.

Adristyrannus

- Missing Usage Instructions for: Organic: Neem oil spray (3-5 ml per liter of water) (No usage info), Organic: NSKE, pheromone traps (No usage info), Chemical: Chlorantraniliprole (No usage info), Chemical: Emamectin benzoate (No usage info)
-

Aleurocanthus_spiniferus

- Missing Usage Instructions for: Organic: Neem oil (3-5 ml/L) or NSKE 5% (No usage info), Organic: Encourage natural enemies (e.g., Encarsia spp.) (No usage info), Organic: Yellow sticky traps (No usage info), Chemical: Imidacloprid / Thiamethoxam (No usage info), Chemical: Buprofezin for nymph control (No usage info)
-

Ampelophaga

- Missing Usage Instructions for: Organic: Spinosad (Derived from soil bacteria) (No usage info), Chemical: Carbaryl (No usage info), Chemical: Chlorpyrifos (No usage info)
-

Aphis_citricola_Vander_Goot

- Missing Usage Instructions for: Organic: Insecticidal Soap (Potassium salts) (No usage info), Organic: Beauveria bassiana (Fungal bio-pesticide) (No usage info), Chemical: Imidacloprid (No usage info), Chemical: Dimethoate (No usage info)
-

Apolygus_lucorum

- Missing Usage Instructions for: Organic: Matrine (0.3%): Spray early morning | 2-3 ml per liter (No usage info), Organic: Beauveria bassiana: Apply during humidity | 3-5 g per liter (No usage info), Organic: Pyrethrum: Mist at dusk | 2-4 ml per liter (No usage info), Chemical: Imidacloprid (17.8 SL): Spray pre-bloom | 0.5 ml per liter (No usage info), Chemical: Acetamiprid (20 SP): Apply on foliage | 0.5 g per liter (No usage info), Chemical: Lambda-cyhalothrin (5 EC): Use during outbreaks | 1 ml per liter (No usage info)
-

Apple_Apple_scab

- Missing Usage Instructions for: Organic: Wettable Sulfur: Spray pre-rain | 3-5 g per liter Copper Fungicide: Apply dormant stage | 2-3 g per liter (No usage info), Chemical: Mancozeb (75 WP): Spray preventative | 2.5 g per liter (No usage info), Homemade: Baking Soda Mix: Mist thoroughly | 5 g per liter (+ few drops oil) (No usage info)
-

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

Apple_Black_rot

- Missing Usage Instructions for: Organic: Pruning & Sanitation : Cut out all cankers and dead wood during the dormant season (winter). Prune at least 6-8 inches below the visible edge of the infection. (No usage info), Chemical: Captan or Myclobutanil: (No usage info), Homemade: Vinegar Spray (Acidification): (No usage info)

Apple_Cedar_apple_rust

- Missing Usage Instructions for: Organic: Sulfur sprays. (No usage info), Organic: Remove nearby galls on cedar trees. (No usage info), Chemical: Myclobutanil (No usage info), Chemical: Mancozeb (No usage info), Homemade: Garlic water spray. (No usage info)

Apple_healthy

- Missing Symptoms
- Missing Organic Treatments
- Missing Chemical (Inorganic) Treatments
- Missing Homemade Treatments

Background_without_leaves

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info), Homemade: Soap or botanical spray: Prepare mild soap or plant extract solution and spray directly on infested areas. (No usage info)

Bactrocera_tsuneonis

- Missing Usage Instructions for: Organic: OrganicX (No usage info), Chemical: TestInorganic1 (No usage info), Homemade: TestHomemade1 (No usage info)

Beet_spot_flies

- Missing Usage Instructions for: Organic: Floating Row Covers: Application: This is the best prevention. Cover your crops with a lightweight garden fabric immediately after planting. This prevents the adult flies from landing and laying eggs on the leaves. (No usage info), Chemical: Pyrethroid Sprays (e.g., Permethrin): Usage: Contact insecticide. Application: Must be timed perfectly to hit the adult flies or the larvae before they enter the leaf. Once they are inside the "mine," these chemicals are largely ineffective. (No usage info), Homemade: Neem & Soap Wash: Recipe: 1 tsp Neem oil, 1/2 tsp liquid dish soap, and 1 liter of water. Application: Spray the undersides of leaves every 7 days. While it won't

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

always kill larvae inside the leaf, it acts as a strong repellent for the adult flies looking to lay egg (No usage info)

Blueberry_healthy

- Missing Symptoms
 - Missing Organic Treatments
 - Missing Chemical (Inorganic) Treatments
 - Missing Homemade Treatments
-

Brevipalpus_lewisi_McGregor

- Missing Usage Instructions for: Organic: Horticultural Oils: Usage: 1% to 2% oil emulsion. Application: Provides excellent control by suffocating the mites and their eggs. Ensure "thorough coverage," meaning the spray should be dripping off the undersides of leaves and fruit. (No usage info), Chemical: Spirodiclofen: Usage: An insect growth regulator that prevents mites from molting and laying viable eggs. (No usage info), Homemade: Canola Oil Mixture: Recipe: 1 cup Canola oil mixed with 1 tbsp dish soap. Use 2 tsp of this concentrate per 1 liter of water. Application: Acts as a homemade "smothering" oil. Effective for small-scale infestations on garden citrus. (No usage info)

Ceroplastes_rubens

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)

Cherry_Powdery_mildew

- Missing Usage Instructions for: Organic: Sulfur fungicide (No usage info), Chemical: Myclobutanil (No usage info), Homemade: Milk spray (No usage info)

Cherry_healthy

- Missing Symptoms
 - Missing Organic Treatments
 - Missing Chemical (Inorganic) Treatments
 - Missing Homemade Treatments
 - Missing Usage Instructions for: Entire record missing
-

Chlumetia_transversa

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)
-

Cicadella_viridis

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)
-

Colomerus_vitis

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)
-

Corn_Cercospora_leaf_spot_Gray_leaf_spot

- Missing Usage Instructions for: Organic: Neem oil / biofungicide (No usage info), Chemical: Azoxystrobin (No usage info), Homemade: Crop residue removal (No usage info)
-

Corn_Common_rust

- Missing Usage Instructions for: Organic: Neem oil (No usage info), Chemical: Propiconazole (No usage info), Homemade: Garlic extract (No usage info)
-

Corn_Northern_Leaf_Blight

- Missing Usage Instructions for: Organic: Trichoderma (No usage info), Chemical: Chlorothalonil (No usage info), Homemade: Neem extract (No usage info)
-

Corn_healthy

- Missing Symptoms
 - Missing Organic Treatments
 - Missing Chemical (Inorganic) Treatments
 - Missing Homemade Treatments
 - Missing Usage Instructions for: Entire record missing
-

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

Dacus_dorsalisHendel

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)

Dasineura_sp

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)

Deporaus_marginatus_Pascoe

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)

Erythroneura_apicalis

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)

Grape_Black_rot

- Missing Usage Instructions for: Organic: Copper spray. (No usage info), Organic: Remove mummies. (No usage info), Chemical: Mancozeb (No usage info), Chemical: Myclobutanil (No usage info), Homemade: Baking soda mix. (No usage info)

Grape_Esca_(Black_Measles)

- Missing Usage Instructions for: Organic: Prune and remove infected vines. (No usage info), Organic: Wound protection. (No usage info), Chemical: No effective chemical cure (No usage info), Chemical: focus on prevention. (No usage info), Homemade: None reliable. (No usage info)

Grape_Leaf_blight_(Isariopsis_Leaf_Spot)

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

- Missing Usage Instructions for: Organic: Sulfur or Copper fungicides. (No usage info), Chemical: Mancozeb (No usage info), Homemade: Neem oil. (No usage info)
-

Grape_healthy

- Missing Symptoms
 - Missing Organic Treatments
 - Missing Chemical (Inorganic) Treatments
 - Missing Homemade Treatments
-

Icerya_purchasi_Maskell

- Missing Usage Instructions for: Organic: Horticultural Oil: Spray dormant stage | 20-30 ml per liter (No usage info), Chemical: Malathion (50 EC): Apply outbreaks | 2 ml per liter (No usage info), Homemade: Alcohol Swab: Wipe visible scales | Use undiluted (70% isopropyl) (No usage info)
-

Lawana_imitata_Melichar

- Missing Usage Instructions for: Organic: Metarhizium anisopliae: Spray evenings | 5 g per liter (No usage info), Chemical: Dimethoate (30 EC): Spray infested stems | 2 ml per liter (No usage info), Homemade: Garlic-Chili Extract: Spray deterrent | 50 ml per liter (No usage info)
-

Locustoidea

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)
-

Lycorma_delicatula

- Missing Usage Instructions for: Organic: Spinosad: Spray contact only | 1-2 ml per liter (No usage info), Chemical: Dinotefuran: Bark spray/drench | Follow label (approx 3g/L) (No usage info), Homemade: Vinegar Spray: Mist on weeds | 250 ml vinegar per liter (careful on crops) (No usage info)
-

Mango_flat_beak_leafhopper

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info), Homemade: Soap or botanical spray: Prepare mild soap or plant extract solution and spray directly on infested areas. (No usage info)
-

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

Miridae

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info), Homemade: Soap or botanical spray: Prepare mild soap or plant extract solution and spray directly on infested areas. (No usage info)

Nipaecoccus_vastalor

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info), Homemade: Soap or botanical spray: Prepare mild soap or plant extract solution and spray directly on infested areas. (No usage info)

Orange_Haunglongbing_(Citrus_greening)

- Missing Usage Instructions for: Organic: Control psyllids (vectors). (No usage info), Organic: Remove infected trees immediately. (No usage info), Chemical: Imidacloprid (for vector control). (No usage info), Homemade: None. Tree removal is required. (No usage info)

Papilio_xuthus

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)

Parlatoria_zizyphus_Lucus

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)

Peach_Bacterial_spot

- Missing Usage Instructions for: Organic: Copper sprays (careful of phytotoxicity). (No usage info), Organic: Oxytetracycline. (No usage info), Chemical: Copper ammonium carbonate (No usage info), Homemade: Garlic/Pepper spray (limited efficacy). (No usage info)

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

Peach_healthy

- Missing Symptoms
 - Missing Organic Treatments
 - Missing Chemical (Inorganic) Treatments
 - Missing Homemade Treatments
-

Pepper_bell_Bacterial_spot

- Missing Usage Instructions for: Organic: Neem + bio copper (No usage info), Chemical: Copper oxychloride (No usage info), Homemade: Baking soda + soap (No usage info)
-

Pepper_bell_healthy

- Missing Symptoms
 - Missing Organic Treatments
 - Missing Chemical (Inorganic) Treatments
 - Missing Homemade Treatments
 - Missing Usage Instructions for: Entire record missing
-

Phyllocnistis_citrella_Stainton

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)
-

Phyllocoptes_oleiverus_ashmead

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)
-

Pieris_canidia

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)
-

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

Polyphagotars_onemus_latus

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)

Potato_Early_blight

- Missing Usage Instructions for: Organic: Copper fungicides. (No usage info), Organic: Serenade (Bacillus subtilis). (No usage info), Chemical: Chlorothalonil (No usage info), Chemical: Mancozeb (No usage info), Homemade: Baking soda spray. (No usage info)

Potato_Late_blight

- Missing Usage Instructions for: Organic: Copper fungicides (preventative, apply every 5-7 days). (No usage info), Organic: Remove and destroy all infected plants immediately (do not compost). (No usage info), Organic: Serenade (Bacillus subtilis) can suppress mild infections. (No usage info), Chemical: Chlorothalonil (Bravo) (No usage info), Chemical: Mancozeb (No usage info), Chemical: Mefenoxam (Ridomil Gold) for commercial use (No usage info), Homemade: None reliable for late blight. Immediate removal is the only home remedy to stop spread. (No usage info)

Potato_healthy

- Missing Symptoms
 - Missing Organic Treatments
 - Missing Chemical (Inorganic) Treatments
 - Missing Homemade Treatments
-

Potosiabre_vitarsis

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)

Prodenia_litura

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

Pseudococcus_comstocki_Kuwana

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)

Raspberry_healthy

- Missing Symptoms
- Missing Organic Treatments
- Missing Chemical (Inorganic) Treatments
- Missing Homemade Treatments

Rhytidodera_bowrinii_white

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)

Rice_Stemfly

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)

Salurnis_marginella_Guerr

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info)

Scirtothrips_dorsalis_Hood

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info), Homemade: Soap or botanical spray: Prepare mild soap or plant extract solution and spray

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

directly on infested areas. (No usage info)

Soybean_healthy

- Missing Symptoms
 - Missing Organic Treatments
 - Missing Chemical (Inorganic) Treatments
 - Missing Homemade Treatments
-

Squash_Powdery_mildew

- Missing Usage Instructions for: Organic: Neem oil. (No usage info), Organic: Sulfur. (No usage info), Chemical: Myclobutanil (No usage info), Homemade: Milk spray (1 part milk : 2 parts water). (No usage info), Homemade: Baking soda. (No usage info)
-

Sternochetus_frigidus

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info), Homemade: Soap or botanical spray: Prepare mild soap or plant extract solution and spray directly on infested areas. (No usage info)
-

Strawberry_Leaf_scorch

- Missing Usage Instructions for: Organic: Remove old leaves. (No usage info), Organic: Copper fungicide. (No usage info), Chemical: Captan (No usage info), Homemade: Chamomile tea spray (mild). (No usage info)
-

Strawberry_healthy

- Missing Symptoms
 - Missing Organic Treatments
 - Missing Chemical (Inorganic) Treatments
 - Missing Homemade Treatments
-

Tetradacus_c_Bactrocera_minax

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info), Homemade: Soap or botanical spray: Prepare mild soap or plant extract solution and spray

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

directly on infested areas. (No usage info)

Thrips

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info), Homemade: Soap or botanical spray: Prepare mild soap or plant extract solution and spray directly on infested areas. (No usage info)

Tomato_Bacterial_spot

- Missing Usage Instructions for: Organic: Copper-based bactericides (apply early at first sign). (No usage info), Organic: Agri-Strep (Streptomycin) if permitted. (No usage info), Organic: Remove severely infected plants. (No usage info), Chemical: Fixed Copper + Mancozeb (tank mix is more effective than copper alone) (No usage info), Chemical: Actigard (systemic acquired resistance inducer) (No usage info), Homemade: Diluted hydrogen peroxide spray (temporary reduction of surface bacteria). (No usage info)

Tomato_Early_blight

- Missing Usage Instructions for: Organic: Copper fungicide. (No usage info), Organic: Neem oil. (No usage info), Chemical: Chlorothalonil (No usage info), Chemical: Mancozeb (No usage info), Homemade: Baking soda mix (1 tbsp soda, 1 tsp soap, 1 gal water). (No usage info)

Tomato_Late_blight

- Missing Usage Instructions for: Organic: Copper spray (preventative only, apply before rain). (No usage info), Organic: Pull and bag infected plants immediately. (No usage info), Organic: Biofungicides (Bacillus amyloliquefaciens). (No usage info), Chemical: Chlorothalonil (Daconil) (No usage info), Chemical: Propamocarb (No usage info), Chemical: Cymoxanil (No usage info), Homemade: No effective home cure. Prevention and removal are key. (No usage info)

Tomato_Leaf_Mold

- Missing Usage Instructions for: Organic: Copper fungicide. (No usage info), Organic: Reduce humidity. (No usage info), Chemical: Chlorothalonil (No usage info), Homemade: Vinegar solution (caution). (No usage info)

Tomato_Septoria_leaf_spot

- Missing Usage Instructions for: Organic: Copper fungicide. (No usage info), Organic: Remove lower

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

infected leaves. (No usage info), Chemical: Chlorothalonil (No usage info), Homemade: Baking soda spray. (No usage info)

Tomato_Spider_mites_Two_spotted_spider_mite

- Missing Usage Instructions for: Organic: Neem oil (No usage info), Chemical: Abamectin (No usage info), Homemade: Water + soap spray (No usage info)

Tomato_Target_Spot

- Missing Usage Instructions for: Organic: Copper fungicide. (No usage info), Chemical: Chlorothalonil (No usage info), Chemical: Azoxystrobin (No usage info), Homemade: Baking soda mix. (No usage info)

Tomato_Tomato_YellowLeaf_Curl_Virus

- Missing Usage Instructions for: Organic: Control whiteflies (Neem/Soap). (No usage info), Organic: Remove infected plants (No cure). (No usage info), Chemical: Imidacloprid (for vector control). (No usage info), Homemade: Yellow sticky traps for whiteflies. (No usage info)

Tomato_Tomato_mosaic_virus

- Missing Usage Instructions for: Organic: Remove and burn infected plants immediately (No cure). (No usage info), Organic: Sanitize tools. (No usage info), Chemical: None. (No usage info), Homemade: Milk dip for tools (prevents spread). (No usage info)

Tomato_healthy

- Missing Symptoms
- Missing Usage Instructions for: Organic: Verification_Test_Medicine (No usage info), Chemical: Copper Fungicide (No usage info), Homemade: Baking Soda Spray (No usage info)

Toxoptera_aurantii

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info), Homemade: Soap or botanical spray: Prepare mild soap or plant extract solution and spray directly on infested areas. (No usage info)

Toxoptera_citricidus

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

- Missing Usage Instructions for: Organic: Neem oil (2%): Spray uniformly on affected plant parts every 10-14 days during infestation. (No usage info), Chemical: Chemical control: Apply recommended insecticide/fungicide at label dose when pest population exceeds economic threshold level. (No usage info), Homemade: Soap or botanical spray: Prepare mild soap or plant extract solution and spray directly on infested areas. (No usage info)

Trialeurodes_vaporariorum

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

Unaspis_yanonensis

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

Xylotrechus

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

alfalfa_plant_bug

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

alfalfa_seed_chalcid

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

alfalfa_weevil

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

aphids

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

army_worm

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

asiatic_rice_borer

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

beet_army_worm

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

beet_fly

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

beet_weevil

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

bird_cherry-oataphid

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

black_cutworm

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

blister_beetle

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info)

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

brown_plant_hopper

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

cabbage_army_worm

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

cerodonta_denticornis

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

corn_borer

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

english_grain_aphid

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

affected plant parts. (No usage info)

flax_budworm

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

flea_beetle

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

grain_spreader_thrips

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

green_bug

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

grub

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

large_cutworm

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

legume_blister_beetle

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

longlegged_spider_mite

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

lytta_polita

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

meadow_moth

- Missing Usage Instructions for: Organic: Neem-based formulations (2%): Spray on foliage at 10-14 day intervals during infestation period. (No usage info), Chemical: Chemical control: Use recommended insecticide or fungicide as per crop and label instructions when threshold levels are crossed. (No usage info), Homemade: Homemade remedy: Prepare soap or plant-extract solution and spray directly on affected plant parts. (No usage info)

mole_cricket

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)

odontothrips_loti

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)

paddy_stem_maggot

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)

parathrene_regalis

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)

peach_borer

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)

penthaleus_major

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)

red_spider

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)

rice_gall_midge

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)

rice_leaf_caterpillar

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)

rice_leaf_roller

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)

rice_leafhopper

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)

rice_shell_pest

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)

rice_water_weevil

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)

sericaorient_alismots_chulsky

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)

small_brown_plant_hopper

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)

tarnished_plant_bug

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade:

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas.
(No usage info)

therioaphis_maculata_Buckton

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas.
(No usage info)

wheat_blossom_midge

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas.
(No usage info)

wheat_phloeothrips

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas.
(No usage info)

wheat_sawfly

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas.
(No usage info)

white_backed_plant_hopper

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas.

Pest & Disease Data Gap Report

Generated on: 2026-02-18 22:14:23

(No usage info)

white_margined_moth

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)

wireworm

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)

yellow_cutworm

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)

yellow_rice_borer

- Missing Usage Instructions for: Organic: Neem oil / biopesticide (2%): Spray uniformly on affected plant parts every 10-14 days. (No usage info), Chemical: Chemical pesticide: Apply recommended formulation at label dose when economic threshold is exceeded. (No usage info), Homemade: Homemade spray: Prepare mild soap or botanical extract solution and spray directly on infested areas. (No usage info)
