

Pest Name	Symptoms	Organic	Inorganic	Homemade
Background without leaves	Background without leaves infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Ceroplastes rubens	Ceroplastes rubens infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Chlumetia transversa	Chlumetia transversa infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Cicadella viridis	Cicadella viridis infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Colomerus vitis	Colomerus vitis infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Dacus dorsalisHendel	Dacus dorsalisHendel infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Dasineura sp	Dasineura sp infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Deporaus marginatus Pascoe	Deporaus marginatus Pascoe infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.

Pest Name	Symptoms	Organic	Inorganic	Homemade
Erythroneura apicalis	Erythroneura apicalis infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Locustoidea	Locustoidea infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Mango flat beak leafhopper	Mango flat beak leafhopper infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Miridae	Miridae infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Nipaecoccus vastalor	Nipaecoccus vastalor infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Papilio xuthus	Papilio xuthus infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Parlatoria zizyphus Lucus	Parlatoria zizyphus Lucus infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Phyllocnistis citrella Stainton	Phyllocnistis citrella Stainton infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.

Pest Name	Symptoms	Organic	Inorganic	Homemade
Phyllocoptes oleiverus ashmead	Phyllocoptes oleiverus ashmead infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Pieris canidia	Pieris canidia infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Polyphagotars onemus latus	Polyphagotars onemus latus infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Potosiabre vitarsis	Potosiabre vitarsis infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Prodenia litura	Prodenia litura infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Pseudococcus comstocki Kuwana	Pseudococcus comstocki Kuwana infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Rhytidodera bowrinii white	Rhytidodera bowrinii white infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Salurnis marginella Guerr	Salurnis marginella Guerr infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.

Pest Name	Symptoms	Organic	Inorganic	Homemade
Scirtothrips dorsalis Hood	Scirtothrips dorsalis Hood infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Sternochetus frigidus	Sternochetus frigidus infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Tetradacus c Bactrocera minax	Tetradacus c Bactrocera minax infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Thrips	Thrips infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Toxoptera aurantii	Toxoptera aurantii infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Toxoptera citricidus	Toxoptera citricidus infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Trialeurodes vaporariorum	Trialeurodes vaporariorum infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
Unaspis yanonensis	Unaspis yanonensis infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.

Pest Name	Symptoms	Organic	Inorganic	Homemade
Xylotrechus	Xylotrechus infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
alfalfa plant bug	alfalfa plant bug infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
alfalfa seed chalcid	alfalfa seed chalcid infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
alfalfa weevil	alfalfa weevil infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
aphids	aphids infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
army worm	army worm infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
asiatic rice borer	asiatic rice borer infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
beet army worm	beet army worm infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.

Pest Name	Symptoms	Organic	Inorganic	Homemade
beet fly	beet fly infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
beet weevil	beet weevil infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
bird cherry-oataphid	bird cherry-oataphid infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
black cutworm	black cutworm infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
blister beetle	blister beetle infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
brown plant hopper	brown plant hopper infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
cabbage army worm	cabbage army worm infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
cerodonta denticornis	cerodonta denticornis infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.

Pest Name	Symptoms	Organic	Inorganic	Homemade
corn borer	corn borer infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
english grain aphid	english grain aphid infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
flax budworm	flax budworm infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
flea beetle	flea beetle infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
grain spreader thrips	grain spreader thrips infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
green bug	green bug infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
grub	grub infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
large cutworm	large cutworm infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
legume blister beetle	legume blister beetle infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.

Pest Name	Symptoms	Organic	Inorganic	Homemade
longlegged spider mite	longlegged spider mite infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
lytta polita	lytta polita infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
meadow moth	meadow moth infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
mole cricket	mole cricket infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
odontothrips loti	odontothrips loti infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
paddy stem maggot	paddy stem maggot infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
parathrene regalis	parathrene regalis infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
peach borer	peach borer infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.

Pest Name	Symptoms	Organic	Inorganic	Homemade
penthaleus major	penthaleus major infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
red spider	red spider infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
rice gall midge	rice gall midge infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
rice leaf caterpillar	rice leaf caterpillar infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
rice leaf roller	rice leaf roller infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
rice leafhopper	rice leafhopper infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
rice shell pest	rice shell pest infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
rice water weevil	rice water weevil infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.

Pest Name	Symptoms	Organic	Inorganic	Homemade
sericaorient alismots chulsky	sericaorient alismots chulsky infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop■specific recommended formulation at label■prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
small brown plant hopper	small brown plant hopper infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop■specific recommended formulation at label■prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
tarnished plant bug	tarnished plant bug infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop■specific recommended formulation at label■prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
therioaphis maculata Buckton	therioaphis maculata Buckton infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop■specific recommended formulation at label■prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
wheat blossom midge	wheat blossom midge infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop■specific recommended formulation at label■prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
wheat phloeothrips	wheat phloeothrips infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop■specific recommended formulation at label■prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
wheat sawfly	wheat sawfly infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop■specific recommended formulation at label■prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
white backed plant hopper	white backed plant hopper infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop■specific recommended formulation at label■prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.

Pest Name	Symptoms	Organic	Inorganic	Homemade
white margined moth	white margined moth infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
wireworm	wireworm infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
yellow cutworm	yellow cutworm infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.
yellow rice borer	yellow rice borer infestation shows symptoms such as discoloration, deformities, wilting, stunted growth, and yield reduction.	Neem oil / biopesticide (2%): Spray thoroughly on affected plant parts every 10–14 days.	Chemical pesticide: Apply crop-specific recommended formulation at label-prescribed dose when ETL is crossed.	Homemade remedy: Use mild soap solution or botanical extract and spray directly on infested areas.