


# Estimation



## Statistical Reasoning and Quantitative Methods

François Briatte & Ivaylo Petev

Session 5

# Outline


---

The properties of the standard normal distribution allow for statistical **inference**: the **estimation**, at a certain level of **confidence**, of the unobserved **population** parameters, using observed **sample** parameters.

But first. . .

## Assignment No. 1

### Estimation




# Assignment No. 1

---

## Univariate statistics

- Introduction
- Datasets
- Distributions
- Estimation

## Assignment No. 1



## Bivariate statistics


- Significance
- Crosstabulation
- Correlation
- Linear regression

## Assignment No. 2

## Statistical modelling

- Basics
- Extensions
- Diagnostics
- Conclusion

## Final paper



## How to proceed

---

**First of all, complete the projects table:** <http://goo.gl/brYmB>

**This table has to be complete for grading purposes.** It has to include your **names, class day and time, topic and dataset.**

Then:

1. Write your **research design** (using the paper template).
2. Write up your do-file and export your **summary statistics**.
3. **Replicate** your do-file and finalize your paper around 3 to 5 pages.

**Finally, email us your work,** **copying your partner to the email**, with the email subject “SRQM: Assignment No. 1, Briatte and Petev” (**substitute your own names**). The deadline is **our next meeting**.

## Step 1: Research design

---

Start a text document in about 2 to 4 paragraphs:

- **Topic:** describe your empirical problem in the form of a **research question** and **hypotheses**.
- **Data:** describe your dataset with its **source**, **sample size**, **sampling strategy** and **variables**.

You can revise your hypotheses and selection of variables in future drafts. For now, stick to writing only a few paragraphs, aiming at concision (short and precise descriptions).

The Stata Guide covers all necessary steps in Sections 1–9, with additional formatting instructions in Section 13 and a summary of instructions for this draft in Section 14.

## Topic and dataset

---

- Your **topic** formulates a research question and derives it into testable **hypotheses** between your dependent and independent variables. Cite previous research if you know any.
- Your **dataset** is a **fully referenced and documented** sample for which you have read the documentation to understand what each variable measures.

Note that the total **number of observations** on which you will work at later stages might decrease because of missing data, so make sure that you are initially working on the **largest possible sample**.

Use the **count**, **fre** and **su** commands to inspect missing data, and ignore variables for which there is an excess of missing data.

## Variables and description

---

- Your **variables** have been fully **described**, as well as **renamed** and/or **recoded** if necessary.
- Your **description** of the variables is a **summary statistics table** with **additional observations** in your text.

Note that the **number of variables** in your project is determined by your research design, and might evolve if you revise your hypotheses.

For example, you might select something like 6 to 12 independent variables and later focus on only some of them, or even add new ones. You will also be able to revise your do-file accordingly.

## Step 2: Summary statistics

---

Continue with an additional 2 to 4 paragraphs:

- **Summary statistics:** describe all variables in a few words.
- **Normality:** assess how normal your dependent variable is, and describe its potential transformation.

For tables and graphs:

- The **summary statistics table** has to describe *both* your **continuous** and **categorical** variables. The table is required to appear in your paper.
- The **histogram** that shows the **distribution of the dependent variable** can be completed by other plots showing interactions with independent variables.



## The `tsst` command

---

\* Short example.

```
use datasets/nhis2009, clear
```

\* Quick help.

```
tsst using test.txt
```

\* Quick test.

```
tsst using test.txt, su(age) fr(sex health) replace
```

Run `draft1.do` for a full example. Look at formatting instructions in the Stata Guide, Section 13.4, which includes an alternative to `tsst` if the command does not work. You can also load it manually:

\* Run this line before the example above if `tsst` fails.

```
run programs/tsst.ado
```

## Additional plots

---

- The **histogram** and **diagnostic plots** that show the distribution and (ab)normality of your dependent variable are required. Also try the `gladder` command to find any potential transformation.
- **Box plots** (`gr box` and `gr hbox`), **dot plots** and **bar plots** (`gr dot` and `gr bar`) with the `over` option can also be used to split your dependent variable over categorical independent variables.
- **Spline plots** with the `spineplot` command are also recommended if you have categorical independent variables to cross-visualize.

Your plots appear only in your do-file, except if you have *specific* observations to make about them. In your paper, include 0 to 2 plots.

## Step 3: Replication

---

Assignment No. 1 consists of:

- **A short paper** named in the format `Briatte_Petev_1.pdf`.
- **A short do-file** named in the format `Briatte_Petev_1.do`.

The do-file will include all that is needed to replicate your results: subsetting, variable renaming and recoding, summary statistics, normality tests, and tabular and graphical visualizations.

**Replicate your do-file** before sending it, to make sure that it executes properly and produces the results displayed in your analysis.

## Further help

---

- Course-specific help:

- Stata Guide
- Session do-files
- Course slides

- General help:


- Handbook chapters
- Stata documentation (`help command`)
- Online tutorials

Everything is systematically archived on the course website:

<http://f.briatte.org/teaching/quantil/>

Happy coding!


And now, estimation.



## Standard normal distribution $\mathcal{N}(0, 1)$

---


- $\mu \pm 1\sigma$  contains approximately **68%** of all values.
- $\mu \pm 2\sigma$  contains approximately **95%** of all values.
- $\mu \pm 3\sigma$  contains approximately **99%** of all values.



## Probability density function

---

- $Pr(\mu - 1\sigma < \mu < \mu + 1\sigma) \approx .68$
- $Pr(\mu - 2\sigma < \mu < \mu + 2\sigma) \approx .95$
- $Pr(\mu - 3\sigma < \mu < \mu + 3\sigma) \approx .99$



# Estimation

---

The normal distribution is used for **statistical inference**, i.e. for estimating the **population** parameters from the **sample** parameters.


| Parameter          | Notation  |            |
|--------------------|-----------|------------|
|                    | Sample    | Population |
| Mean               | $\bar{X}$ | $\mu$      |
| Standard deviation | $s$       | $\sigma$   |

This operation **generalizes** the values held by the sample to the population under consideration, under certain **levels of confidence**.



$$P(B_j|A) = \frac{P(B_j)P(A|B_j)}{P(A)}$$

$$P\left(\bigcup_{j=1}^n E_j\right) \leq \sum_{j=1}^n P(E_j)$$



$$\chi^2 = \sum_{i=1}^N (n_i - u_i)^2 P(n_i, N)$$

# Central Limit Theorem

---

Formally,  $\mu$  is the **population mean**, which is unobserved, and  $\bar{X}$  is the **sample mean**, which is observable by analysing the data.

The **Central Limit Theorem** (CLT) states that, if repeated samples are drawn from the population, their respective means  $\bar{X}_1, \bar{X}_2, \dots, \bar{X}_n$  will be **normally distributed** around the population mean  $\mu$ :

$$\text{CLT} : \sqrt{N} \left( \frac{1}{N} \sum_{i=1}^N \bar{X}_i - \mu \right) \xrightarrow{d} \mathcal{N}(0, \sigma^2)$$

The CLT holds **regardless of the distribution** of the variable  $X$  under examination, which is, like, totally awesome.



# Law of Large Numbers

---

Formally,  $\sigma$  is the **population standard deviation**, which is unobserved, and  $s$  is the **sample standard deviation**, which is observable by analysing the data.

The **Law of Large Numbers** states that, the larger the sample, the closer its mean will be to the population mean. The **standard error** of that estimate derives from the standard deviation of the variable.

The **standard error of the mean** (SEM), which estimates the standard deviation of the population from the standard deviation in the sample, will hence (slowly) decrease with sample size:

$$\text{SEM} = SE_{\bar{X}} = \frac{s}{\sqrt{N}} \text{ (sample)}$$



# Confidence intervals

---

Given these properties, the population mean  $\mu$  can be estimated from a sample  $X$  of  $N$  (statistically independent) observations.

**When we observe a normal distribution**,  $\mu \pm 2\sigma$  contains approximately 95% of all values. The exact number used for estimation at **95% confidence**, called the **z-score**, is  $z = 1.96$ .

**When the sample values of  $X$  are normally distributed**,  $\bar{X} \pm 1.96 \cdot SE_{\bar{X}}$  contains 95% of the possible values of  $\mu$ .

These bounds define a **95% confidence interval**:

$$\bar{X} - 1.96 \cdot SE_{\bar{X}} < \mu < \bar{X} + 1.96 \cdot SE_{\bar{X}}$$

- In 2.5% of cases,  $\mu < \bar{X} - 1.96$ .
- In 2.5% of cases,  $\mu > \bar{X} + 1.96$ .



## Stata implementation

---

Summary statistics for current worldwide fertility rates:

`su births`

| Variable | Obs | Mean     | Std. Dev. | Min | Max   |
|----------|-----|----------|-----------|-----|-------|
| births   | 186 | 3.138294 | 1.649826  | 1.1 | 7.446 |

Standard error and 95% confidence interval:

`ci births`

| Variable | Obs | Mean     | Std. Err. | [95% Conf. Interval] |          |
|----------|-----|----------|-----------|----------------------|----------|
| births   | 186 | 3.138294 | .1209711  | 2.899634             | 3.376954 |

- Use the `level(99)` option for a **larger** 99% confidence interval.
- Use the `binomial` option if the variable is **dichotomous** (binary).

## Stata implementation: Q&A

---

- **Q: Why is the 99% confidence interval larger?**

A: The 99% CI uses  $z = 2.58$  to include 99% of all possible values of  $\mu$  in the interval. Its calculation with  $\bar{X} \pm 2.58 \cdot SE_{\bar{X}}$  therefore includes more values on both sides of  $\bar{X}$ .

This is called the **precision-accuracy trade-off**: higher confidence for  $\mu$  is obtained at the expense of a more precise value for  $\mu$ .

**Maximising sample size** will attenuate the trade-off.

- **Q: Why do we apply a 'binomial' option to binary variables?**

A: Binary outcomes of “yes/no” variables form a distribution with no intermediate values that cannot be normal. Instead, the discrete probability of independent 0/1 outcomes (**Bernoulli trials**) is given by the **binomial distribution**.

We will focus on estimation based on the normal distribution.

## Back to estimation


Using the normal distribution as a probability density function and standardised z-scores to select a level of confidence:

| Parameter          | Notation  |            |
|--------------------|-----------|------------|
|                    | Sample    | Population |
| Observations       | $N$       | unobserved |
| Mean               | $\bar{X}$ | $\mu$      |
| Standard deviation | $s$       | $\sigma$   |
| Standard error     | $SE$      |            |

Estimating  $\mu$  at 95% or 99% confidence:

$$Pr(\mu \in \bar{X} \pm z = 1.96 \cdot SE_{\bar{X}}) = .95$$

$$Pr(\mu \in \bar{X} \pm z = 2.58 \cdot SE_{\bar{X}}) = .99$$



## Back to estimation


---

For a variable  $X$ , estimating  $\mu$  from  $\bar{X}$  relies on three prerequisites that you will apply to all your variables:

- **Assess the distribution's normality** i.e. whether  $X \sim \mathcal{N}(\mu, \sigma)$ .
- **Select the standardised z-score** to fix a **level of confidence**.
- **Minimise the sampling error** by maximising the **sample size** ( $N$ ).

This technique guides all **point estimation** that we perform in this course. It is called **Maximum Likelihood Estimation** (MLE).

Following the assumptions of MLE pleases **Estimation Cat**, the first of two cats in The Prophecy.





# The Prophecy

---

The Prophecy states that the powers of Estimation Cat (white) are limited by those of **Significance Cat** (black).

- **Estimation** provides a parameter and its **standard error**.  
Relationships between variables can be modelled as parameters.
- **Statistical significance** provides its **probability level**, i.e. the probability that the estimated parameter is different from zero.

**Parametric statistics** work by confronting the awesome powers of Estimation Cat and Significance Cat into a **model**.

This course is an introduction to statistical modelling using frequentist statistics, a.k.a The Prophecy.



