

# Lecture 7: Rare Variant Analysis: Collapsing Tests, Kernel (Variance Component) Tests and Omnibus Tests

Instructors: Joelle Mbatchou and Loic Yengo

Summer Institute in Statistical Genetics 2022

## Lecture Overview

1. Limitations of GWAS
2. Rationale for Rare Variant Analysis
3. Challenges
4. Collapsing/Burden Tests
5. Variance Component Tests
6. Omnibus Tests

## GWAS: Missing Heritability

- ▶ GWAS primarily focus on common variants ( $MAF \geq 5\%$ ) whose effects are small.
- ▶ **Missing heritability:** Significant GWAS SNPs explain a small proportion of disease heritability.
- ▶ Possible reasons:
  - ▶ GxG and GxE interactions?
  - ▶ Many common causal variants: Each with a small effect?
  - ▶ Epigenetics?
  - ▶ **Rare variants?**

## Why rare variants?

- Most of human variants are rare.
- Functional variants tend to be rare.

### Article

**Table 1 | Number of variants in 40,722 unrelated individuals in TOPMed**


	All unrelated individuals (n = 40,722)		A
	Total	Singletons (%)	
<b>Total variants</b>	<b>384,127,954</b>	<b>203,994,740 (53)</b>	
SNVs	357,043,141	189,429,596 (53)	
Indels	27,084,813	14,565,144 (54)	
<b>Novel variants</b>	<b>298,373,330</b>	<b>191,557,469 (64)</b>	
SNVs	275,141,134	177,410,620 (64)	
Indels	23,232,196	14,146,849 (61)	
<b>Coding variation</b>	<b>4,651,453</b>	<b>2,523,257 (54)</b>	
Synonymous	1,435,058	715,254 (50)	
Nonsynonymous	2,965,093	1,648,672 (56)	
Stop/essential splice	97,217	60,347 (62)	
Frameshift	104,704	71,577 (68)	
In-frame	51,997	29,110 (56)	

Novel variants are taken as variants that were not present in dbSNP build 149, the most recent dbSNP version v

Talium et al., *Nature* 2021

## Challenges in Association Studies for Rare Variants

- ▶ Compared to common variant studies, **individual SNP analysis in rare variant studies is seriously underpowered.**  
→ How many subjects are needed to achieve 80% of power ( $\alpha = 10^{-6}$ ) by single variant test?



- ▶ A lot more rare variants than common variants → larger multiple testing burden

## Challenges in Association Studies for Rare Variants

- ▶ Individual rare variant tests are underpowered
- ▶ Need **cost-effective study designs** to genotype a large number of individuals
- ▶ Need **powerful statistical methods and strategies** to test for associations
  - ▶ Region based analysis: genes, moving windows, networks/pathways
  - ▶ Integrate with bioinformatics: Incorporate functional information

## Region Based Analysis of Rare Variants

- ▶ Gene (or Region) based tests
- ▶ Strategy:
  - ▶ Identify all observed variants within a sequenced (sub)-region.
  - ▶ Regions: gene, regulatory region, ...
  - ▶ Test the joint effect of rare/common variants.

## Regression Models

- ▶  $p$  variants in a certain region.
- ▶ SNPs in a region  $\mathbf{G}_i = (g_{i1}, g_{i2}, \dots, g_{ip})'$ , ( $g_{ij} = 0, 1, 2$ )
- ▶ Covariates  $\mathbf{X}_i$  : age, gender, PC scores (for population stratification).
- ▶ Continuous/binary traits:

$$\mu_i / \log \left( \frac{\mu_i}{1 - \mu_i} \right) = \alpha_0 + \mathbf{X}_i' \boldsymbol{\alpha} + \mathbf{G}_i' \boldsymbol{\beta}$$

- ▶ Joint test of no genetic effect in region:

$$H_0 : \boldsymbol{\beta} = (\beta_1, \dots, \beta_p) = \mathbf{0}$$



## Major Classes of Tests

- ▶ Burden/Collapsing tests
- ▶ Supervised/Adaptive Burden/Collapsing tests
- ▶ Variance component (similarity) based tests
- ▶ Omnibus tests

## Collapsing/Burden Tests - Principle


- ▶ If  $p$  is large, multivariate test  $\beta = 0$  is not powerful ( $\text{df}=p$ ).
- ▶ Collapsing: Suppose  $\beta_1 = \dots = \beta_p = \beta$

$$\begin{aligned}\mu_i / \log \left( \frac{\mu_i}{1 - \mu_i} \right) &= \alpha_0 + \mathbf{X}_i' \boldsymbol{\alpha} + \mathbf{G}_i' \boldsymbol{\beta} \\ &= \alpha_0 + \mathbf{X}_i^T \boldsymbol{\alpha} + C_i \beta\end{aligned}$$

- ▶  $C_i = g_{i1} + \dots + g_{ip}$  : **genetic burden/score**
- ▶ Test  $H_0 : \beta = 0$  ( $\text{df}=1$ )
- ▶ **Key assumption:** all rare variants in region are causal variants with the same effect sizes and association directions.

## Burden Tests

- Collapse rare variants

Y	G <sub>1</sub>	G <sub>2</sub>	G <sub>3</sub>	G <sub>4</sub>		C
1	1	0	0	0		1
1	0	1	0	0		1
1	0	0	1	1		2
.	.	.	.	.		.
.	.	.	.	.		.
.	.	.	.	.		.
0	0	0	0	0		0
0	0	0	0	0		0
0	0	0	0	0		0

## Burden Tests

- ▶ Many different types of tests exist based on different aggregation rules to get  $C_i$ 
  - ▶ Reflects assumptions on genetic architecture
- ▶ **Existence of any rare variants can cause loss of function of a region** (e.g. CAST)

$$C_i = \begin{cases} 1 & \text{if } \sum_{j=1}^p g_{ij} > 0 \\ 0 & \text{if } \sum_{j=1}^p g_{ij} = 0 \end{cases}$$

- ▶ **Dominant genetic model** (e.g.. MZ-test/GRANVIL)

$$C_i = \sum_{j=1}^p I(g_{ij} > 0)$$

## Weighted Burden

- ▶ Assume that **rarer variants have larger effects**
- ▶ Suppose  $\beta_j = w_j\beta$ , where  $w_j = w(MAF_j)$ .
  - ▶ Ex:  $w(MAF_j) = 1/\sqrt{MAF_j(1 - MAF_j)}$  (Madsen and Browning).
- ▶ Weighted count of rare variants

$$C_i = w_1g_{i1} + \cdots + w_pg_{ip}$$

## Power of Burden Tests

- ▶ Power of burden tests depends on
  - ▶ Number of associated variants
  - ▶ Number of non-associated variants
  - ▶ Direction of the effects.
- ▶ **Powerful if most variants are causal and have effects in the same direction.**

## Variance component test

- ▶ Burden tests are not powerful, if
  - ▶ there exist variants with different association directions
  - ▶ many non-causal variants
- ▶ Variance component tests have been proposed to address this limitation.

## Sequence Kernel Association Test (SKAT)

- Recall the original regression models:

$$\mu_i / \log \left( \frac{\mu_i}{1 - \mu_i} \right) = \alpha_0 + \mathbf{X}_i^T \boldsymbol{\alpha} + \mathbf{G}_i^T \boldsymbol{\beta}$$

- Assume  $\beta_j \sim \text{dist.}(0, w_j^2 \tau)$ .
- $H_0 : \beta_1 = \dots = \beta_p = 0 \iff H_0 : \tau = 0$ .
- 1df test!



## Sequence Kernel Association Test (SKAT)

- ▶ Score test statistic for  $\tau = 0$ :

$$Q_{SKAT} = (\mathbf{y} - \hat{\boldsymbol{\mu}}_0)' \mathbf{K} (\mathbf{y} - \hat{\boldsymbol{\mu}}_0),$$

- ▶  $\mathbf{K} = \mathbf{G}\mathbf{W}\mathbf{W}\mathbf{G}'$  : weighted linear kernel (where  $\mathbf{W} = \text{diag}[w_1, \dots, w_p]$ ).
- ▶ It is a  $N \times N$  similarity matrix

# SKAT

- $Q_{SKAT}$  is a **weighted sum of single variant score statistics**

$$\begin{aligned} Q_{SKAT} &= (\mathbf{y} - \hat{\boldsymbol{\mu}}_0)' \mathbf{G} \mathbf{W} \mathbf{W} \mathbf{G}' (\mathbf{y} - \hat{\boldsymbol{\mu}}_0) \\ &= \sum_{j=1}^p w_j^2 [\mathbf{g}_j' (\mathbf{y} - \hat{\boldsymbol{\mu}}_0)] = \sum_{j=1}^p w_j^2 S_j^2 \end{aligned}$$


- $S_j$  is a score test statistic in the SNP  $j$  only model:

$$\mu_i / \log \left( \frac{\mu_i}{1 - \mu_i} \right) = \alpha_0 + \mathbf{X}_i^T \boldsymbol{\alpha} + g_{ij} \beta_j$$

- Under  $H_0$ ,  $Q_{SKAT}$  (asymptotically) follows a **mixture of  $\chi^2$  distribution**  $\sum_{j=1}^p \lambda_j \chi_{1,j}^2$

# Burden vs SKAT

- ▶ Power simulations: 5% of the variants in region are causal & vary the directions of effects
- ▶ SKAT remains powerful even if variants have different effect directions

Lee et al., *Am J Hum Genet* 2011

## SKAT vs. Collapsing

- ▶ Collapsing tests are more powerful when a large % of variants are causal and effects are in the same direction.
- ▶ SKAT is more powerful when a small % of variants are causal, or the effects have mixed directions.
- ▶ Both scenarios can happen when scanning the genome.
- ▶ Best test to use depends on the underlying biology.
  - Difficult to choose which test to use in practice.

## SKAT vs. Collapsing

- ▶ Collapsing tests are more powerful when a large % of variants are causal and effects are in the same direction.
- ▶ SKAT is more powerful when a small % of variants are causal, or the effects have mixed directions.
- ▶ Both scenarios can happen when scanning the genome.
- ▶ Best test to use depends on the underlying biology.
  - Difficult to choose which test to use in practice.

We want to develop a unified test that works well in both situations → Omnibus tests

## Combine Test Statistics: Unified Test Statistics

Lee (2012). *Biostatistics*

- ▶ Combined Test of Burden tests and SKAT

$$Q_{\rho} = (1 - \rho)Q_{SKAT} + \rho Q_{Burden}, \quad 0 \leq \rho \leq 1.$$

- ▶  $Q_{\rho}$  includes SKAT and burden tests.
  - ▶  $\rho = 0$ : SKAT
  - ▶  $\rho = 1$ : Burden

# SKATO

- ▶ Model:


$$\mu_i / \log \left( \frac{\mu_i}{1 - \mu_i} \right) = \alpha_0 + \mathbf{X}_i^T \boldsymbol{\alpha} + \mathbf{G}_i^T \boldsymbol{\beta}$$

where  $\beta_j/w_j$  follows any arbitrary distribution with mean 0 and variance  $\tau$  and the correlation among  $\beta_j$ 's is  $\rho$ .

- ▶ Special cases:
  - ▶ SKAT:  $\rho = 0$
  - ▶ Burden:  $\rho = 1$
- ▶ Set a grid of values for  $\rho$  in  $[0, 1]$  and pick  $\rho$  which maximizes power
- ▶ SKATO p-value is obtained through numerical integration

# SKATO vs Burden/SKAT

- SKATO remains powerful across all scenarios



Lee et al., *Biostatistics* 2012



## Aggregated Cauchy Association Test: ACAT

- ▶ Based on the Cauchy combination method to combine a set of p-values  $\{p_j\}$ :

$$T_{ACAT} = \sum_j w_j \tan\{\pi(0.5 - p_j)\}$$

- ▶ Computing p-value is extremely fast

$$\text{p-value} \approx 0.5 - \frac{\arctan\{T_{ACAT}/w\}}{\pi}, \quad w = \sum_j w_j$$

- ▶ Insensitive to correlation in the p-values (can be unknown)

## Aggregated Cauchy Association Tests: ACATV & ACATO

### ▶ ACATV


- ▶ Apply ACAT to single variant p-values from rare variants
- ▶ More powerful when fewer variants are associated (i.e. sparse alternative)
- ▶ SKAT & Burden can lose substantial power under this scenario

### ▶ ACATO

- ▶ Apply ACAT to combine the p-values of SKAT, Burden and ACATV
- ▶ Omnibus test which should work well whether
  - ▶ Effects are in same direction & many variants are associated
  - ▶ Effects are in different directions
  - ▶ Very few variants are causal

## ACAT/SKAT/Burden

- ACATO remains powerful across all scenarios



Liu et al., *AJHG* 2019

## Summary

- ▶ Region based tests can increase the power of rare variants analysis.
- ▶ Relative performance of rare variant tests depends on underlying disease models
- ▶ Combined tests (omnibus tests), e.g, SKATO/ACATO, are more robust and powerful in different scenarios