



basic education

Department:
Basic Education
REPUBLIC OF SOUTH AFRICA

NATIONAL SENIOR CERTIFICATE

GRADE 12

MATHEMATICS P2

NOVEMBER 2022

MARKS: 150

TIME: 3 hours

This question paper consists of 13 pages and 1 information sheet.


INSTRUCTIONS AND INFORMATION

Read the following instructions carefully before answering the questions.

1. This question paper consists of 10 questions.
2. Answer ALL the questions in the SPECIAL ANSWER BOOK provided.
3. Clearly show ALL calculations, diagrams, graphs, etc. which you have used in determining your answers.
4. Answers only will NOT necessarily be awarded full marks.
5. You may use an approved scientific calculator (non-programmable and non-graphical), unless stated otherwise.
6. If necessary, round off answers correct to TWO decimal places, unless stated otherwise.
7. Diagrams are NOT necessarily drawn to scale.
8. An information sheet with formulae is included at the end of the question paper.
9. Write neatly and legibly.

QUESTION 1

The matric class of a certain high school had to vote for the chairperson of the RCL (representative council of learners). The scatter plot below shows the IQ (intelligence quotient) of the 10 learners who received the most votes and the number of votes that they received.



Before the election, the popularity of each of these ten learners was established and a popularity score (out of a 100) was assigned to each. The popularity scores and the number of votes of the same 10 learners who received the most votes are shown in the table below.


Popularity score (x)	32	89	35	82	50	59	81	40	79	65
Number of votes (y)	9	22	10	21	11	15	20	12	19	16

- 1.1 Calculate the:
 - 1.1.1 Mean number of votes that these 10 learners received (2)
 - 1.1.2 Standard deviation of the number of votes that these 10 learners received (1)
- 1.2 The learners who received fewer votes than one standard deviation below the mean were not invited for an interview. How many learners were invited? (2)
- 1.3 Determine the equation of the least squares regression line for the data given in the table. (3)
- 1.4 Predict the number of votes that a learner with a popularity score of 72 will receive. (2)
- 1.5 Using the scatter plot and table above, provide a reason why:
 - 1.5.1 IQ is not a good indicator of the number of votes that a learner could receive (1)
 - 1.5.2 The prediction in QUESTION 1.4 is reliable (1)

[12]

QUESTION 2

A company conducted research among all its employees on what percentage of their monthly salary was spent on fuel in a particular month. The data is represented in the ogive (cumulative frequency graph) below.




- 2.1 How many people are employed at this company? (1)
- 2.2 Write down the modal class of the data. (1)
- 2.3 How many employees spent more than 22,5% of their monthly salary on fuel? (2)
- 2.4 An employee spent R2 400 of his salary on fuel in that particular month. Determine the monthly salary of this employee if he spends 7% of his salary on fuel. (2)
- 2.5 The monthly salaries of these employees remains constant and the number of litres of fuel used in each month also remains constant. If the fuel price increases from R21,43 per litre to R22,79 per litre at the beginning of the next month, how will the above ogive change? (2)

[8]

QUESTION 3

In the diagram, $A(4; 2)$, $B(6; -4)$ and $C(-2; -3)$ are vertices of $\triangle ABC$. T is the midpoint of CB . The equation of line AC is $5x - 6y = 8$. The angle of inclination of AB is α . $\triangle DCT$ is drawn such that $CD \parallel BA$. The lines AC and DT intersect at S , the y -intercept of AC . P , F and R are the x -intercepts of DC , AC and AB respectively.



3.1 Calculate the:

3.1.1 Gradient of AB (2)

3.1.2 Size of α (2)

3.1.3 Coordinates of T (2)

3.1.4 Coordinates of S (2)

3.2 Determine the equation of CD in the form $y = mx + c$. (3)

3.3 Calculate the:


3.3.1 Size of \hat{DCA} (4)

3.3.2 Area of $\triangle POSC$ (5)

[20]

QUESTION 4

In the diagram, $M(3 ; -5)$ is the centre of the circle having PN as its diameter. KL is a tangent to the circle at $N(7 ; -2)$.



4.1 Calculate the coordinates of P . (2)

4.2 Determine the equation of:

4.2.1 The circle in the form $(x-a)^2 + (y-b)^2 = r^2$ (3)

4.2.2 KL in the form $y = mx + c$ (5)

4.3 For which values of k will $y = -\frac{4}{3}x + k$ be a secant to the circle? (4)

4.4 Points $A(t ; t)$ and B are not shown on the diagram.

From point A , another tangent is drawn to touch the circle with centre M at B .

4.4.1 Show that the length of tangent AB is given by $\sqrt{2t^2 + 4t + 9}$. (2)

4.4.2 Determine the minimum length of AB . (4)
[20]

QUESTION 5

5.1 Given that $\sqrt{13} \sin x + 3 = 0$, where $x \in (90^\circ; 270^\circ)$.

Without using a calculator, determine the value of:

5.1.1 $\sin(360^\circ + x)$ (2)

5.1.2 $\tan x$ (3)

5.1.3 $\cos(180^\circ + x)$ (2)

5.2 Determine the value of the following expression, **without using a calculator**:

$$\frac{\cos(90^\circ + \theta)}{\sin(\theta - 180^\circ) + 3 \sin(-\theta)} \quad (5)$$

5.3 Determine the general solution of the following equation:

$$(\cos x + 2 \sin x)(3 \sin 2x - 1) = 0 \quad (6)$$

5.4 Given the identity: $\cos(x + y) \cdot \cos(x - y) = 1 - \sin^2 x - \sin^2 y$

5.4.1 Prove the identity. (4)

5.4.2 Hence, determine the value of $1 - \sin^2 45^\circ - \sin^2 15^\circ$, **without using a calculator**. (3)

5.5 Consider the trigonometric expression: $16 \sin x \cdot \cos^3 x - 8 \sin x \cdot \cos x$


5.5.1 Rewrite the expression as a single trigonometric ratio. (4)

5.5.2 For which value of x in the interval $x \in [0^\circ; 90^\circ]$ will $16 \sin x \cdot \cos^3 x - 8 \sin x \cdot \cos x$ have its minimum value? (1)

[30]

QUESTION 6

In the diagram below, the graphs of $f(x) = \tan x$ and $g(x) = 2\sin 2x$ are drawn for the interval $x \in [-180^\circ; 180^\circ]$. A($60^\circ; k$) and B are two points of intersection of f and g .




- 6.1 Write down the period of g . (1)
- 6.2 Calculate the:
- 6.2.1 Value of k (1)
- 6.2.2 Coordinates of B (1)
- 6.3 Write down the range of $2g(x)$. (2)
- 6.4 For which values of x will $g(x+5^\circ) - f(x+5^\circ) \leq 0$ in the interval $x \in [-90^\circ; 0^\circ]$? (2)
- 6.5 Determine the values of p for which $\sin x \cdot \cos x = p$ will have exactly two real roots in the interval $x \in [-180^\circ; 180^\circ]$. (3)
- [10]**

QUESTION 7

AB is a vertical flagpole that is $\sqrt{5}p$ metres long. AC and AD are two cables anchoring the flagpole. B, C and D are in the same horizontal plane.

$BD = 2p$ metres, $\hat{ACD} = x$ and $\hat{ADC} = 45^\circ$.




- 7.1 Determine the length of AD in terms of p . (2)
- 7.2 Show that the length of $CD = \frac{3p(\sin x + \cos x)}{\sqrt{2} \sin x}$. (5)
- 7.3 If it is further given that $p = 10$ and $x = 110^\circ$, calculate the area of $\triangle ADC$. (3)

[10]

QUESTION 8

- 8.1 In the diagram, O is the centre of the circle. $MNPR$ is a cyclic quadrilateral and SN is a diameter of the circle. Chord MS and radius OR are drawn. $\hat{M}_2 = 64^\circ$.




Determine, giving reasons, the size of the following angles:

8.1.1 \hat{P} (2)

8.1.2 \hat{M}_1 (2)

8.1.3 \hat{O}_1 (2)


- 8.2 In the diagram, $\triangle ABG$ is drawn. D and E are midpoints of AB and AG respectively. AG and BG are produced to C and H respectively. F is a point on BC such that $FG \parallel CH$.



- 8.2.1 Give a reason why $DE \parallel BH$. (1)
- 8.2.2 If it is further given that $\frac{FC}{BF} = \frac{1}{4}$, $DE = 3x - 1$ and $GH = x + 1$, calculate, giving reasons, the value of x . (6)
- [13]

QUESTION 9


- 9.1 In the diagram, O is the centre of a circle. OD bisects chord AB .



Prove the theorem that states that the line from the centre of a circle that bisects a chord is perpendicular to the chord, i.e. $OD \perp AB$.

(5)

- 9.2 In the diagram, E, B, F, S and P are points on the circle centred at O . GB is a tangent to the circle at B . FE is produced to meet the tangent at G . OT is drawn such that T is the midpoint of EF . GO and BO are drawn. BS is drawn through T . $PS \parallel GF$.



Prove, giving reasons, that:

- 9.2.1 $OTBG$ is a cyclic quadrilateral

(5)


- 9.2.2 $\hat{GOB} = \hat{S}$

(4)

[14]

QUESTION 10

In the diagram, PQRS is a cyclic quadrilateral. KP is a tangent to the circle at P. C and D are points on chords PQ and PS respectively and CD produced meets RS produced at A. CA \parallel QS. RC is drawn. $\hat{P}_1 = \hat{R}_2$.



Prove, giving reasons, that:

10.1 $\hat{S}_1 = \hat{T}_2$ (4)

10.2 $\frac{AD}{AR} = \frac{AS}{AC}$ (5)

10.3 $AC \times SD = AR \times TC$ (4)
[13]

TOTAL: 150

INFORMATION SHEET

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

$$A = P(1 + ni)$$

$$A = P(1 - ni)$$

$$A = P(1 - i)^n$$

$$A = P(1 + i)^n$$

$$T_n = a + (n - 1)d$$

$$S_n = \frac{n}{2} [2a + (n - 1)d]$$

$$T_n = ar^{n-1}$$

$$S_n = \frac{a(r^n - 1)}{r - 1}; r \neq 1$$

$$S_\infty = \frac{a}{1 - r}; -1 < r < 1$$

$$F = \frac{x[(1 + i)^n - 1]}{i}$$

$$P = \frac{x[1 - (1 + i)^{-n}]}{i}$$

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x + h) - f(x)}{h}$$

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

$$M\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$$

$$y = mx + c$$

$$y - y_1 = m(x - x_1)$$

$$m = \frac{y_2 - y_1}{x_2 - x_1}$$

$$m = \tan \theta$$

$$(x - a)^2 + (y - b)^2 = r^2$$

$$\text{In } \triangle ABC: \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

$$a^2 = b^2 + c^2 - 2bc \cdot \cos A$$

$$\text{area } \triangle ABC = \frac{1}{2} ab \cdot \sin C$$

$$\sin(\alpha + \beta) = \sin \alpha \cdot \cos \beta + \cos \alpha \cdot \sin \beta$$

$$\sin(\alpha - \beta) = \sin \alpha \cdot \cos \beta - \cos \alpha \cdot \sin \beta$$

$$\cos(\alpha + \beta) = \cos \alpha \cdot \cos \beta - \sin \alpha \cdot \sin \beta$$

$$\cos(\alpha - \beta) = \cos \alpha \cdot \cos \beta + \sin \alpha \cdot \sin \beta$$

$$\cos 2\alpha = \begin{cases} \cos^2 \alpha - \sin^2 \alpha \\ 1 - 2\sin^2 \alpha \\ 2\cos^2 \alpha - 1 \end{cases}$$

$$\sin 2\alpha = 2\sin \alpha \cdot \cos \alpha$$

$$\bar{x} = \frac{\sum x}{n}$$

$$\sigma^2 = \frac{\sum_{i=1}^n (x_i - \bar{x})^2}{n}$$

$$P(A) = \frac{n(A)}{n(S)}$$

$$P(A \text{ or } B) = P(A) + P(B) - P(A \text{ and } B)$$

$$\hat{y} = a + bx$$

$$b = \frac{\sum (x - \bar{x})(y - \bar{y})}{\sum (x - \bar{x})^2}$$



basic education

Department:
Basic Education
REPUBLIC OF SOUTH AFRICA

**NATIONAL
SENIOR CERTIFICATE/
NASIONALE
SENIOR SERTIFIKAAT**

GRADE 12/GRAAD 12

MATHEMATICS P2/WISKUNDE V2

NOVEMBER 2022

MARKING GUIDELINES/NASIENRIGLYNE

MARKS/PUNTE: 150

**These marking guidelines consist of 24 pages.
*Hierdie nasienriglyne bestaan uit 24 bladsye.***

NOTE:


- If a candidate answers a question **TWICE**, only mark the **FIRST** attempt.
- If a candidate has crossed out an attempt of a question and not redone the question, mark the crossed out version.
- Consistent accuracy applies in **ALL** aspects of the marking memorandum. Stop marking at the second calculation error.
- Assuming answers/values in order to solve a problem is **NOT** acceptable.

NOTA:

- As 'n kandidaat 'n vraag **TWEE KEER** beantwoord, merk slegs die **EERSTE** poging.
- As 'n kandidaat 'n antwoord van 'n vraag doodtrek en nie oordoen nie, merk die doodgetrekte poging.
- Volgehoue akkuraatheid word in **ALLE** aspekte van die memorandum toegepas. Hou op nasien by die tweede berekeningsfout.
- Aanvaar van antwoorde/waardes om 'n probleem op te los, word **NIE** toegelaat nie.

GEOMETRY/MEETKUNDE	
S	A mark for a correct statement (A statement mark is independent of a reason)
	<i>'n Punt vir 'n korrekte bewering</i> (<i>'n Punt vir 'n bewering is onafhanklik van die rede</i>)
R	A mark for the correct reason (A reason mark may only be awarded if the statement is correct)
	<i>'n Punt vir 'n korrekte rede</i> (<i>'n Punt word slegs vir die rede toegeken as die bewering korrek is</i>)
S/R	Award a mark if statement AND reason are both correct
	<i>Ken 'n punt toe as die bewering EN rede beide korrek is</i>


QUESTION/VRAAG 1




Popularity score (x) <i>Gewildheidspunt (x)</i>	32	89	35	82	50	59	81	40	79	65
Number of votes (y) <i>Getal stemme (y)</i>	9	22	10	21	11	15	20	12	19	16

1.1.1	$\bar{y} = \frac{155}{10}$ $= 15,5$ <div style="border: 1px solid black; padding: 5px; display: inline-block;">ANSWER ONLY: Full marks</div>	✓ 155 ✓ answer (2)
1.1.2	SD = 4,59	✓ answer (1)
1.2	$\bar{y} - SD$ $= 15,5 - 4,59$ $= 10,91$ $\therefore 10 - 2 = 8 \text{ learners}$	✓ value of $\bar{y} - SD$ ✓ answer (2)
1.3	$a = 1,7709...$ $b = 0,2243...$ $\hat{y} = 1,77 + 0,22x$	✓ a ✓ b ✓ equation (3)
1.4	$\hat{y} = 1,77 + 0,22(72)$ $= 17,61$ $\approx 18 \text{ votes}$ <p>OR/OF</p> $\hat{y} = 17,92 \approx 18 \text{ votes}$	✓ substitution ✓ answer (2) ✓✓ answer (2)
1.5.1	Points are all scattered therefore low correlation and unrealistic prediction./ <i>Punte is versprei daarom 'n lae korrelasie en onrealistiese voorspelling.</i>	✓ R (1)
1.5.2	$r = 0,98$ /correlation very strong/ <i>korrelasie baie sterk</i> \therefore a reliable prediction/ <i>'n betroubare voorspelling</i>	✓ S (1)
[12]		

QUESTION/VRAAG 2



2.1	60 employees	✓ answer (A)	(1)
2.2	$20 < x \leq 25$	✓ answer	(1)
2.3	$60 - 34 = 26$ employees <div>ANSWER ONLY: Full marks</div>	✓ 34 ✓ answer	(2)
2.4	$\text{Salary} = \frac{100}{7} \times 2400$ $\text{Salary} = \text{R}34\,285,71$ <div>ANSWER ONLY: Full marks</div>	✓ method ✓ answer	(2)
2.5	\therefore Ogive/Cumulative frequency graph will shift to the right/will become steeper. \therefore Ogief/Kumulatiewe frekwensie grafiek sal na regs skuif/sal steiler wees.	✓✓ answer	(2)
[8]			

QUESTION/VRAAG 3


3.1.1	$m_{AB} = \frac{2 - (-4)}{4 - 6}$ OR $m_{AB} = \frac{-4 - 2}{6 - 4}$ $m_{AB} = -3$ <div style="border: 1px solid black; padding: 2px; display: inline-block;">ANSWER ONLY: Full marks</div>	✓ substitution ✓ answer (2)
3.1.2	$\tan \alpha = m_{AB} = -3$ $\alpha = 108,43^\circ$ <div style="border: 1px solid black; padding: 2px; display: inline-block;">ANSWER ONLY: Full marks</div>	✓ $\tan \alpha = m_{AB} = -3$ ✓ answer (2)
3.1.3	$T\left(\frac{x_1 + x_2}{2}; \frac{y_1 + y_2}{2}\right)$ $T\left(\frac{-2 + 6}{2}; \frac{-3 - 4}{2}\right)$ $T\left(2; \frac{-7}{2}\right)$	✓ $x_T = 2$ ✓ $y_T = \frac{-7}{2}$ (2)
3.1.4	$5(0) - 6y = 8$ $y = -\frac{4}{3}$ $S\left(0; \frac{-4}{3}\right)$	✓ $x_S = 0$ ✓ $y_S = \frac{-4}{3}$ (2)
3.2	$m_{CD} = m_{AB} = -3$ $-3 = -3(-2) + c$ OR $y - (-3) = -3(x - (-2))$ $c = -9$ $y = -3x - 9$ $y = -3x - 9$	✓ gradient ✓ substitution of $C(-2; -3)$ ✓ equation (3)

3.3.1	$5x - 6y = 8$ $y = \frac{5}{6}x - \frac{8}{6}$ $\tan \theta = m_{AC} = \frac{5}{6}$ $\theta = 39,81^\circ$ $\hat{A} = 108,43^\circ - 39,81^\circ$ $= 68,62^\circ$ $\hat{DCA} = 68,62^\circ$ [alt \angle s ; DC AB]	$\checkmark \tan \theta = m_{AC} = \frac{5}{6}$ $\checkmark \theta = 39,81^\circ$ $\checkmark \hat{A} = 68,62^\circ$ \checkmark answer (4)
3.3.2	<p>P(-3;0) and F(1,6 ; 0)</p> <p>Area POSC = Area ΔFPC – Area ΔOFS</p> $= \frac{1}{2}(4,6)(3) - \frac{1}{2}(1,6)\left(\frac{4}{3}\right)$ $= 6,9 - 1,07$ $= 5,83 \text{ units}^2$ <p>OR/OF</p> <p>P(-3;0)</p> $FC = \sqrt{\left(-2 - \frac{8}{5}\right)^2 + (-3 - 0)^2} = \frac{3\sqrt{61}}{5}$ $\text{Area } \Delta \text{PFC} = \frac{1}{2}(\text{PF})(\text{FC})\sin \hat{\text{OFS}}$ $= \frac{1}{2}\left(\frac{23}{5}\right)\left(\frac{3\sqrt{61}}{5}\right)\sin 39,81^\circ$ $= 6,90$ $\text{Area } \Delta \text{OFS} = \frac{1}{2}\left(\frac{8}{5}\right)\left(\frac{4}{3}\right)$ $= 1,07$ <p>Area POSC = 6,90 – 1,07</p> $= 5,83 \text{ units}^2$ <p>OR/OF</p>	\checkmark P(-3;0) \checkmark method $\checkmark \frac{1}{2}(4,6)(3)$ $\checkmark \frac{1}{2}(1,6)\left(\frac{4}{3}\right)$ \checkmark answer (5) \checkmark P(-3;0) $\checkmark \frac{1}{2}\left(\frac{23}{5}\right)\left(\frac{3\sqrt{61}}{5}\right)\sin 39,81^\circ$ $\checkmark \frac{1}{2}\left(\frac{8}{5}\right)\left(\frac{4}{3}\right)$ \checkmark method \checkmark answer (5)

	<div data-bbox="339 248 884 539" data-label="Figure"> </div> <p>P(-3;0)</p> <p>Area of POSC = Area of OSCR + Area of $\triangle PRC$</p> $= \frac{1}{2} \left(\frac{4}{3} + 3 \right) \times 2 + \frac{1}{2} (1 \times 3)$ $= \frac{35}{6}$ $= 5,83 \text{ units}^2$ <p>OR/ OF</p> <div data-bbox="339 1111 884 1413" data-label="Figure"> </div> <p>P(-3;0)</p> <p>Area POSC = Area ROSW + Area $\triangle PRC$ + Area $\triangle WSC$</p> $= \left(\frac{4}{3} \right) (2) + \frac{1}{2} (1)(3) + \frac{1}{2} (2) \left(\frac{5}{3} \right)$ $= \frac{35}{6}$ $= 5,83 \text{ units}^2$ <p>OR/OF</p>	<p>✓ P(-3;0)</p> <p>✓ method</p> <p>✓ $\frac{1}{2} \left(\frac{4}{3} + 3 \right) \times 2$ ✓ $\frac{1}{2} (1 \times 3)$</p> <p>✓ answer (5)</p> <p>✓ P(-3;0)</p> <p>✓ method</p> <p>✓ $\frac{1}{2} (1)(3)$</p> <p>✓ $\left(\frac{4}{3} \right) (2) + \frac{1}{2} (2) \left(\frac{5}{3} \right)$</p> <p>✓ answer (5)</p>
--	---	---

	$P(-3;0)$ $\text{Area of } \triangle PSC = \frac{1}{2}(PC)(CS) \sin \hat{DCA}$ $= \frac{1}{2}(\sqrt{10})\left(\frac{\sqrt{61}}{3}\right) \sin 68,62^\circ$ $= 3,833..$ $\text{Area of } \triangle POS = \frac{1}{2}(PO)(OS)$ $= \frac{1}{2}(3)\left(\frac{4}{3}\right)$ $= 2$ $\text{Area POSC} = 3,833... + 2$ $= 5,83\text{units}^2$	$\checkmark P(-3;0)$ $\checkmark \frac{1}{2}(\sqrt{10})\left(\frac{\sqrt{61}}{3}\right) \sin 68,62^\circ$ $\checkmark \frac{1}{2}(3)\left(\frac{4}{3}\right)$ $\checkmark \text{ method}$ $\checkmark \text{ answer}$ <div style="text-align: right;">(5)</div>
		[20]

QUESTION/VRAAG 4



4.1	$P(x; y); N(7; -2); M(3; -5)$ $\frac{x+7}{2}=3 \quad \frac{y-2}{2}=-5$ $x=-1 \quad y=-8$ $P(-1; -8)$	$\checkmark \quad x_p = -1 \quad \checkmark \quad y_p = -8$ (2)
4.2.1	$r^2 = (7-3)^2 + (-2-(-5))^2$ OR/OR $r^2 = (-1-3)^2 + (-8-(-5))^2$ $r^2 = 25$ $(x-3)^2 + (y+5)^2 = 25$	\checkmark substitution into distance formula $\checkmark \quad (x-3)^2 + (y+5)^2$ $\checkmark \quad r^2 = 25$ (3)
4.2.2	$m_{\text{radius}} = \frac{-5-(-2)}{3-7} = \frac{3}{4}$ $m_{\text{tangent}} = -\frac{4}{3}$ [radius \perp tangent/raaklyn \perp radius] $-2 = -\frac{4}{3}(7) + c$ OR $y-(-2) = -\frac{4}{3}(x-7)$ $c = \frac{22}{3}$ $y = -\frac{4}{3}x + \frac{22}{3}$	\checkmark substitution $\checkmark \quad m_{\text{radius}} = \frac{-3}{-4} = \frac{3}{4}$ $\checkmark \quad m_{\text{tangent}} = -\frac{4}{3}$ \checkmark substitution of m and $N(7; -2)$ \checkmark equation (5)
4.3	$-8 = -\frac{4}{3}(-1) + c$ $\therefore c = -\frac{28}{3}$ $-\frac{28}{3} < k < \frac{22}{3}$	\checkmark subst m and P \checkmark value of c $\checkmark \checkmark$ answer (4)

4.4.1	$AB^2 = AM^2 - MB^2$ $AB^2 = \left[(t-3)^2 + (t+5)^2 \right] - 5^2$ $= t^2 - 6t + 9 + t^2 + 10t + 25 - 25$ $AB = \sqrt{2t^2 + 4t + 9}$	✓ substitution into Pythagoras ✓ simplification (A) (2)
4.4.2	$t = \frac{-4}{2(2)}$ $= -1$ <p>Minimum at $t = -1$</p> $AB = \sqrt{2(-1)^2 + 4(-1) + 9}$ $AB = \sqrt{7}$ <p>OR/OF</p> $4t + 4 = 0$ $t = -1$ <p>Minimum at $t = -1$</p> $AB = \sqrt{2(-1)^2 + 4(-1) + 9}$ $AB = \sqrt{7}$ <p>OR/OF</p> <p>Length of $AB = \sqrt{2t^2 + 4t + 9}$</p> $= \sqrt{2\left(t^2 + 2t + \frac{9}{2}\right)}$ $= \sqrt{2\left[(t+1)^2 + \frac{7}{2}\right]}$ $= \sqrt{2(t+1)^2 + 7}$ <p>Minimum at $t = -1$</p> $AB = \sqrt{2(-1)^2 + 4(-1) + 9}$ $AB = \sqrt{7}$	✓ substitution into correct formula ✓ $t = -1$ ✓ substitution ✓ answer (4) ✓ derivative = 0 ✓ $t = -1$ ✓ substitution ✓ answer (4) ✓ completing of the square ✓ $t = -1$ ✓ substitution ✓ answer (4)
		[20]


QUESTION/VRAAG 5

5.1.1	$\sin(360^\circ + x)$ $= \sin x$	$\checkmark + \checkmark \sin x$ (2)
5.1.2	$x\text{-coordinate} = \sqrt{(\sqrt{13})^2 - (-3)^2}$ $= -2$ $\tan x = \frac{-3}{-2}$ $= \frac{3}{2}$ OR/OF $x\text{-coordinate} = \sqrt{(\sqrt{13})^2 - (3)^2}$ $= 2$ $\tan x = \frac{3}{2}$	$\checkmark\checkmark$ substitution \checkmark method $\checkmark\checkmark$ substitution \checkmark method (3)
5.1.3	$\cos(180^\circ + x)$ $= -\cos x$	$\checkmark - \checkmark \cos x$ (2)
5.2	$\frac{\cos(90^\circ + \theta)}{\sin(\theta - 180^\circ) + 3\sin(-\theta)}$ $= \frac{-\sin \theta}{\sin(-(180^\circ - \theta)) - 3\sin \theta}$ $= \frac{-\sin \theta}{-\sin \theta - 3\sin \theta}$ $= \frac{-\sin \theta}{-4\sin \theta}$ $= \frac{1}{4}$	$\checkmark - \sin \theta$ $\checkmark - 3\sin \theta$ $\checkmark - \sin \theta$ \checkmark simplification \checkmark answer (5)


5.3	$(\cos x + 2\sin x)(3\sin 2x - 1) = 0$ $\cos x + 2\sin x = 0$ or $3\sin 2x - 1 = 0$ $\tan x = -\frac{1}{2}$ $\sin 2x = \frac{1}{3}$ $\text{ref } \angle = 26,565\dots^\circ$ $\text{ref } \angle = 19,471\dots^\circ$ $x = 153,43^\circ + k.180^\circ; k \in \mathbb{Z}$ $x = 9,74^\circ + k.180^\circ; k \in \mathbb{Z}$ OR/OF or $x = 153,43^\circ + k.360^\circ; k \in \mathbb{Z}$ $x = 80,26^\circ + k.180^\circ;$ $k \in \mathbb{Z}$ or $x = 333,43^\circ + k.360^\circ; k \in \mathbb{Z}$	✓ both equations ✓ $\tan x = -\frac{1}{2}$ ✓ $\sin 2x = \frac{1}{3}$ ✓ $x = 153,43^\circ$ OR $x = 153,43^\circ \& 333,43^\circ$ ✓ $x = 9,74^\circ \& 80,26^\circ$ ✓ $+ k.180^\circ; k \in \mathbb{Z}$
		(6)
5.4.1	$\text{LHS} = \cos(x+y) \cdot \cos(x-y)$ $= [\cos x \cdot \cos y - \sin x \cdot \sin y][\cos x \cdot \cos y + \sin x \cdot \sin y]$ $= \cos^2 x \cdot \cos^2 y - \sin^2 x \cdot \sin^2 y$ $= (1 - \sin^2 x)(1 - \sin^2 y) - \sin^2 x \cdot \sin^2 y$ $= 1 + \sin^2 x \cdot \sin^2 y - \sin^2 x - \sin^2 y - \sin^2 x \cdot \sin^2 y$ $= 1 - \sin^2 x - \sin^2 y = \text{RHS}$	✓ expansion ✓ simplification ✓ square identity ✓ product
		(4)
5.4.2	$1 - \sin^2 45^\circ - \sin^2 15^\circ$ $= \cos(45^\circ + 15^\circ) \cdot \cos(45^\circ - 15^\circ)$ $= \cos 60^\circ \cdot \cos 30^\circ$ $= \left(\frac{1}{2}\right)\left(\frac{\sqrt{3}}{2}\right)$ $= \frac{\sqrt{3}}{4}$ OR/OF	✓ identifying x and y ✓ substitution ✓ answer
		(3)

5.5.1	$16 \sin x \cdot \cos^3 x - 8 \sin x \cdot \cos x$ $= 8 \sin x \cdot \cos x (2 \cos^2 x - 1)$ $= 4 \sin 2x (\cos 2x)$ $= 2 \sin 4x$ <p>OR/OF</p> $16 \sin x \cdot \cos^3 x - 8 \sin x \cdot \cos x$ $= 16 \cos^2 x \left(\frac{1}{2} \sin 2x \right) - 8 \left(\frac{1}{2} \sin 2x \right)$ $= 8 (2 \cos^2 x - 1) \left(\frac{1}{2} \sin 2x \right)$ $= 4 \sin 2x \cdot \cos 2x$ $= 2 \sin 4x$	<p>✓ factorisation</p> <p>✓ $4 \sin 2x$ ✓ $\cos 2x$</p> <p>✓ double angle</p> <p>(4)</p> <p>✓ factorisation</p> <p>✓ $4 \sin 2x$ ✓ $\cos 2x$</p> <p>✓ double angle</p> <p>(4)</p>
5.5.2	$16 \sin x \cdot \cos^3 x - 8 \sin x \cdot \cos x = 2 \sin 4x$ <p>Minimum at $x = 67,5^\circ$</p>	<p>✓ answer</p> <p>(1)</p>
		[30]

QUESTION/VRAAG 6




6.1	180°	✓ answer (1)
6.2.1	$k = \sqrt{3} = 1,73$	✓ answer (1)
6.2.2	$B(-120^\circ; \sqrt{3})$	✓ $x = -120^\circ$ (1)
6.3	Range of g : $y \in [-2; 2]$ Range of $2g(x)$: $y \in [-4; 4]$ OR/OF <div style="border: 1px solid black; padding: 5px; display: inline-block;">ANSWER ONLY: Full marks</div> Range of g : $-2 \leq y \leq 2$ Range of $2g(x)$: $-4 \leq y \leq 4$	✓ $y \in [-2; 2]$ ✓ answer (2) ✓ $-2 \leq y \leq 2$ ✓ answer (2)
6.4	$x \in [-65^\circ; -5^\circ]$ OR/OF $-65^\circ \leq x \leq -5^\circ$	✓✓ $x \in [-65^\circ; -5^\circ]$ (2) ✓✓ $-65^\circ \leq x \leq -5^\circ$ (2)
6.5	$\sin x \cdot \cos x = p$ $4 \sin x \cdot \cos x = 4p$ $2 \sin 2x = 4p$ $4p = \pm 2$ $\therefore p = -\frac{1}{2} \text{ or } \frac{1}{2}$ <div style="border: 1px solid black; padding: 5px; display: inline-block;">ANSWER ONLY: Full marks</div>	✓ $2 \sin 2x = 4p$ ✓ $4p = \pm 2$ ✓ answers (3)
[10]		

QUESTION/VRAAG 7

7.1	$AD^2 = AB^2 + BD^2$ $AD^2 = (\sqrt{5}p)^2 + (2p)^2$ $AD^2 = 9p^2$ $AD = 3p$	✓ substitution in Pythagoras ✓ answer (2)
7.2	$\frac{CD}{\sin(135^\circ - x)} = \frac{3p}{\sin x}$ $CD = \frac{3p \sin(135^\circ - x)}{\sin x}$ $CD = \frac{3p(\sin 135^\circ \cos x - \cos 135^\circ \sin x)}{\sin x}$ $CD = \frac{3p(\sin 45^\circ \cos x + \cos 45^\circ \sin x)}{\sin x}$ $CD = \frac{3p\left(\frac{\sqrt{2}}{2} \cos x + \frac{\sqrt{2}}{2} \sin x\right)}{\sin x}$ $CD = \frac{3p\left(\frac{\sqrt{2}}{2}\right)(\cos x + \sin x)}{\sin x}$ $CD = \frac{3p(\sin x + \cos x)}{\sqrt{2} \sin x}$	✓ correct use of sine rule ✓ $135^\circ - x$ ✓ compound angle ✓ special values ✓ factorisation (5)


QUESTION/VRAAG 8

8.1



8.1.1	$\hat{P} = 116^\circ$ [opp \angle s of cyclic quad/teenoorst. \angle e van kvh]	✓ S ✓ R (2)
8.1.2	$\hat{M}_1 + 64^\circ = 90^\circ$ [\angle in semi-circle/ \angle in halwe sirkel] $\hat{M}_1 = 26^\circ$	✓ R ✓ S (2)
8.1.3	$\hat{O}_1 = 52^\circ$ [\angle at centre = 2 x \angle at circumference/midpts. \angle = 2 x omtreks. \angle]	✓ S ✓ R (2)

8.2




8.2.1	Midpt theorem/ <i>Midpt. Stelling</i> OR/OF Converse prop intercept theorem	✓ R ✓ R (1)
8.2.2	$BG = 2DE$ or $6x - 2$ [<i>Midpt theorem/Midpt. stelling</i>] $BG = 6x - 2$ $\frac{GH}{BG} = \frac{FC}{BF}$ [<i>line one side of Δ OR</i> <i>prop theorem; $FG \parallel CH$ /</i> <i>lyn een sy v. Δ]</i> $\frac{x+1}{6x-2} = \frac{1}{4}$ $4x + 4 = 6x - 2$ $2x = 6$ $x = 3$ OR/OF	✓ S ✓ R ✓ S ✓ R ✓ equation into x ✓ answer (6)

	$\frac{BF}{FC} = \frac{BG}{GH}$ <p>[line \parallel one side of Δ OR prop theorem; $FG \parallel CH$ / <i>lyn \parallel een sy v. Δ</i></p> $\frac{AE}{AG} = \frac{DE}{BG}$ <p>[$\triangle ADE \parallel \triangle ABG$]</p> $BG = 4x + 4$ $\frac{1}{2} = \frac{3x-1}{4x+4}$ $\therefore 4x + 4 = 6x - 2$ $\therefore x = 3$	<p>✓ S ✓ R</p> <p>✓ S ✓ R</p> <p>✓ equation into x</p> <p>✓ answer</p> <p>(6)</p>
		[13]


QUESTION/VRAAG 9

9.1




9.1.1	<p>Construction: Draw OA and OB In $\triangle ADO$ and $\triangle BDO$ $OA = OB$ [radii/radiusse] $OD = OD$ [common side/gemeenskaplike sy] $AD = DB$ [given/gegee] $\therefore \triangle ADO \equiv \triangle BDO$ [S;S;S] ADB is a straight line $\therefore \hat{D}_1 = \hat{D}_2$ $\therefore OD \perp AB$</p> <p>OR/OF Construction: Draw OA and OB In $\triangle ADO$ and $\triangle BDO$ $AD = DB$ [given/gegee] $\hat{A} = \hat{B}$ [\angles opp; \angles sides / \anglee teenoor gelyke sye] $OA = OB$ [radii/radiusse] $\therefore \triangle ADO \equiv \triangle BDO$ [S;\angle;S] ADB is a straight line $\therefore \hat{D}_1 = \hat{D}_2$ $\therefore OD \perp AB$</p> <p>$\triangle ADO \equiv \triangle BDO$ [\angles on a str line/\anglee op 'n reguitlyn]</p>	<p>✓ construction</p> <p>✓ first pair of sides ✓ other 2 pairs ✓ R</p> <p>✓ R</p> <p>(5)</p> <p>✓ construction</p> <p>✓ first pair of sides</p> <p>✓ other 2 pairs ✓ R</p> <p>✓ R</p> <p>(5)</p>
-------	--	---

9.2



9.2.1	$\hat{O}TG = 90^\circ$ $\hat{O}BG = 90^\circ$ $\therefore \hat{O}TG = \hat{O}BG = 90^\circ$ $\therefore OTBG$ is a cyclic quadrilateral	[line from centre to midpt of chord/ <i>midpt. sirkel; midpt. koord</i>] [tan \perp radius/ <i>raaklyn \perp radius</i>] [line subtends equal \angle s OR converse \angle s in the same segment/ <i>lyn onderspan gelyke \anglee</i>]	✓ S ✓ R ✓ S ✓ R ✓ R	(5)
9.2.2	$\hat{S} = \hat{B}TG$ But $\hat{B}TG = \hat{G}OB$ $\hat{G}OB = \hat{S}$	[corresp \angle s; $GF \parallel PS$ / <i>ooreenk. \angles; $GF \parallel PS$] [\angles in the same segment/\anglee in dies. <i>sirkelsegment</i>] </i>	✓ S ✓ R ✓ S ✓ R	(4)
[14]				

QUESTION/VRAAG 10

10.1	$\hat{P}_1 = \hat{Q}_1$ $\hat{S}_1 = \hat{Q}_1 + \hat{Q}_2$ $\therefore \hat{S}_1 = \hat{P}_1 + \hat{Q}_2$ $\hat{T}_2 = \hat{R}_2 + \hat{Q}_2$ but $\hat{P}_1 = \hat{R}_2$ $\hat{T}_2 = \hat{P}_1 + \hat{Q}_2$ $\therefore \hat{S}_1 = \hat{T}_2 = \hat{P}_1 + \hat{Q}_2$	[tan-chord theorem/ <i>∠ tussen raaklyn en koord</i>] [ext \angle of cyclic quad/ <i>buite \angle v. kvh</i>] [ext \angle of Δ / <i>buite \angle v. Δ</i>] [given/ <i>gegee</i>]	✓ S ✓ S / R ✓ S ✓ S
10.2	In ΔASD and ΔACR $\hat{A} = \hat{A}$ $\hat{S}_1 = \hat{T}_2$ $\hat{T}_2 = \hat{C}_2$] $\therefore \hat{S}_1 = \hat{C}_2$ $\hat{D}_1 = \hat{R}_1$ $\Delta ASD \parallel \Delta ACR$ $\therefore \frac{AD}{AR} = \frac{AS}{AC}$ OR/OF	[common \angle / <i>gemeenskaplike \angle</i>] [proven/ <i>reeds bewys</i>] [alt \angle s; $QS \parallel CA$ / <i>verw. \anglee; $QS \parallel CA$</i>] [sum of \angle s in Δ / <i>\angle v. Δ</i>] [corresponding sides in proportion/ <i>ooreenstemmende sy in dies. verhouding</i>]	✓ identifying Δ 's ✓ S ✓ S / R ✓ S ✓ S

(4)

(5)

	<p>In $\triangle ASD$ and $\triangle ACR$</p> <p>$\hat{A} = \hat{A}$ [common \angle/gemeenskaplike \angle]</p> <p>$\hat{S}_1 = \hat{T}_2$ [proven/gegee]</p> <p>$\hat{T}_2 = \hat{C}_2$ [alt \angles; $QS \parallel CA$/verw. \anglee; $QS \parallel CA$]</p> <p>$\therefore \hat{S}_1 = \hat{C}_2$</p> <p>$\triangle ASD \parallel \triangle ACR$ [\angle; \angle; \angle]</p> <p>$\therefore \frac{AD}{AR} = \frac{AS}{AC}$ [corresponding sides in proportion/ ooreenstemmende sy in dies. verhouding]</p>	<p>✓ identifying \triangle's</p> <p>✓ S</p> <p>✓ S/R</p> <p>✓ S</p> <p>✓ R</p>
10.3	<p>$\frac{AS}{AC} = \frac{SD}{CR}$ [$\triangle ASD \parallel \triangle ACR$]</p> <p>$\therefore AS = \frac{AC \times SD}{CR}$</p> <p>$\frac{AS}{AR} = \frac{CT}{CR}$ [line \parallel one side of \triangle OR prop theorem; TS \parallel CA/lyn \parallel een sy v. \triangle]</p> <p>$\therefore AS = \frac{AR \times CT}{CR}$</p> <p>$\therefore \frac{AC \times SD}{CR} = \frac{AR \times CT}{CR}$</p> <p>$\therefore AC \times SD = AR \times CT$</p>	<p>✓ S</p> <p>✓ S ✓ R</p> <p>✓ equating</p>
		(5)
		(4)
		[13]

TOTAL/TOTAAL: 150