

MATH 6304 - Theory of Matrices

Joel Sleeba
University of Houston
joelsleeba1@gmail.com

September 4, 2025

Contents

Contents	1
1 Introduction	2
1.1 Review of Linear Algebra	3
1.2 Eigenvalues and Eigenvectors	4
1.3 Similarity	4
1.4 Diagonazability	4
1.5 Warm Up	6
1.6 Simultaneous diagonalization	6

26/08/2025

Office Hours : Tuesday 10 - 11 AM, Wednesday 1 - 2 PM

Chapter 1

Introduction

Let A be a $m \times n$ matrix and D a diagonal $n \times n$ matrix with entries d_1, d_2, \dots, d_n ,

$$A = \begin{bmatrix} | & | & \cdots & | \\ a_1 & a_2 & \cdots & a_n \\ | & | & \cdots & | \end{bmatrix}$$

Then

$$AD = \begin{bmatrix} | & | & \cdots & | \\ d_1 a_1 & d_2 a_2 & \cdots & d_n a_n \\ | & | & \cdots & | \end{bmatrix}$$

and if

$$B = \begin{bmatrix} - & b_1 & - \\ - & b_2 & - \\ & \vdots & \\ - & b_n & - \end{bmatrix}$$

then

$$DB = \begin{bmatrix} - & d_1 b_1 & - \\ - & d_2 b_2 & - \\ & \vdots & \\ - & d_n b_n & - \end{bmatrix}$$

Do the same for upper triangular matrices and add some context to the multiplications.

Notice that every time you left multiply, you play with column, and when you right multiply, you play with the rows.

Exercise 1.0.1. Let A be an $n \times n$ matrix over \mathbb{C} . Let $\omega = e^{\frac{2\pi i}{n}}$. Then prove that

$$A' = \frac{1}{n} \sum_{k=0}^n (U^*)^k A U^k$$

preserve all the diagonal entries of A and kills the rest of entries. That is $A' = \text{Diag}(A)$

1.1 Review of Linear Algebra

- Rank-Nullity Theorem
- Orthogonality
- Orthogonal projection is the closest point on the subspace from the given vector.

28/08/2025

Definition 1.1.1. If $A \in M_n$, $A = (a_{i,j})_{i,j=1}^n$, we let

$$\text{trace}(A) = \sum_{j=1}^n a_{jj}$$

The determinant is

$$\det(A) = \sum_{\sigma \in S_n} \text{sgn}(\sigma) a_{1,\sigma(1)} a_{2,\sigma(2)} \cdots a_{n,\sigma(n)}$$

Remark 1.1.1. If $A = [a_1 a_2 \dots a_n]$, then $\det(A) = f(a_1, a_2, \dots, a_n)$ is the only function that is linear in each a_i , alternating (swapping columns doesn't alter the value), and normalized ($\det(I) = 1$).

This is useful to show that for $A, B \in M_n$, $\det(AB) = \det(A)\det(B)$.

Moreover if $A = \begin{bmatrix} B & C \\ 0 & D \end{bmatrix}$, then $\det(A) = \det(B)\det(D)$.

We also have

$$\det(A) = \sum_{i,j=1}^n (-1)^{i+j} a_{i,j} \det(A_{i,j})$$

Where $A_{i,j}$ is the submatrix with i th row and j th column removed from A .

1.2 Eigenvalues and Eigenvectors

Definition 1.2.1. Eigenvalue, Eigenvector, Spectrum of a matrix

1.3 Similarity

Definition 1.3.1. A matrix $B \in M_n$ is similar to $A \in M_n$, if there is an invertible $S \in M_n$ such that $B = S^{-1}AS$. This defines an equivalence relation.

Theorem 1.3.1. If $A, B \in M_n$ are similar. Then their characteristic polynomial $P_A = P_B$.

Remark 1.3.1. characteristic polynomial is not characteristic upto similarity, because

$$A = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \text{ and } B = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$$

02/09/2025

1.4 Diagonalizability

Can we find conditions for diagonalizability.

Theorem 1.4.1. Let $A \in M_n(\mathbb{C})$, $p_A(t) = \prod_{j=1}^n (t - \lambda_j)$, and $\lambda_i \neq \lambda_j$ for $j \neq i$, then A is diagonalizable.

Proof. We'll show that there's a linearly independent set of n eigenvectors. Let $x_j \in \mathbb{C}^n$ such that $Ax_j = \lambda_j x_j$. If $\{x_1, x_2, \dots, x_n\}$ were linearly dependent, then there is a linear combination

$$\alpha_1 x_{j_1} + \alpha_2 x_{j_2} + \dots + \alpha_r x_{j_r} = 0$$

with $r \leq n$, and all $\alpha_j \neq 0$. Let r be smallest such $r \leq n$, and assume with possible renumbering that $j_i = i$. Then applying A to the linear combination gives us

$$A(\alpha_1 x_1 + \alpha_2 x_2 + \dots + \alpha_n x_n) = \alpha_1 \lambda_1 x_1 + \alpha_2 \lambda_2 x_2 + \dots + \alpha_n \lambda_n x_n = 0$$

multiplying the previous equation with λ_r and then subtracting gives us

$$\alpha_1 (\lambda_1 - \lambda_r) x_1 + \alpha_2 (\lambda_2 - \lambda_r) x_2 + \dots + \alpha_r (\lambda_r - \lambda_r) x_r = 0$$

which contradicts the minimality of r . □

Unfortunately this is just a sufficient condition, as it excludes the following matrix.

$$\begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

Definition 1.4.1. If for $A \in M_n(\mathbb{C})$,

$$p_A(t) = (t - \lambda_1)^{m_1} (t - \lambda_2)^{m_2} \dots (t - \lambda_r)^{m_r}$$

then we say that λ_j has algebraic multiplicity m_j . We call $\text{null}(\lambda_j I - A)$, the geometric multiplicity of λ_j

Lemma 1.4.1. If $A \in M_n$ has eigenvalue λ , and $p_A(t) = (t - \lambda)^m q(t)$, with $q(\lambda) \neq 0$, then $r = \text{null}(\lambda I - A) \leq m$

Proof. Choose a basis $\{x_1, x_2, \dots, x_r\}$ of eigenvectors, spanning $E_\lambda = \{x \in \mathbb{C}^n : Ax = \lambda x\}$. Complete it to a basis $\{x_1, x_2, \dots, x_n\}$ of \mathbb{C}^n . Let $S = [x_1, x_2, \dots, x_n]$.

Then $AS = [\lambda x_1, \lambda x_2, \dots, \lambda x_r, y_{r+1}, \dots, y_n]$ with some vectors y_{r+1}, \dots, y_n . Then $S^{-1}AS = \text{verify}$, and we get

$$\begin{aligned} \det(tI - A) &= \det(tI - S^{-1}AS) \\ &= (t - \lambda)^r \det(tI - C) \end{aligned}$$

Thus we conclude that algebraic multiplicity of λ is at least equal to r . □

Remark 1.4.1. See that the sum of all the algebraic multiplicity of the eigenvalues of $A \in M_n(\mathbb{C})$ is n .

Theorem 1.4.2. The matrix $A \in M_n(\mathbb{C})$ is diagonalizable if and only if the algebraic and geometric multiplicities are equal for each eigenvalue.

Proof. We note that given two eigenvalues $\lambda_j \neq \lambda_k$, then their eigenspaces E_i, E_j intersect trivially. Thus if $\{v_1, v_2, \dots, v_{r_1}\}$ and $\{u_1, u_2, \dots, u_{r_2}\}$ form a basis for E_{λ_1} and E_{λ_2} respectively, then $\{v_1, v_2, \dots, v_{r_1}, u_1, u_2, \dots, u_{r_2}\}$ is linearly independent. Iterating this way, we get a basis for $E_1 + E_2 \dots + E_n$ with dimension $r = \sum_{i=1}^k r_i$.

If algebraic and geometric multiplicities equal then $r = n$, and we have a basis of eigenvectors. Otherwise if $r < n$, then we do not have such a basis of eigenvectors. And since existence of a basis of eigenvectors characterizes diagonalizability, this characterizes diagonalizability. □

Next lecture, we'll look when multiple matrices can be simultaneously diagonalizable with the same S matrix.

1.5 Warm Up

Assume we know that A is diagonalizable. Let $p_0, p_1, p_2, \dots, p_n \in \mathbb{C}$ and consider

$$B := P(A) = p_0 I + p_1 A + p_2 A^2 + \dots + p_n A^n$$

Is B diagonalizable?

Proof. Yes. Because if $A = S^{-1}DS$, then $A^n = S^{-1}D^nS$, and therefore

$$B = S^{-1}(p_0 I + p_1 D + p_2 D^2 + \dots + p_n D^n)S$$

□

1.6 Simultaneous diagonalization

Theorem 1.6.1. *Let A, B be diagonalizable. Then $AB = BA$ if and only if they are simultaneously diagonalizable by the same S .*

Proof. Let $D_A = S^{-1}AS$, and $B' = S^{-1}BS$, where D_A is a diagonal matrix. Without loss of generality, assume that common eigenvalues appear together in D_A . If not choose S with an additional permutation of the rows.

Assuming $AB = BA$, we get

$$\begin{aligned} D_A B' &= S^{-1} A S S^{-1} B S \\ &= S^{-1} A B S \\ &= S^{-1} B A S \\ &= S^{-1} B S S^{-1} A S \\ &= B' D_A \end{aligned}$$

If $B' = [b'_{i,j}]_{i,j=1}^n$, then by $D_A B' = B' D_A$, from the diagonal structure of D_A , we get

$$\tilde{\lambda}_i b'_{i,j} = b'_{i,j} \tilde{\lambda}_j$$

where $\tilde{\lambda}_i$ is the i -th diagonal entry on D_A . So, we have

$$(\tilde{\lambda}_i - \tilde{\lambda}_j) b'_{i,j} = 0$$

which shows that if $\tilde{\lambda}_i \neq \tilde{\lambda}_j$, then $b'_{i,j} = 0$. Thus we get that

$$B' = \begin{bmatrix} B'_1 & & & \\ & B'_2 & & \\ & & \ddots & \\ & & & B'_r \end{bmatrix}$$

Since B and B' are diagonalizable, so is each B'_i . Taking matrices T_1, T_2, \dots, T_r that diagonalize B'_1, B'_2, \dots, B'_r respectively, let

$$T = \begin{bmatrix} T_1 & & & \\ & T_2 & & \\ & & \ddots & \\ & & & T_r \end{bmatrix}$$

Then,

$$T^{-1}BT = \begin{bmatrix} T_1^{-1}B'_1T_1 & & & \\ & T_2^{-1}B'_2T_2 & & \\ & & \ddots & \\ & & & T_r^{-1}B'_rT_r \end{bmatrix} = \begin{bmatrix} D'_1 & & & \\ & D'_2 & & \\ & & \ddots & \\ & & & D'_r \end{bmatrix}$$

where each D'_i is a diagonal block. Also,

$$T^{-1}D_AT = \begin{bmatrix} T_1^{-1}\lambda_1IT_1 & & & \\ & T_2^{-1}\lambda_2IT_2 & & \\ & & \ddots & \\ & & & T_r^{-1}\lambda_rIT_r \end{bmatrix} = D_A$$

This implies $D_A = T^{-1}S^{-1}AST$, and $D_B = T^{-1}S^{-1}BST$ are both diagonal.

Converse is left as an exercise

□

Next, we consider simultaneous diagonalization for a family of matrices.

Definition 1.6.1. A family $F \subset M_n$ is a commuting family if for each $A, B \in F$, $AB = BA$.

Definition 1.6.2. A subspace $W \subset \mathbb{C}^n$ is called an A -invariant subspace for some $A \in M_n$ if $Aw \in W$ for all $w \in W$. If $F \subset M_n$, then W is called F -invariant if for each $A \in F$, W is A -invariant.

Lemma 1.6.1. If $W \subset \mathbb{C}^n$ is A -invariant for some $A \in M_n$, and suppose that $\dim(W) \geq 1$, then there is an $x \in W \setminus \{0\}$ such that $Ax = \lambda x$.

Proof. We consider $B := A|_W$. Then $B : W \rightarrow W$ has an eigenvector since it has atleast one eigenvalue by the fundamental theorem of algebra. □

Lemma 1.6.2. If $F \subset M_n$ is a commuting family, then there exists an $x \in \mathbb{C}^n$ such that for each $A \in F$, $Ax = \lambda_A x$.

Proof. Choose W to be an F -invariant subspace of minimum, non-zero dimension. Existence of W is guaranteed since we can choose $W = \mathbb{C}^n$.

Next, we show that any $x \in W \setminus \{\mathbf{0}\}$ is an eigenvector for each $A \in F$. Assume this is not true. Then there is a $y \in W \setminus \{\mathbf{0}\}$, and an $A \in F$, such that $Ay \notin \mathbb{C}y$. Since W is A -invariant by the setup, by previous lemma, we get that there is a $x \in W \setminus \{\mathbf{0}\}$ such that $Ax = \lambda_x x$ for some $\lambda \in \mathbb{C}$.

Let $W_0 = \{z \in W : Az = \lambda z\}$. By $y \notin W_0$, we get that $W_0 \neq W$. But for any $B \in F$, by invariance of W_0 , $Bx \in W$, and for $u \in W_0$,

$$A(Bu) = B(Au) = \lambda Bu$$

We observe $Bu \in W_0$, thus B maps W_0 to W_0 , so W_0 is F -invariant. We have derived a contradiction with the minimality of W . \square

Remark 1.6.1. This implies that commuting families have at least one common eigenvector

Definition 1.6.3. A simultaneously diagonalizable family is a family $F \subset M_n$ such that there exists $S \in M_n$ for which $S^{-1}AS$ is diagonal for each $A \in F$

Theorem 1.6.2. Let $F \subset M_n$ be a family of diagonalizable matrices, then F is a commuting family if and only if it is simultaneously diagonalizable.

Proof. \square