

MATH 6321**Theory of Functions of a Real Variable
Spring 2025**

First name: _____ Last name: _____

Points:

--

Assignment 2, due Thursday, February 6, 10am

Please staple this cover page to your homework. When asked to prove something, make a careful step-by-step argument. You can quote anything we covered in class in support of your reasoning.

Problem 1

Let $1 < p < \infty$. Prove that for $f, g \in L^p(\mathbb{R})$, if $\|f\|_p = \|g\|_p = 1$ and $f \neq g$ (they differ on a set of positive measure), then $\|f + g\|_p < 2$.

Problem 2

Consider $C([0; 1])$, equipped with the sup-norm. Show that the affine subspace

$$M = \{f \in C([0; 1]) : \int_0^1 f(t) dt = 1\}$$

is a closed and convex subset of $C([0; 1])$ which does not contain an element of minimal norm.

Problem 3

Consider $L^1([0; 1])$ (as space of equivalence classes of functions that are identical up to sets with vanishing Lebesgue measure m) and

$$M = \{f \in L^1([0; 1]) : \int_0^1 f dm = 1\}$$

and show that M has infinitely many elements of minimal norm (viewed as functions, they differ on sets of non-zero measure).

Problem 4

Consider $C([0; 1])$, equipped with the sup-norm. Let for $n \in \mathbb{N}$,

$$X_n = \{f \in C([0; 1]) : \text{there is } x \in [0; 1] \text{ s.t. for each } y \in [0; 1] : |f(x) - f(y)| \leq \frac{1}{n} |x - y|\}$$

Prove that each X_n has empty interior. Use this to prove that there is a G_δ set of nowhere differentiable functions in $C([0; 1])$. Hint: Why is each X_n closed? To show X_n has empty interior, take $f \in X_n$ and use uniform continuity to approximate it with a piecewise linear, continuous function g . Next, consider $h(x) = g(x) + \sin(Nx)$ or functions that have similar oscillatory behavior.