

Numerical Analysis Qualifier

prepared by

Jing Li & Lothar Reichel

Kent State University

August, 2011

INSTRUCTIONS: Do any (and only) 8 of the following 10 problems.

Throughout this exam $\|\cdot\|$ denotes the Euclidean vector norm or the associated induced matrix norm.

1. Floating-point arithmetic: By the periodicity of the cosine function, the following relation should hold for all integers k :

$$\cos\left(\frac{\pi}{4} + 2k\pi\right) = \frac{1}{\sqrt{2}}.$$

Yet when the left-hand side is evaluated in MATLAB with $k=10^8$, we obtain

$$\left|\cos\left(\frac{\pi}{4} + 2 \cdot 10^8 \pi\right) - \frac{1}{\sqrt{2}}\right| = 8.2 \cdot 10^{-8}.$$

Explain.

2. Stability: Let X and Y be normed vector spaces. Let $\tilde{f} : X \rightarrow Y$ denote an algorithm for the solution of the problem $f : X \rightarrow Y$.

- (a) What does it mean that the algorithm \tilde{f} is *stable* for each $x \in X$?
- (b) What does it mean that the algorithm \tilde{f} is *backward stable* for each $x \in X$?

3. Gaussian elimination: Consider the problem of solving $Ax = b$, where $A \in \mathbb{R}^{n \times n}$ is nonsingular and $x, b \in \mathbb{R}^n$, by Gaussian elimination with partial pivoting (GEPP).

- (a) Let $PA = LU$, where P is a permutation matrix. Let \tilde{L} and \tilde{U} be the computed factors by GEPP. Then

$$\tilde{L}\tilde{U} = PA + E,$$

where the matrix E satisfies

$$\frac{\|E\|}{\|L\|\|U\|} = \mathcal{O}(\varepsilon_{\text{machine}})$$

and $\varepsilon_{\text{machine}}$ denotes machine epsilon. Is GEPP backward stable? Justify your answer.

- (b) Let $x = A^{-1}b$ and let \tilde{x} be the approximate solution computed with GEPP. Give a bound for $\|x - \tilde{x}\|$. Justify your bound.
4. The singular value decomposition:
- (a) What is the singular value decomposition of a matrix $A \in \mathbb{R}^{m \times n}$, $m \geq n$? Describe the matrices involved.
 - (b) Express the norm of A and the condition number of A in terms of the singular values of A .
 - (c) Give the matrix of rank one that best approximates A .
5. Interpolation: The error formula for polynomial interpolation to a smooth function $f(t)$ by a polynomial of degree at most $n - 1$ at n distinct points $t_1 < \dots < t_n$ is given by

$$f(t) - P_{n-1}(t) = \frac{f^{(n)}(\theta(t))}{n!} (t - t_1) \cdots (t - t_n).$$

- (a) Prove the above formula.
 - (b) What are the zeros of the Chebyshev polynomial T_n and what is their significance in the context of polynomial interpolation?
 - (c) Let all nodes t_j live in the interval $[-1, 1]$ and let $f(t) = \exp(t)$. Will the interpolating polynomials P_{n-1} converge to f on the interval $[-1, 1]$ as the number of interpolation points is increased? Justify your answer.
6. Orthogonal Polynomials: Define the inner product

$$(f, g) = \int_{-1}^1 f(x)g(x)w(x) dx, \tag{1}$$

where $w(x)$ is a positive weight function, and consider the family of orthogonal polynomials p_0, p_1, p_2, \dots with respect to this inner product. Here p_j is of degree j and has leading coefficient one.

- (a) Give the form of the three-term recurrence relation that these polynomials satisfy.
 - (b) Prove the existence of the three-term recurrence relation. Give expressions for the coefficients in the recurrence relation.
7. Gauss quadrature: Let w be the weight function of the inner product (1).

- (a) Describe the Gauss quadrature rule for approximating the integral

$$\int_a^b f(x)w(x)dx \approx \sum_{k=1}^n w_k f(x_k).$$

How are the nodes related to the orthogonal polynomials of problem 6?

- (b) Prove that this rule is exact for all polynomials of degree strictly less than $2n$.
8. Newton's method: Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a nonlinear differentiable function. The equation $f(x) = 0$ can be solved by Newton's method.
- (a) Describe Newton's method for the solution of $f(x) = 0$.
- (b) Define quadratic convergence of an iterative method.
- (c) Show that Newton's method yields quadratic convergence.
9. Gram-Schmidt Orthogonalization: Consider the $m \times n$ matrix A with linearly independent columns and $m \geq n$. Describe the modified Gram-Schmidt procedure for orthogonalizing the columns of A .
10. Projectors:
- (a) Give the defining equation for a projector P .
- (b) Show that $I - P$ also is a projector.
- (c) Show that $I - P$ projects onto the nullspace of P .
- (d) What is an orthogonal projector? How do the range and nullspace of an orthogonal projector relate?
- (e) Give a formula for the orthogonal projector onto the range of $A \in \mathbb{R}^{m \times n}$, $m > n$. The matrix A is assumed to have linearly independent columns.