

# Tutorial – Mathematics for Social Scientists

Winter semester 2024/25

Introduction to Probability

# To do

- short extrema recap
- weekly recap
- real world applications
- hands on practice
- questions

# Chapter 8 | Extrema - Recap

# Extrema Recap


**Does  $f(x) = 3x + 2$  have extrema?**

→ No,  $f(x)$  increases in a strictly monotonic fashion and if we inspect it in its entirety, it does not have extrema

**... what about  $f(x) = 3x + 2$  for  $x \in [1, 2]$ ?**

→ Yes! Once we zoom into an interval and only inspect parts of  $f(x)$ , we can find extrema!

→  $f(x)$  then has a global min at  $P(1 | 5)$  and a global max at  $P(2 | 8)$



# Endpoint Extrema

## Endpoint extrema are valid extrema

→ they occur only(!) at the boundary points of an interval – if we inspect  $f(x)$  at such an interval

Endpoints can be **both** local and global extrema

→ an extreme point is considered 'local' when we find no min or max within the direct neighbourhood of our extreme point that is higher/lower in magnitude

→ it is considered 'global', when it is the highest/lowest in magnitude over the defined domain!

The defined **domain** may be the function in its entirety – or just the specified interval!

# Chapter 9 | An Introduction to Probability

# Probability vocabulary

## Probability

- a measure of an event's likelihood

## Likelihood

- the plausibility of an event to occur

## Sample space

- a set of all possible outcomes

## Outcome

- a realized event of e.g., an experiment/ the DGP

## (Random) Event

- uncertain outcome, follows probability distribution

**Probability of an event**  $\Pr(e) = \frac{\# \text{ outcomes of interest}}{\# \text{ outcomes in sample space}}$

# Hands on – probability definitions

**Task:** Given one roll of a fair die, what is the probability that the number rolled is less than 3?

- what is an outcome?
- what is the sample space  $S$ ?
- what is the event  $e$ ?
- what is the  $\Pr(e)$ ?



# Hands on – probability definitions

## Solution:

- what is an outcome?  
→ one side of the die – a ‘number’
- what is the sample space  $S$ ?  
→  $S = \{1, 2, 3, 4, 5, 6\}$
- what is the event  $e$ ?  
→  $e = \{1, 2\}$
- what is  $\Pr(e)$ ?  
→  $\Pr(e) = \frac{\# \text{ outcomes of interest}}{\# \text{ outcomes in sample space}} = \frac{2}{6} = 0.\overline{33}$

# Hands On – Venn diagram refresher

**Task:** Draw the Venn Diagrams for the events A and B in the overall sample space S for:

- the complement of A  $A^C$  or  $\bar{A}$
- the subset of A and B  $A \subset B$
- the Union of A and B  $A \cup B$
- the intersection of A and B  $A \cap B$

**Hint:** You have seen all of these already in class...

# Hands On – Venn diagram refresher

**Solution:**

# Probability rules – joint probabilities

## Addition rule

- applies whenever an event is the union of two other events

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

## Multiplication rule

- applies whenever an event is the intersection of events A and B  
→ then events A and B must occur simultaneously

$$\begin{aligned} P(A \cap B) &= P(A) \cdot P(B|A) \\ &= P(B) \cdot P(A|B) \end{aligned}$$

# Addition and multiplication rule

# Probability rules

## Complement rule

- applies whenever we ‘flip’ an event

$$P(A') = 1 - P(A)$$

$$P(A) + P(A') = 1$$

## Conditional rule

- applies whenever an event needs another event to occur first!

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

# Independence, mutually exclusive, collectively exhaustive

## Independence

- two events A and B are considered **independent** if the **occurrence** of **one event** **does NOT depend** on the **occurrence** of the **other**
- $P(A) = P(A|B)$

## Mutually exclusive


- **mutually exclusive events** have **NO outcome** in **common** – they **CANNOT** occur together!  
→ also called **disjoint events**

- **Collectively exhaustive**

- the **union** of **all events** covers the **entire sample space** – at least one event in the sample space **MUST** occur

# Independence and mutually exclusive events

Which of these depicts independent events?



mutually exclusive:

$P(A \cap B) = 0$   
NO intersection  
NO outcomes in common!

Independence:

$$P(A) = P(A|B)$$

I can both throw a coin head's up AND I can assure you, I will have lunch!

$$P(A \cap B) = P(A) \cdot P(B)$$

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

$$P(A|B) = \frac{P(A) \cdot P(B)}{P(B)}$$



# Hands on – probability rules


**Task:** The sample space for throwing a die is  $S = \{1, 2, 3, 4, 5, 6\}$ . Suppose events A, B and C are defined as follows:

A = Getting an even number  $\{2, 4, 6\}$

B = Getting at least 5 =  $\{5, 6\}$

C = Getting at most 3 =  $\{1, 2, 3\}$

Find the probability of each of these events and its complement. Then, find the union, intersection and conditional probability of each pair of events.



AP Statistics Edition 2020 Princeton Review, p. 213


# Hands on – probability rules

## Solution:

- $A$  = Getting an even number  $\{2, 4, 6\}$
- $B$  = Getting at least 5 =  $\{5, 6\}$
- $C$  = Getting at most 3 =  $\{1, 2, 3\}$

## Probability:

- $P(A) = \frac{3}{6} = 0.5$
- $P(B) = \frac{2}{6} = 0.\bar{3}$
- $P(C) = \frac{3}{6} = 0.5$



AP Statistics Edition 2020 Princeton Review, p. 213

# Hands on – probability rules

## Solution:

- $A$  = Getting an even number  $\{2, 4, 6\}$
- $B$  = Getting at least 5  $= \{5, 6\}$
- $C$  = Getting at most 3  $= \{1, 2, 3\}$

## Complement:

$A'$  = Getting an odd number  $= \{1, 3, 5\}$


- $P(A') = \frac{3}{6} = 0.5 = 1 - P(A)$

$B'$  = Getting a number less than 5  $= \{1, 2, 3, 4\}$

- $P(B') = \frac{4}{6} = 0.\bar{6} = 1 - P(B)$

$C'$  = Getting a number larger than 3  $= \{4, 5, 6\}$

- $P(C') = \frac{3}{6} = 0.5 = 1 - P(C)$



AP Statistics Edition 2020 Princeton Review, p. 213

# Hands on – probability rules

## Solution:

- $A$  = Getting an even number  $\{2, 4, 6\}$
- $B$  = Getting at least 5 =  $\{5, 6\}$
- $C$  = Getting at most 3 =  $\{1, 2, 3\}$

## Union:

$(A \cup B)$  = Getting an even nr. or one  $\geq 5 = \{2, 4, 5, 6\}$


- $P(A \cup B) = \frac{4}{6} = 0.\bar{6}$

$(A \cup C)$  = Getting an even nr. or one  $\leq 3 = \{1, 2, 3, 4, 6\}$

- $P(A \cup C) = \frac{5}{6} = 0.8\bar{3}$

$(B \cup C)$  = Getting a nr. that is at most 3 or at least 5 or both =  $\{1, 2, 3, 5, 6\}$

- $P(B \cup C) = \frac{5}{6} = 0.8\bar{3}$



AP Statistics Edition 2020 Princeton Review, p. 213

# Hands on – probability rules

## Solution:

- $A$  = Getting an even number  $\{2, 4, 6\}$
- $B$  = Getting at least 5 =  $\{5, 6\}$
- $C$  = Getting at most 3 =  $\{1, 2, 3\}$

## Intersection:

$(A \cap B)$  = Getting an even nr. that is at least 5 =  $\{6\}$

- $P(A \cap B) = \frac{1}{6} = 0.1\bar{6}$


$(A \cap C)$  = Getting an even nr. that is at most 3 =  $\{2\}$

- $P(A \cap C) = \frac{1}{6} = 0.1\bar{6}$

$(B \cap C)$  = Getting a nr. that is at most 3 and at least 5 =  $\{\}$

- $P(B \cap C) = \frac{0}{6} = 0.00$

→ **B and C are mutually exclusive events!**



AP Statistics Edition 2020 Princeton Review, p. 213



# Hands on – probability rules

## Solution:

- $A$  = Getting an even number  $\{2, 4, 6\}$
- $B$  = Getting at least 5  $= \{5, 6\}$
- $C$  = Getting at most 3  $= \{1, 2, 3\}$

## Conditional event:

$(A|C)$  = Getting an even nr. that it is at most 3  $= \{2\}$


- $P(A|C) = \frac{1}{3} = 0.\bar{3}$

$(A|B)$  = Getting an even nr. given it is  $\geq 5 = \{6\}$

- $P(A|B) = \frac{1}{2} = 0.5$

$(B|C)$  = Getting at least 5 given that the nr. is at most 3  $= \emptyset$

- $P(B|C) = 0$



AP Statistics Edition 2020 Princeton Review, p. 213

# Probability rules

## Want even more practice?

- come up with your own scenarios and go to <https://www.cuemath.com/data/probability-rules/> to fact check your results! 😊

# Combinations and permutations

## Combinations

- Order **DOES NOT** matter!

$$\binom{n}{k} = \frac{n!}{k!(n-k)!} = n_{C_r}$$

→ use when you have to ‘**select**’  
k things out of n

## Permutations

- Order **DOES** matter!

$$P(n, k) = \frac{n!}{(n-k)!} = n_{P_r}$$

→ use when you have to  
‘**arrange**’ k things out of n



# Hands on – combinations and permutations

**Task:** Let there be a bag with three coloured balls (one red, one yellow, one blue), how many ways are there to draw two balls given:

- order does not matter?
- order does matter?

# Hands on – combinations and permutations

**Solution:** Let there be a bag with three coloured balls (one red, one yellow, one blue), how many ways are there to draw two balls given:

- order does not matter?

{(red & yellow), (red & blue), (yellow & blue)}

$$\binom{3}{2} = \frac{3!}{2!(3-2)!} = \frac{6}{2} = 3$$


- order does matter?

{(red & yellow), (yellow & red), (red & blue), (blue & red), (yellow & blue), (blue & yellow)}


$$P(3, 2) = \frac{3!}{(3-2)!} = \frac{6}{1} = 6$$

# Real world applications – molecules

**Task:** Let a linear hydrocarbon of nine singly bonded carbon atoms be given, at the end of which there is an OH group.



- a) How many **different isomers** can be obtained by **substituting two chlorine atoms**, if only one chlorine atom may be placed at each carbon atom?
- b) How many **different isomers** do you get if you use **one chlorine atom** and **one bromine atom** instead of two chlorine atoms?




→**NOTE:** The images do not exactly depict the molecule of the task 😊

→**Isomers:** molecules **constituted** of the **same number** of atoms **per element** – but **put together differently**

# Real world applications – molecules

**Task:** Let a linear hydrocarbon of nine singly bonded carbon atoms be given, at the end of which there is an OH group.




- a) How many **different isomers** can be obtained by **substituting two chlorine atoms**, if only one chlorine atom may be placed at each carbon atom?
- b) How many **different isomers** do you get if you use **one chlorine atom** and **one bromine atom** instead of two chlorine atoms?

$$\frac{9!}{7! \cdot 2!} = 36 \text{ different isomers}$$

$$\frac{9!}{7! \cdot 1! \cdot 1!} = 72 \text{ different isomers}$$

→**NOTE:** The images do not exactly depict the molecule of the task ☺



# Bayes' Theorem

## Remember the probability rules?

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{P(B|A) \cdot P(A)}{P(B)}$$

## Multiplication rule:

- $P(A \cap B) = P(A|B)P(B) = P(B|A)P(A)$

## Law of total probability:

- events  $B_i$  to  $B_n$  partition the entire sample space  $S$

$$P(A) = \sum_{i=1}^N P(A \cap B_i) = \sum_{i=1}^N P(A|B_i) \cdot P(B_i)$$

## Bayes' Theorem

- events  $B_i$  to  $B_n$  partition the entire sample space  $S$
- $P(A) > 0$
- then for  $i = 1, \dots, n$ :

$$P(B_i|A) = \frac{P(A|B_i)P(B_i)}{P(A)}$$

# Bayes' Theorem

$$\bullet P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{\frac{P(A \cap B)}{P(A)} P(A)}{P(B)} = \frac{P(B|A) \cdot P(A)}{P(B)}$$

- we are interested in the probability of two events A and B with  $P(A)$  and  $P(B) > 0$  occurring
- especially the probability of A occurring, given that B occurred via the probability of B occurring, given A
- joint and conditional probabilities

# Bayes' Theorem

$$P(A|B) = \frac{P(B|A)P(A)}{P(B|A)P(A) + P(B|\sim A)P(\sim A)}$$

- Note that  $P(B)$  and  $P(\sim B) = 1$ 
  - $P(A|B)$  – the conditional probability of event A given B
  - $P(B|A)$  – the conditional probability of event B given A
  - $P(A)$  – a priori probability of A
  - $P(B)$  – a priori probability of B

# Real world applications – Bayes' Theorem

How do we interpret **Bayes' Theorem** under the idea of **H** being a research **hypothesis** of interest and **E** some **evidence** we observed?

$$P(H|E) = \frac{P(E|H)P(H)}{P(E)}$$

- $P(H)$  the '**priori probability**' of our hypothesis
  - $P(H|E)$  the probability of our hypothesis after we have observed evidence in favour, or against it! ← '**posterior probability**'
  - $\frac{P(E|H)}{P(E)}$  is called **the likelihood ratio** and  $P(E|H)$  the **likelihood** of how well our hypothesis explains the evidence
- under Bayes, this statement reads as: **The probability of our hypothesis after observing evidence is equal to the probability of our hypothesis multiplied with the likelihood ratio**



# Hands on – Bayes' Theorem

**Task:** 1% of women at age forty who participate in routine screening have breast cancer. 80% of women with breast cancer will get a positive test result. 9,6% of women without breast cancer will receive a false positive on their test results. If a woman in this age group receives a positive test result, what is the probability that she actually has breast cancer?

**Tips:**

$$P(A|B) = \frac{P(B|A)P(A)}{P(B|A)P(A) + P(B|\sim A)P(\sim A)}$$

- Define events **A** and **B**!
- Find  $P(A)$ ,  $P(\sim A)$ ,  $P(B|A)$ ,  $P(B|\sim A)$ !

# Hands on – Bayes' rule

**Solution:**

$$P(A|B) = \frac{P(B|A)P(A)}{P(B|A)P(A) + P(B|\sim A)P(\sim A)}$$

A = breast cancer

B = positive test

$$P(A) = 0.01, P(\sim A) = 0.99, P(B|A) = 0.80, P(B|\sim A) = 0.096$$

$$P(A|B) = \frac{0.80 \cdot 0.01}{0.80 \cdot 0.01 + 0.096 \cdot 0.99} = \frac{0.008}{0.10304} = 0.0776 = 7.76\%$$

→ On average, 7.76% of women who receive a positive test result actually have breast cancer.

# Real world applications – Bayesian Stats

- **Statistics**

- Bayesian Stats – available here at uni (usually)

- **Bioinformatics**

- e.g., used to classify and sort complex & noisy data like DNA samples

- **Your own e-mail accounts** 😊

- using the ‘naïve Bayes’ classificatory’ email programs are capable to filter incoming mails and sort them into folders – including unwanted spam! This is why before deleting spam that got into your main folders, **you should always classify them as spam!** You will train your mailing-program this way

# Odds and odds ratio

## Odds

- the odds of ONE event are given by

$$Pr(y) = \frac{Pr(y)}{Pr(\sim y)}$$

## Odds ratio

- the odds ratio of TWO events are given by

$$\frac{\frac{Pr(x_1)}{Pr(\sim x_1)}}{\frac{Pr(x_2)}{Pr(\sim x_2)}}$$

→ expresses relationship between two independent events occurring!

→ important for MLE

# Hands on – odds and odds ratio

## Task:

1. If  $P(y) = 0.60$ , what are the odds that  $y$  occurs?
2. If the odds of  $x_1$  are 2:1 and the odds of  $x_2$  are 1:2, what is the odds ratio of  $x_1 : x_2$ ?

# Hands on – odds and odds ratio

## **Solution:**

1. If  $P(y) = 0.60$ , what are the odds that  $y$  occurs?

$$P(y) = \frac{0.60}{0.40} = 1.5 \quad [1.5:1]$$

2. If the odds of  $x_1$  are 2:1 and the odds of  $x_2$  are 1:2, what is the odds ratio of  $x_1:x_2$ ?

$$\frac{\frac{2}{1}}{\frac{1}{2}} = \frac{4}{1} = 4$$

# Relative risk – RR

## Intuition:

- origin lies in **medical field**: Do certain groups of people fall ill more easily due to a given risk factor?
- ‘the ratio of two probabilities’ **ranging from 0 to  $\infty$**

## Interpretation

- $RR < 1$  - risk factor **decreases** risk of falling ill / prob in first ‘group’ of ratio is larger
- $RR = 1$  - risk factor has **no apparent effect/risk factor is equal for two groups**
- $RR > 1$  - risk factor **increases** risk of falling ill / prob in first ‘group’ of ratio is smaller

## Example

- $P(\text{war} | \text{one major power}) / P(\text{war} | \sim \text{major power}) = 1.95$   
→ **the group with at least one major power is more likely to start a war!**
- $P(\text{war} | \text{both autocracies}) / P(\text{war} | \text{mixed}) = 0.67$   
→ **the mixed group is more likely to start a war!**

# Relative risk – RR

## Formula:

$$RR := \frac{P(\# \text{ of cases exposed to risk factor})}{P(\# \text{ of cases NOT exposed to risk factor})}$$

## Percentage change:

$$\widehat{RR} = \frac{\frac{a}{(a+c)}}{\frac{b}{(b+d)}}$$

	With risk factor	Without risk factor
Positive cases	a	b
Negative cases	c	d



# Hands on – more practice

## Task:

- 1) If A, B and C are mutually exclusive and collectively exhaustive events and  $P(A) = 0.25$  and  $P(B) = 0.35$ , what is the joint probability of A or C?  $P(A \cup C)$ ?
- 2) Let  $P(A) = 0.3$  and  $P(A \cup B) = 0.5$ . Find  $P(B)$ , assuming both events are independent
- 3) Solve  $\binom{8}{5}$
- 4) Winning a lottery requires drawing 6 numbers from a possibility of 49 numbers, where the order of numbers being drawn does not matter. What is the probability of winning the lottery?

# Hands on – more practice

## Solution:

1)

$$P(C) = 1 - P(A) - P(B) = 0$$

$$P(A \cup C) = P(A) + P(C) = 0.25 + 0.4 = 0.65$$

2)

$$P(A \cup B) = P(A) + P(B) - P(A)P(B)$$

$$0.5 = 0.3 + P(B) - 0.3 \cdot P(B)$$

$$P(B)(1 - 0.3) = 0.2 \leftarrow \text{factorisation!}$$

$$P(B) = \frac{2}{7}$$

# Hands on – more practice

3) Solve  $\binom{8}{5}$

$$\binom{8}{5} = \frac{8!}{5!(8-5)!} = \frac{6 \cdot 7 \cdot 8}{6} = 56$$

4) Winning a lottery requires drawing 6 numbers from a possibility of 49 numbers, where the order of numbers being drawn does not matter. What is the probability of winning the lottery?

$$P(A) = \frac{1}{\binom{49}{6}}$$

$$\binom{49}{6} = \frac{49!}{6! \cdot 43!} = 13,983,816$$

$$P(A) = \frac{1}{13,983,816} = 0.0000000715$$

# Time for your questions

- Any questions during the week?
  - [joerdis.strack@uni-konstanz.de](mailto:joerdis.strack@uni-konstanz.de)

