

# Inter-Rater Reliability (Krippendorff's $\alpha$ ) — stratified by judge-subject overlap

| Measure              | $\alpha$ (all) | $\alpha$ (cross-model) | $\alpha$ (same-model) | $\Delta$ self–cross |
|----------------------|----------------|------------------------|-----------------------|---------------------|
| Ward Total (0–10)    | 0.913          | 0.909                  |                       | -0.256              |
| Ward: symmetry       | 0.881          | 0.879                  |                       | -0.007              |
| Ward: sensory_organs | 0.551          | 0.538                  |                       | -0.008              |
| Ward: locomotion     | 0.904          | 0.903                  |                       | 0.004               |
| Ward: body_plan      | 0.848          | 0.845                  |                       | 0.003               |
| Ward: skin_covering  | 0.632          | 0.623                  |                       | -0.079              |
| Ward: reproduction   | 0.937          | 0.940                  |                       | -0.036              |
| Ward: metabolism     | 0.782          | 0.777                  |                       | -0.047              |
| Ward: communication  | 0.692          | 0.678                  |                       | -0.043              |
| Ward: habitat        | 0.761          | 0.771                  |                       | -0.029              |
| Ward: cognition      | 0.857          | 0.852                  |                       | -0.013              |