

MATH 3503 - Winter 2025

Joey Bernard

2024-07-05

Table of contents

Preface	1
1 Introduction	3
2 First Order ODEs	5
2.1 Introduction	5
2.2 Seperable ODEs	5
2.3 Linear Equations and Integrating Factors	6
2.4 Exact/Homogeneous	6
2.5 Bernoulli	8
3 Second Order ODEs	11
3.1 Homogeneous with Constant Coefficients	11
3.1.1 Case 1	12
3.1.2 Case 2	12
3.1.3 Case 3	13
3.2 Initial Value Problem	13
3.3 Non-homogeneous Differential Equations with Constant Coefficients	13
3.3.1 Undetermined Coefficients	14
3.3.2 Variation of Parameters	14
3.4 Applications	14
4 Systems of ODEs	15
4.1 Fundamentals and Examples	15
4.2 Physical Examples	15
4.3 Ideas and Notation	15

Table of contents

4.4	Eigenvalue Method Examples	15
4.5	Complex Eigenvalues	15
4.6	Vector Fields	15
4.7	Second Order Systems	15
4.8	Repeated Eigenvalues	15
4.9	Matrix Exponentials	15
4.10	Non-homogeneous Equations	15
5	Fourier Series	17
5.1	Boundary Value Problem	17
5.2	Orthogonality of Eigenfunctions	17
5.3	Eigenfunction Decomposition and Trigonometric Series . . .	17
5.4	Sawtooth and Square Wave Examples	17
5.5	More Series	17
6	Laplace Transforms	19
6.1	Introduction	19
6.2	Examples	19
6.3	Inverse Laplace Transform	19
6.4	Solving Homogeneous ODEs	19
6.5	Solving Non-homogeneous ODEs	19
6.6	Unit Step Function	19
6.7	More Transforms of ODEs	19
6.8	Convolution	19
6.9	Impulse Functions	19
7	Summary	21
	References	23

Preface

These notes are for the Winter 2025 term of the course MATH 3503 at the University of New Brunswick. These notes will cover more advanced calculus topics for engineers.

To learn more about Quarto books visit <https://quarto.org/docs/books>.

1 Introduction

This is a book created from markdown and executable code.

See Knuth (1984) for additional discussion of literate programming.

2 First Order ODEs

2.1 Introduction

First order Ordinary Differential Equations (ODEs) are the types of equations that you were exposed to in your first calculus class. These are derivative functions, made up of first derivatives of one independent variable. In the most general case, these look like

$$\frac{dy}{dx} = f(x, y)$$

In the special case where $f(x, y)$ is actually only a function of x (i.e., $f(x, y) = f(x)$), then the most general form of a solution looks like

$$y(x) = \int f(x) dx$$

In this section, we will look at various techniques that can be used to solve other forms of first order ODEs. In all of the following cases, we may or may not have an initial condition. When we do, this initial condition can be used to determine any constants of integration that pop up during calculating the solution.

2.2 Seperable ODEs

One of the simplest forms of first order ODE is the seperable form. This happens when you can write the differential equation in the form of

$$y' = f(x)g(y)$$

2 First Order ODEs

When you have an equation that looks like this, you can rearrange it to look like

$$\frac{dy}{dx} = f(x)g(y)$$
$$\frac{1}{g(y)}dy = f(x)dx$$

Solving the equation becomes as simple as integrating both sides of the rearranged equation.

$$\int \frac{1}{g(y)}dy = \int f(x)dx$$

2.3 Linear Equations and Integrating Factors

The next step in complexity for first order ODEs is if we have an extra function of x and y . The general form in this case looks like

$$y' + p(x)y = f(x)$$

This type of equation is considered linear because there are no higher powers of y or y' . In order to solve these types of equations, we need to use an integrating factor, defined by

$$I(x) = e^{\int p(x)dx}$$

This extra factor can now get used in solving the differential equation by plugging it into

$$y = \frac{1}{I(x)} \int I(x) * f(x)dx$$

2.4 Exact/Homogeneous

Exact equations are special first order ODEs that are derived from a potential function. These are common types of equations that occur in physics and engineering. In these types of cases, you have a potential field that

2.4 Exact/Homogeneous

defines the behaviour described by the differential equation. In the general case, we may have the potential defined by

$$F(x, y)$$

In this general case, we can look at the total derivative of $F(x, y)$.

$$dF = \frac{\partial F}{\partial x} dx + \frac{\partial F}{\partial y} dy$$

This should look familiar from the MATH 2513 class, when you covered partial differential equations. We can use shorter nomenclature and rewrite this as

$$dF = M(x, y)dx + N(x, y)dy$$

If we assume that the potential function is a constant function, then we know that $dF = 0$, and we can rewrite our equation as

$$M(x, y)dx + N(x, y)dy = 0$$

This now looks like something that we may run into when working with first order ODEs. Specifically, this is a homogeneous ODE. If we can write our problem equation in this form, we can work backwards to discover what the solution is. The first step is to integrate either M or N . You usually pick whichever one is simpler to work with. In the M case, we end up with

$$F = \int M dx + C(y)$$

The important part is the extra term of $C(y)$. You then take this solution for $F(x, y)$ and take the derivative with respect to y . You then set this equal to N , in order to solve for what $C(y)$ is. You can also start by integrating N first, and you end up with an extra term of $C(x)$.

$$F = \int N dy + C(x)$$

We can now take the derivative with respect to x and set it equal M . In this way, we can solve for $C(x)$.

In some cases, an equation of the form

2 First Order ODEs

$$Mdx + Ndy = 0$$

actually isn't exact. If we aren't sure, we can test for exactness by checking to see if we have

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial x}$$

This comes from the fact that both M and N are partial derivatives of the potential function F. If you remember from MATH 2513, we have

$$\begin{aligned}\frac{\partial M}{\partial y} &= \frac{\partial N}{\partial x} \\ \frac{\partial^2 F}{\partial y \partial x} &= \frac{\partial^2 F}{\partial x \partial y}\end{aligned}$$

So, if we fail this test, what can we do? In these cases, we can multiply by an integrating factor to get it into an exact form. There is a derivation of how to get the following equation, but we can use

$$P(x) = \frac{M_y - N_x}{N}$$

to find a function. This becomes our integrating factor, that we can substitute into

$$e^{\int P(x)dx}(Mdx + Ndy) = 0$$

This new equation is now exact, and we can use the above technique to solve for a solution.

2.5 Bernoulli

There are several classes of equations where we can use a change of variable to help make the differential equation simpler to solve. One of these types are Bernoulli equations. These have a general form of

$$y' + p(x)y = q(x)y^n$$

In this case, we can start by dividing out both sides of the equation by y^n . This gives us

2.5 Bernoulli

$$y^{-n}y' + p(x)y^{1-n} = q(x)$$

We can now do a change of variables to

$$u = y^{1-n}$$

with the first derivative being

$$u' = (1-n)y^{-n}y'$$

Substituting this back into the original differential equation, we end up with

$$\frac{u'}{(1-n)y^{-n}} + p(x)u = q(x)$$

The resulting new differential equation will then need to be solved using one of the other techniques you learned. In some cases, it may be separable, in some cases you may need to use an integrating factor. Once you have a solution for u , you can then substitute it back into the change of variable you did above.

3 Second Order ODEs

Second order ODEs are differential equations where we may have second derivatives of y with respect to x . The general form looks like

$$A(x)y'' + B(x)y' + C(x)y = F(x)$$

Usually, we divide through so that the y'' term is all by itself. If we do this, we end up with

$$y'' + p(x)y' + q(x)y = f(x)$$

3.1 Homogeneous with Constant Coefficients

The first class of second order ODEs we will look at are homogeneous equations. This is the special case of setting $f(x) = 0$. We will also further constrain the equations for this section by only looking at constant coefficients. This special case looks like

$$y'' + by' + cy = 0$$

In the most general case, there should be two linearly independent solutions (y_1 and y_2) that combine to make the general solution. This general solution looks like

$$y = C_1y_1 + C_2y_2$$

You can check for linear independence by using the Wronskian. If we have linear independence, then we have

3 Second Order ODEs

$$\begin{vmatrix} y_1' & y_2' \\ y_1 & y_2 \end{vmatrix} \neq 0$$

In order to find these two solutions to the differential equation, we start by setting up the characteristic equation that is associated with the differential equation. This characteristic equation is given by

$$ar^2 + br + c = 0$$

We can then find out what the two values for r are. Because this is just a quadratic equation, it is simply

$$r_1, r_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

There are three cases when solving for r .

- Case 1 - r_1 and r_2 are real and unique
- Case 2 - r_1 and r_2 are equal
- Case 3 - r_1 and r_2 are complex

3.1.1 Case 1

If we have case 1, we can easily construct our solution. In this case, it would look like

$$y = C_1 e^{r_1 x} + C_2 e^{r_2 x}$$

3.1.2 Case 2

If we have case 2, we can't simply use the same method as for case 1. We would end up with two equal equations. In this case, we can separate y_1 and y_2 by multiplying one of them by x . This would give us the solution

$$y = C_1 e^{rx} + C_2 x e^{rx}$$

3.1.3 Case 3

Case 3 is a new one if you have never considered the imaginary case for the quadratic equation. In this case, the solution to the characteristic equation is

$$r_{1,2} = \alpha \pm \beta i$$

In this case, we use the real and imaginary parts of r to make up our solution for y . The general form becomes

$$y = C_1 e^{\alpha x} \cos(\beta x) + C_2 e^{\alpha x} \sin(\beta x)$$

3.2 Initial Value Problem

In all of the above cases, we would have constants of integration that would show up in the solutions. In the first order ODE case, a single initial value is enough to define the constant of integration that appears. In the second order case, we will need two initial values. These take the form of

$$\begin{aligned} y(x_0) &= y_0 \\ y'(x_0) &= y'_0 \end{aligned}$$

These can be plugged into the general solution for y to find a particular solution.

3.3 Non-homogeneous Differential Equations with Constant Coefficients

Up to now, we have been working with homogeneous equations where $f(x)=0$. In many physical situations, we end up with a non-homogeneous differential equation. The $f(x)$ term is often called a driving function or forcing function.

3 Second Order ODEs

In order to solve these types of equations, we start by assuming that the equation is homogeneous. In this way, we can find a complimentary solution, y_c that solves the equation when we pretend that $f(x)=0$. The second step is to come up with a second solution, y_p , called the particular solution. We will look at two options of finding this particular solution.

3.3.1 Undetermined Coefficients

The first method is to use the method of undetermined coefficients.

3.3.2 Variation of Parameters

3.4 Applications

4 Systems of ODEs

4.1 Fundamentals and Examples

4.2 Physical Examples

4.3 Ideas and Notation

4.4 Eigenvalue Method Examples

4.5 Complex Eigenvalues

4.6 Vector Fields

4.7 Second Order Systems

4.8 Repeated Eigenvalues

4.9 Matrix Exponentials

4.10 Non-homogeneous Equations

5 Fourier Series

5.1 Boundary Value Problem

5.2 Orthogonality of Eigenfunctions

5.3 Eigenfunction Decomposition and Trigonometric Series

5.4 Sawtooth and Square Wave Examples

5.5 More Series

6 Laplace Transforms

6.1 Introduction

6.2 Examples

6.3 Inverse Laplace Transform

6.4 Solving Homogeneous ODEs

6.5 Solving Non-homogeneous ODEs

6.6 Unit Step Function

6.7 More Transforms of ODEs

6.8 Convolution

6.9 Impulse Functions

7 Summary

In summary, this book has no content whatsoever.

References

Knuth, Donald E. 1984. “Literate Programming.” *Comput. J.* 27 (2): 97–111. <https://doi.org/10.1093/comjnl/27.2.97>.

