



Winter survey 2020 R/V Johan Hjort

Report generated by: Johanna Fall

08/03/2020


Contents

Cruise tracks and stations	3
Sampling depth for trawl hauls	4
Mean bottom depth during trawl hauls	4
Mean fishing depth during trawl hauls	5
Catch rates - biomass and number per nautical mile trawled for species registered in Sea2Data	6
Species diversity	6
Number of species identified versus the number of stations sampled	6
Average catch rate by species for the 20 species with highest catch rates	7
Bottom trawl	7
Bottom trawl excluding sponges	8
Spatial variation in catches of common species	9
Capelin	9
Polar cod	10
Blue whiting	12
Herring	13
Haddock	14
Cod	16

Redfish	17
Length distributions	18
Length-weight relationships	20
Length-age relationships	22
GSI versus maturity stage of female cod and haddock	24
GSI versus length and maturity stage for female haddock	25
GSI versus length and maturity stage for female cod	26
Acoustic registrations	27
Example echograms	27
Depth-integrated acoustic backscatter	27
Total backscatter	27
Pelagic species	28
Demersal species	28
Acoustic backscatter in depth channels	29
Distance from the surface to weighted depth of acoustic registrations	29
Distance from the seafloor (or max scrutinized depth) to weighted depth of acoustic registrations	31
CTD	33
Summary of measurements	33
Variation in temperature and salinity with bathymetry and geographical location	34
Density in the water column	35
Light in the water column	36

Cruise tracks and stations

Cruise tracks from the position log with points indicating start positions for different sampling gear. The points are jittered slightly for better visual representation:




Sampling depth for trawl hauls


153 trawl hauls were taken during the survey, of which 148 were bottom trawl, and 5 pelagic trawl. The trawl hauls covered a total distance of 264 km (142.5 nmi).

The sampling stations were located in areas with bottom depths from 48.7 m to 479.3 m, and the fishing depth varied from 25.7 m to 480.7 m.

Mean bottom depth during trawl hauls



Mean fishing depth during trawl hauls




Catch rates - biomass and number per nautical mile trawled for species registered in Sea2Data

Species diversity



A total of 75 species were registered in Sea2Data during the survey.

Number of species identified versus the number of stations sampled





Average catch rate by species for the 20 species with highest catch rates

Bottom trawl




Bottom trawl excluding sponges




Spatial variation in catches of common species
Capelin

Biomass | 2020 | *Mallotus villosus*




Number of individuals | 2020 | *Mallotus villosus*




Polar cod

Biomass | 2020 | *Boreogadus saida*




Number of individuals | 2020 | *Boreogadus saida*




Blue whiting

Biomass | 2020 | *Micromesistius poutassou*




Number of individuals | 2020 | *Micromesistius poutassou*




Herring

Biomass | 2020 | *Clupea harengus*




Number of individuals | 2020 | *Clupea harengus*




Haddock

Biomass | 2020 | *Melanogrammus aeglefinus*




Number of individuals | 2020 | *Melanogrammus aeglefinus*




Cod

Biomass | 2020 | *Gadus morhua*




Number of individuals | 2020 | *Gadus morhua*




Redfish

Biomass | 2020 | Redfish species




Number of individuals | 2020 | Redfish species




Length distributions


The following figures shows the length distribution of individuals sampled during the survey.




Histogram of length





Length-weight relationships






Length-age relationships






GSI versus maturity stage of female cod and haddock




GSI versus length and maturity stage for female haddock

The vertical line represents the size (40 cm) over which gonad samples should be taken if GSI < 2 % (horizontal line).



GSI versus length and maturity stage for female cod

The vertical line represents the size (60 cm) over which most females with GSI > 1.5 % (horizontal line) are maturing (Skjæraasen et al. 2012).




Acoustic registrations

A total of 5781.944 km (3122 nmi) of acoustic transects were scrutinized.

Example echograms


Depth-integrated acoustic backscatter

Total backscatter




Pelagic species

Winter survey Johan Hjort | 2020 | 38 kHz




Demersal species

Winter survey Johan Hjort | 2020 | 38 kHz




Acoustic backscatter in depth channels


Distance from the surface to weighted depth of acoustic registrations




Demersal species



Distance from the seafloor (or max scrutinized depth) to weighted depth of acoustic registrations





Demersal species




CTD

Summary of measurements



52 CTD casts were done during the survey, covering a total depth of 7.5 km.



Variation in temperature and salinity with bathymetry and geographical location



Density in the water column



Light in the water column

