

# Question 1

Joseph Froelicher

February 26, 2020

## Part A

$$\ln \left( \frac{p_i}{1 - p_i} \right) = \mathbf{X}_j \beta$$

$$\hat{\beta}_0 = \ln \left( \frac{p(y=1|x_j=0)}{1 - p(y=1|x_j=0)} \right) = \ln \left( \frac{(619/2416)}{1 - (619/2416)} \right) = -1.065769$$

$$\hat{\beta}_1 = (\hat{\beta}_0 + \hat{\beta}_1) - \hat{\beta}_0 = \ln \left( \frac{p(y=1|x_j=1)}{1 - p(y=1|x_j=1)} \right) - \hat{\beta}_0 = \ln \left( \frac{(355/771)}{1 - (355/771)} \right) - (-1.065769) = 0.9072015$$

$$\hat{\beta}_2 = (\hat{\beta}_0 + \hat{\beta}_2) - \hat{\beta}_0 = \ln \left( \frac{p(y=1|x_j=2)}{1 - p(y=1|x_j=2)} \right) - \hat{\beta}_0 = \ln \left( \frac{(162/731)}{1 - (162/731)} \right) - (-1.065769) = -0.1905151$$

## Part B

$$L(Y_i|p) = \prod_{i=1}^N \binom{n_i}{Y_i} p^{Y_i} (1-p)^{n_i-Y_i}$$

$$L(Y_i|p) = \prod_{i=1}^N \binom{n_i}{Y_i} \left( \frac{e^{\mathbf{X}\beta}}{1 + e^{\mathbf{X}\beta}} \right)^{Y_i} \left( 1 - \frac{e^{\mathbf{X}\beta}}{1 + e^{\mathbf{X}\beta}} \right)^{n_i-Y_i}$$

$$L(Y_i|p) = \left( \frac{e^{\mathbf{X}\beta}}{1 + e^{\mathbf{X}\beta}} \right)^{\sum_{i=1}^n Y_i} \left( 1 - \frac{e^{\mathbf{X}\beta}}{1 + e^{\mathbf{X}\beta}} \right)^{n - \sum_{i=1}^n Y_i}$$

$$\ln(L(Y_i|p)) = \ln \left( \left( \frac{e^{\mathbf{X}\beta}}{1 + e^{\mathbf{X}\beta}} \right)^{\sum_{i=1}^n Y_i} \left( 1 - \frac{e^{\mathbf{X}\beta}}{1 + e^{\mathbf{X}\beta}} \right)^{n - \sum_{i=1}^n Y_i} \right)$$

$$l(Y_i|p) = \sum_{i=1}^n \left( \ln \binom{n}{Y_i} + Y_i \ln \left( \frac{e^{\mathbf{X}\beta}}{1 + e^{\mathbf{X}\beta}} \right) + (n - Y_i) \ln \left( 1 - \frac{e^{\mathbf{X}\beta}}{1 + e^{\mathbf{X}\beta}} \right) \right)$$

$$l(Y_i|p) = \left[ \ln \binom{n_1}{Y_1} + Y_1 \ln \left( \frac{e^{\hat{\beta}_0}}{1 + e^{\hat{\beta}_0}} \right) + (n_1 - Y_1) \ln \left( 1 - \frac{e^{\hat{\beta}_0}}{1 + e^{\hat{\beta}_0}} \right) \right]$$

$$+ \left[ \ln \binom{n_2}{Y_2} + Y_2 \ln \left( \frac{e^{\hat{\beta}_0 + X_1 \hat{\beta}_1}}{1 + e^{\hat{\beta}_0 + X_1 \hat{\beta}_1}} \right) + (n_2 - Y_2) \ln \left( 1 - \frac{e^{\hat{\beta}_0 + X_1 \hat{\beta}_1}}{1 + e^{\hat{\beta}_0 + X_1 \hat{\beta}_1}} \right) \right]$$

$$+ \left[ \ln \binom{n_3}{Y_3} + Y_3 \ln \left( \frac{e^{\hat{\beta}_0 + X_2 \hat{\beta}_2}}{1 + e^{\hat{\beta}_0 + X_2 \hat{\beta}_2}} \right) + (n_3 - Y_3) \ln \left( 1 - \frac{e^{\hat{\beta}_0 + X_2 \hat{\beta}_2}}{1 + e^{\hat{\beta}_0 + X_2 \hat{\beta}_2}} \right) \right]$$

By hand, the log likelihood of model 1 is -10.8699802. Using the `logLik()` function in R, we get the value of -10.8699802 for the likelihood of model 1.

## Part C

$$\begin{aligned}
 \ln(L(Y_i|p)) &= \ln \left( \left( \frac{e^{\mathbf{X}\beta}}{1 + e^{\mathbf{X}\beta}} \right)^{\sum_{i=1}^n Y_i} \left( 1 - \frac{e^{\mathbf{X}\beta}}{1 + e^{\mathbf{X}\beta}} \right)^{n - \sum_{i=1}^n Y_i} \right) \\
 l(Y_i|p) &= \sum_{i=1}^n \left( \ln \binom{n}{Y_i} + Y_i \ln \left( \frac{e^{\mathbf{X}\beta}}{1 + e^{\mathbf{X}\beta}} \right) + (n - Y_i) \ln \left( 1 - \frac{e^{\mathbf{X}\beta}}{1 + e^{\mathbf{X}\beta}} \right) \right) \\
 l(Y_i|p) &= \left[ \ln \binom{n_1}{Y_1} + Y_1 \ln \left( \frac{e^{\hat{\beta}_0}}{1 + e^{\hat{\beta}_0}} \right) + (n_1 - Y_1) \ln \left( 1 - \frac{e^{\hat{\beta}_0}}{1 + e^{\hat{\beta}_0}} \right) \right]
 \end{aligned}$$

By hand, the log likelihood of model 0 is -4.2654493. Using the `logLik()` function in R, we get the value of -4.2654493 for the likelihood of model 1.

## Part D

$H_0$  : The null, intercept only model, sufficiently models the data.  $H_a$  : The full, covariate model, is significantly better at modeling the data.

$$T_{LR} = -2\ln \left[ \frac{L(p_{H_0})}{L(p_{MLE})} \right]$$

$$T_{LR} = -2\ln \left[ \frac{e^{-4.26545}}{e^{-10.86998}} \right]$$

$$X_{2,1-0.95}^2 = 5.991$$

## [1] 1

$$P\{(T_{LR} = -13.209) > 5.991\} > 0.05$$

$$(T_{LR} = 131.0817) \not\sim X_{2,0.95}^2$$

Fail to reject the null hypothesis. The likelihood ratio test suggested that our full model, does not fit the data significantly better than the null model.

## Part E

The predicted probability of a misconduct violation during the first year in prison for a prisoner with 1 strike is 0.3938046. The predicted probability of a misconduct violation during the first year in prison for a prisoner with 3 strikes is 0.4464219.

## Part F

$$CI = [e^{\hat{\beta} - Z_{1-\frac{\alpha}{2}} \hat{SE}(\hat{\beta})}, e^{\hat{\beta} + Z_{1-\frac{\alpha}{2}} \hat{SE}(\hat{\beta})}]$$

$$CI = [0.9525757, 1.3490558]$$

## Part G

Two reasons why model 1 is better than model 2: 1. We shouldn't try to model discrete observations as continuous variables, as we are doing in model 2. This could result in unexpected model behaviour. 2. The AIC for model 1 is significantly lower ( $> 10$ ) than model 2. There is no indication that model 2 is a better fit for our data than model 1.

## Appendix

```
# part b
data = data.frame(
  "strikes" = c("1", "2", "3"),
  "misconduct" = c(619, 355, 162),
  "no_misconduct" = c(1797, 416, 569)
)

# by hand
p1 = data[1, 2] / (data[1, 2] + data[1, 3])
p2 = data[2, 2] / (data[2, 2] + data[2, 3])
p3 = data[3, 2] / (data[3, 2] + data[3, 3])

n1 = data[1, 2] + data[1, 3]
n2 = data[2, 2] + data[2, 3]
n3 = data[3, 2] + data[3, 3]

c1 = data[1, 2]
c2 = data[2, 2]
c3 = data[3, 2]

w1 = data[1, 3]
w2 = data[2, 3]
w3 = data[3, 3]

ll_m1 = lchoose(n1, c1) + c1 * log(p1) + w1 * log(1 - p1)
ll_m2 = lchoose(n2, c2) + c2 * log(p2) + w2 * log(1 - p2)
ll_m3 = lchoose(n3, c3) + c3 * log(p3) + w3 * log(1 - p3)
like1_hand = sum(ll_m1, ll_m2, ll_m3)

model1 = glm(
  cbind(misconduct, no_misconduct) ~ strikes, data, family = binomial
)

like1_r = logLik(model1)

# part b
n0 = n1 + n2 + n3
c0 = c1 + c2 + c3
w0 = w1 + w2 + w3

model0 = glm(
  cbind(c0, w0) ~ 1, data, family = binomial
)

# by hand
p0 = exp(summary(model0)$coefficients[1, 1]) / (1 + exp(summary(model0)$coefficients[1, 1]))

like0_hand = lchoose(n0, c0) + c0 * log(p0) + w0 * log(1 - p0)

# check
like0_r = logLik(model0)
```

```

# part b
t_lr = -2 * (like0_hand - like1_hand)
pchisq(t_lr, 2, lower.tail = F)

# numeric model 2
data_numeric = data.frame(
  "strikes" = c(1, 2, 3),
  "misconduct" = c(619, 355, 162),
  "no_misconduct" = c(1797, 416, 569)
)

model2 = glm(
  cbind(misconduct, no_misconduct) ~ strikes, data_numeric, family = binomial
)

summary(model2)

# part E
p_1 = summary(model2)$coefficients[1, 1] + summary(model2)$coefficients[2, 1]
p_3 = summary(model2)$coefficients[1, 1] + (3 * summary(model2)$coefficients[2, 1])

# part f
f = exp(p_3) / exp(p_1)
a = 0.05
z = qnorm(1 - (a / 2))
se = 2 * summary(model2)$coefficients[2, 2]
b = p_3 - p_1
ci = c(exp(b - (z * se)), exp(b + (z * se)))

```