

Project 2: Optimal Management

Josephine Walker

May 16, 2013

Collaborators

- Michael Bode (UMelb)
- Hugh Possingham (UQ)

1 Introduction

Communities at the border of Makgadikgadi Pans National Park in Botswana make up part of a social-ecological system which includes people, domestic animals, wild animals, parasites, plants, and other organisms. People interact with the rest of the system by raising livestock, planting crops, and applying management strategies such as building fences, using pesticides, or digging wells. At this point, decisions on the best methods for management are generally not based on economic outcomes. The goal of this project is to develop a way to conduct cost-benefit analyses for various management strategies that incorporates impacts for multiple players in the system - including conservation and tourism as well as livelihoods of the local people. The method I am currently using is called 'qualitative modeling'. The first step is to develop a network model of the ecosystem based on Lotka-Volterra interactions, incorporating uncertain links. Next, I subjected simulations of this network to combinations of press perturbations which represent management strategies. The results of the press perturbations indicate whether certain nodes in the network are likely to increase or decrease following the perturbation. These results will then be incorporated into a cost-benefit equation based on estimated utility functions that describe the value of each node. In this way, management strategies can be compared based on their predicted cost-benefit ratio, as well as the uncertainty of the outcome for each node.

2 Methods

2.1 Defining a network

The network model is defined by linked Lotka-Volterra-type equations. This can be written as a system of differential equations as defined by Hosack, Hayes and Dambacher:

$$\frac{dX_i}{dt} = f_i(X_1, \dots, X_n; p_1, \dots, p_k)$$

$$i = 1 \dots n, k = 1 \dots m$$

where m parameters p_k control the rates of increase or decrease within each growth function f_i [1].

For example, for $n = 2$ the equations could look like this:

$$\frac{dX_1}{dt} = f_1(X_1, X_2; p_1, p_2) = p_1 X_1 - p_2 X_1 X_2$$

$$\frac{dX_2}{dt} = f_2(X_1, X_2; p_3, p_4) = p_3 X_2 - p_4 X_2 X_1$$

The system is linearized by evaluating the first partial derivatives of the growth functions f_i with respect to each X_j at equilibrium, this makes up the $n \times n$ Jacobian matrix \mathbf{A} .

Each element of \mathbf{A} , a_{ij} , therefore represents the effect of changes in X_j (the node in the column) on X_i (the node in the row), and can be written, as in [1], as

$$a_{ij} = \left. \frac{\partial f_i}{\partial X_j} \right|_*$$

where $*$ denotes evaluation at equilibrium.

The values of each individual parameter p_k in the Lotka-Volterra equations or each a_{ij} in \mathbf{A} are not known, but the results of the model are likely to be highly sensitive to the values chosen [1]. Instead of defining values, I defined the biological class of interaction for each cell in the matrix (for example: predator, prey, competition) in order to decide whether a_{ij} will be positive or negative (Table 1). This is one sense in which the model is 'qualitative'.

I defined the model to have 17 nodes representing relevant groups of organisms in the system, including separating parasites into infective and within-host stages. Each possible interaction is defined as one of nine possible classes representing combinations of positive, negative, and no interactions between two nodes:

- **ab**, an abiotic or unidirectional increase, such as the effect water has on everything, a_{ji} is 0
- **uk**, unknown, could be positive or negative or no link
- **mu**, mutualistic, positive impact and a_{ji} is also positive
- **pr**, predator, negative impact and a_{ji} is positive (prey)
- **py**, prey, positive impact and a_{ji} is negative (predator)
- **co**, competition, negative impact and a_{ji} is also negative
- **dd**, density-dependent, negative impact through competition, but only on the diagonal indicating the effect of a node on itself
- **na**, no interaction
- **in**, unidirectional decrease, where a_{ji} is 0

This is based on the approach taken by [2] in their assessment of the impact of eradication of invasive species on Macquarie Island. They define their network in a similar manner and go on to simulate values for each a_{ij} . First, in order to deal with unknown links, they look at each possible combination of x unknown links existing or not (2^x), and simulate 1000 Lyapunov stable (stable around equilibrium, all real parts of eigenvalues are negative) values for \mathbf{A} for each combination, resulting in 131 million matrices.

Instead of taking this approach, which leads to equal weighting of each combination regardless of the likelihood of that combination yielding a stable matrix, I simulate values so that each unknown has an equal probability of being negative, zero, or positive. Various changes to how each a_{ij} is chosen makes it more or less likely that a simulated \mathbf{A} will be stable, but in general many iterations are necessary to reach a large number of stable simulations (25+ unstable matrices generated per stable matrix).

2.2 Perturbations on a network

Simulation of the matrices

2.3 Incorporating management strategies

2.4 Cost-benefit analysis and utility functions

2.5 Differences between communities

McDermott describes how the economics of rural livestock production depend on the degree of intensification of the system as well as the degree of importance of livestock to an individual household economy [3]. In this case, I think that the level of intensification is likely to be similar across villages as most use a very low-input method of raising livestock (letting them graze wherever they want). Instead, in this context the degree of interaction with wildlife will be important. I have made a figure which represents where hypothetical farmers would fall on these spectra (Figure 1).




Figure 1: Hypothetical spectra representing differences between farmers or communities

In practice, I think I can incorporate this issue into the model by comparing a few subsets of simulated models where some defined interaction factor is high/med/low. I have not thought much about this yet, but perhaps I can subset the simulated networks by using cutoffs for strength of links between domestic and wild animals. For example, simulated networks with highly competitive interactions between domestic and wild grazers, and/or with high levels of predation of domestic animals by wild carnivores, would fall into the high interaction category. This could be subsetted from all simulated networks or new networks could be simulated with these additional constraints.

The goal of dividing communities by level of interaction would be to compare how these differences affect the recommended management strategy from the perspective of a particular community. In addition, it will be useful to define whether individual communities may be affected differently by the same management strategy.

2.6 Limitations and some other possible methods

The qualitative modeling method defines a framework for assessment of the economic impact of a management strategy based on the modeled structure of the system. By simulating many possible networks, we incorporate the high levels of uncertainty about the model structure. Essentially, everything is highly uncertain and yet we try to make conclusions anyway. By using a complex model, there are so many parameters to be defined and this contributes to the uncertainty. I have thought a little bit about whether it would be possible to use a simpler model.

From what I have seen, there is only one study which attempts to economically examine the impact of wildlife contact on communities [4]. They do an analysis which incorporates potential costs and benefits to communities of living near the park, but they don't consider the impact of humans on the wildlife or of the impact of different management strategies. I could use a similar method with additional factors included and estimate parameters based on expert opinion instead of using simulated networks.

Another study explicitly models wildebeest population growth in relation to poaching pressure by communities in order to determine the risk of collapse of the wildebeest population under different management strategies [5]. I could potentially do something similar to this, where I focus on transmission of parasites between domestic and wild populations using fairly simple population growth models

(with competition just between two species) which would be possible to parameterize explicitly. This would allow me to estimate costs and benefits of parasite management but would neglect many interactions and effects.

References

- [1] GR Hosack, KR Hayes, and JM Dambacher. Assessing model structure uncertainty through an analysis of system feedback and Bayesian networks. *Ecological Applications*, 18(4):1070–1082, 2008.
- [2] Ben Raymond, Julie Mcinnes, Jeffrey M Dambacher, Sarah Way, and Dana M. Bergstrom. Qualitative modelling of invasive species eradication on subantarctic Macquarie Island. *Journal of Applied Ecology*, 48(1):181–191, February 2011.
- [3] JJ McDermott, TF Randolph, and SJ Staal. The economics of optimal health and productivity in smallholder livestock systems in developing countries. *Revue scientifique et ...*, 18(2):399–424, 1999.
- [4] Petronella Chaminuka, Cheryl M. E. McCrindle, and Henk M. J. Udo. Cattle Farming at the Wildlife/Livestock Interface: Assessment of Costs and Benefits Adjacent to Kruger National Park, South Africa. *Society & Natural Resources*, 25(3):235–250, March 2012.
- [5] CB Barrett and Peter Arcese. Wildlife harvest in integrated conservation and development projects: linking harvest to household demand, agricultural production, and environmental shocks in the. *Land economics*, 74(4):449–465, 1998.

	Water	Grass	Trees	Crops	Domestic Ruminants	Domestic Equids	Wild Ruminants	Wild Equids	Parasites Dom Eq	Parasites Wild Eq	Parasites Dom Rum	Parasites Wild Rum	Infectious Eq Par	Infectious Rum Par	Carnivores	Elephants	People
Water	ab	uk	uk	uk	uk	uk	uk	uk	uk	uk	uk	uk	uk	uk	uk	uk	uk
Grass	ab	dd	co	uk	pr	pr	pr	pr	na	na	na	na	na	na	na	uk	uk
Trees	ab	co	dd	uk	na	na	na	na	na	na	na	na	na	na	na	pr	uk
Crops	ab	uk	uk	dd	uk	uk	co	na	na	na	na	na	na	na	na	pr	mu
Domestic Ruminants	ab	py	uk	uk	dd	co	co	co	pr	na	pr	na	na	na	pr	na	mu
Domestic Equids	ab	py	uk	uk	co	co	co	co	na	na	na	na	na	na	pr	na	mu
Wild Ruminants	ab	py	uk	na	co	co	co	dd	na	pr	na	pr	na	na	pr	uk	na
Wild Equids	ab	py	uk	na	co	co	co	dd	na	pr	na	na	na	na	pr	uk	na
Parasites Dom Eq	ab	uk	na	na	na	py	na	na	dd	na	na	na	mu	na	na	na	na
Parasites Wild Eq	ab	uk	na	na	na	na	na	py	na	dd	na	na	mu	na	na	na	na
Parasites Dom Rum	ab	uk	na	na	py	na	na	na	na	na	dd	na	na	mu	na	na	na
Parasites Wild Rum	ab	uk	na	na	na	na	py	na	na	na	na	dd	na	mu	na	na	na
Infectious Eq Par	ab	ab	na	na	in	in	in	in	mu	mu	na	na	dd	na	na	na	na
Infectious Rum Par	ab	ab	na	na	in	in	in	in	na	na	mu	na	na	dd	na	uk	in
Carnivores	ab	na	uk	na	py	py	py	py	na	na	na	na	na	na	dd	uk	in
Elephants	ab	uk	py	py	na	na	na	na	na	na	na	na	na	na	uk	dd	na
People	ab	uk	uk	mu	mu	mu	na	na	na	na	na	na	na	na	na	na	dd

Table 1: Defined interactions between nodes; ab = abiotic/incidental increase, uk = unknown, mu = mutualistic, pr = predator, py = prey, co = competition, dd = density-dependent (diagonal), na = none, in = incidental reduction