

=====
=====
Group difference of FD and log FC -- Without total CTQ as covariate
=====
=====

Loading FD data...

=====
UF_fd
=====

Descriptive statistics (raw):

	mean	std	count
group			
PTSD	0.403997	0.016449	51
control	0.408739	0.018337	57

Assumption checks

group*mri_age_demeaned interaction not significant (p=0.5376). Homogeneity of slopes assumed.
Levene W=0.281, p=0.597 — homogeneity of variances assumed.
Breusch-Pagan p=0.3642 — homoscedasticity assumed.
Shapiro-Wilk W=0.985, p=0.274 — normality of residuals assumed.

ANCOVA results (HC3-robust, Type II) — UF_fd

	sum_sq	df	F	PR(>F)	np2
C(group)	0.000216	1.0	0.921146	3.393791e-01	0.008697
mri_age_demeaned	0.008623	1.0	36.750629	2.137453e-08	0.259263
Residual	0.024636	105.0	NaN	NaN	0.500000
group effect not significant (p=0.3394)					

OLS coefficient table (HC3 SEs):

	coef	std err	t	P> t	[0.025	0.975]
Intercept	0.4050	0.002	189.999	0.000	0.401	0.409
C(group)[T.control]	0.0029	0.003	0.960	0.339	-0.003	0.009
mri_age_demeaned	-0.0008	0.000	-6.062	0.000	-0.001	-0.001

Adjusted group means (all covariates at their mean):

	group	adjusted_mean	mean	mean_se	mean_ci_lower	mean_ci_upper	obs_ci_lower	obs_ci_upper
0	control	0.407869	0.407869	0.002114	0.403678	0.412061	0.377209	0.438529
1	PTSD	0.404969	0.404969	0.002131	0.400743	0.409195	0.374304	0.435634

=====
CB_fd
=====

Descriptive statistics (raw):

	mean	std	count
group			
PTSD	0.372557	0.012107	51
control	0.375593	0.015093	57

Assumption checks

group*mri_age_demeaned interaction not significant (p=0.7998). Homogeneity of slopes assumed.
Levene W=1.274, p=0.262 — homogeneity of variances assumed.
Breusch-Pagan p=0.2284 — homoscedasticity assumed.
Shapiro-Wilk W=0.989, p=0.552 — normality of residuals assumed.

ANCOVA results (HC3-robust, Type II) — CB_fd

	sum_sq	df	F	PR(>F)	np2
C(group)	0.000199	1.0	1.047678	0.308395	0.009879
mri_age_demeaned	0.000089	1.0	0.469345	0.494798	0.004450
Residual	0.019951	105.0	NaN	NaN	0.500000
group effect not significant (p=0.3084)					

OLS coefficient table (HC3 SEs):

	coef	std err	t	P> t	[0.025	0.975]
Intercept	0.3727	0.002	215.143	0.000	0.369	0.376
C(group)[T.control]	0.0028	0.003	1.024	0.308	-0.003	0.008
mri_age_demeaned	-0.0001	0.000	-0.685	0.495	-0.000	0.000

Adjusted group means (all covariates at their mean):

	group	adjusted_mean	mean	mean_se	mean_ci_lower	mean_ci_upper	obs_ci_lower	obs_ci_upper
0	control	0.375479	0.375479	0.002071	0.371373	0.379585	0.347840	0.403118
1	PTSD	0.372685	0.372685	0.001732	0.369250	0.376120	0.345138	0.400232

=====
ST_fd
=====

Descriptive statistics (raw):

	mean	std	count
group			

PTSD 0.443102 0.037062 51
control 0.453445 0.032223 57

Assumption checks

group*mri_age_demeaned interaction not significant (p=0.9931). Homogeneity of slopes assumed.
Levene W=0.500, p=0.481 — homogeneity of variances assumed.
WARNING: Heteroscedasticity detected (Breusch-Pagan p=0.0152)
Shapiro-Wilk W=0.991, p=0.734 — normality of residuals assumed.

ANCOVA results (HC3-robust, Type II) — ST_fd

	sum_sq	df	F	PR(>F)	np2
C(group)	0.001148	1.0	1.242850	0.267468	0.011698
mri_age_demeaned	0.019456	1.0	21.056436	0.000012	0.167040
Residual	0.097021	105.0	NaN	NaN	0.500000

group effect not significant (p=0.2675)

OLS coefficient table (HC3 SEs):

	coef	std err	t	P> t	[0.025	0.975]
Intercept	0.4450	0.005	95.285	0.000	0.436	0.454
C(group)[T.control]	0.0067	0.006	1.115	0.267	-0.005	0.019
mri_age_demeaned	-0.0016	0.000	-4.589	0.000	-0.002	-0.001

Adjusted group means (all covariates at their mean):

	group	adjusted_mean	mean	mean_se	mean_ci_lower	mean_ci_upper	obs_ci_lower	obs_ci_upper
0	control	0.451737	0.451737	0.003807	0.444188	0.459286	0.390993	0.512480
1	PTSD	0.445012	0.445012	0.004670	0.435751	0.454272	0.384032	0.505992

Loading FC data...

UF_fc

Descriptive statistics (raw):

	mean	std	count
group			
PTSD	0.054032	0.069023	51
control	0.084932	0.077414	57

Assumption checks

group*mri_age_demeaned interaction not significant (p=0.9341). Homogeneity of slopes assumed.
group*t1_volume_demeaned interaction not significant (p=0.6797). Homogeneity of slopes assumed.
Levene W=1.288, p=0.259 — homogeneity of variances assumed.
Breusch-Pagan p=0.2280 — homoscedasticity assumed.
Shapiro-Wilk W=0.990, p=0.593 — normality of residuals assumed.

ANCOVA results (HC3-robust, Type II) — UF_fc

	sum_sq	df	F	PR(>F)	np2
C(group)	0.000015	1.0	0.009072	9.243007e-01	0.000087
mri_age_demeaned	0.000294	1.0	0.180737	6.716191e-01	0.001735
t1_volume_demeaned	0.372159	1.0	228.515307	5.290667e-28	0.687232
Residual	0.169374	104.0	NaN	NaN	0.500000

group effect not significant (p=0.9243)

OLS coefficient table (HC3 SEs):

	coef	std err	t	P> t	[0.025	0.975]
Intercept	0.0699	0.006	12.205	0.000	0.059	0.081
C(group)[T.control]	0.0008	0.008	0.095	0.924	-0.015	0.017
mri_age_demeaned	-0.0002	0.000	-0.425	0.672	-0.001	0.001
t1_volume_demeaned	4.839e-07	3.2e-08	15.117	0.000	4.2e-07	5.47e-07

Adjusted group means (all covariates at their mean):

	group	adjusted_mean	mean	mean_se	mean_ci_lower	mean_ci_upper	obs_ci_lower	obs_ci_upper
0	control	0.070701	0.070701	0.00555	0.059695	0.081706	-0.010080	0.151481
1	PTSD	0.069938	0.069938	0.00573	0.058575	0.081301	-0.010892	0.150768

CB_fc

Descriptive statistics (raw):

	mean	std	count
group			
PTSD	0.017148	0.075854	51
control	0.043794	0.084570	57

Assumption checks

group*mri_age_demeaned interaction not significant (p=0.5494). Homogeneity of slopes assumed.
group*t1_volume_demeaned interaction not significant (p=0.5124). Homogeneity of slopes assumed.

Levene W=0.953, p=0.331 — homogeneity of variances assumed.
Breusch-Pagan p=0.3670 — homoscedasticity assumed.
Shapiro-Wilk W=0.997, p=0.996 — normality of residuals assumed.

ANCOVA results (HC3-robust, Type II) — CB_fc

	sum_sq	df	F	PR(>F)	np2
C(group)	0.000821	1.0	0.529760	4.683432e-01	0.005068
mri_age_demeaned	0.004554	1.0	2.939962	8.939124e-02	0.027492
t1_volume_demeaned	0.392648	1.0	253.506996	1.203092e-29	0.709097
Residual	0.161082	104.0	NaN	NaN	0.500000

group effect not significant (p=0.4683)

OLS coefficient table (HC3 SEs):

	coef	std err	t	P> t	[0.025	0.975]
Intercept	0.0343	0.006	5.538	0.000	0.022	0.047
C(group)[T.control]	-0.0059	0.008	-0.728	0.468	-0.022	0.010
mri_age_demeaned	0.0006	0.000	1.715	0.089	-9.16e-05	0.001
t1_volume_demeaned	5.525e-07	3.47e-08	15.922	0.000	4.84e-07	6.21e-07

Adjusted group means (all covariates at their mean):

	group	adjusted_mean	mean	mean_se	mean_ci_lower	mean_ci_upper	obs_ci_lower	obs_ci_upper
0	control	0.028409	0.028409	0.005027	0.018440	0.038379	-0.050269	0.107087
1	PTSD	0.034343	0.034343	0.006201	0.022045	0.046641	-0.044664	0.113349

ST_fc

Descriptive statistics (raw):

	mean	std	count
group			
PTSD	-0.063659	0.117743	51
control	-0.046215	0.110523	57

Assumption checks

group*mri_age_demeaned interaction not significant (p=0.3987). Homogeneity of slopes assumed.
group*t1_volume_demeaned interaction not significant (p=0.3334). Homogeneity of slopes assumed.
Levene W=0.478, p=0.491 — homogeneity of variances assumed.
Breusch-Pagan p=0.8142 — homoscedasticity assumed.
Shapiro-Wilk W=0.984, p=0.229 — normality of residuals assumed.

ANCOVA results (HC3-robust, Type II) — ST_fc

	sum_sq	df	F	PR(>F)	np2
C(group)	0.000046	1.0	0.006472	9.360350e-01	0.000062
mri_age_demeaned	0.214495	1.0	30.270908	2.712867e-07	0.225447
t1_volume_demeaned	0.300016	1.0	42.340323	2.747524e-09	0.289328
Residual	0.736926	104.0	NaN	NaN	0.500000

group effect not significant (p=0.9360)

OLS coefficient table (HC3 SEs):

	coef	std err	t	P> t	[0.025	0.975]
Intercept	-0.0537	0.013	-4.202	0.000	-0.079	-0.028
C(group)[T.control]	-0.0014	0.018	-0.080	0.936	-0.037	0.034
mri_age_demeaned	0.0047	0.001	5.502	0.000	0.003	0.006
t1_volume_demeaned	4.814e-07	7.4e-08	6.507	0.000	3.35e-07	6.28e-07

Adjusted group means (all covariates at their mean):

	group	adjusted_mean	mean	mean_se	mean_ci_lower	mean_ci_upper	obs_ci_lower	obs_ci_upper
0	control	-0.055130	-0.055130	0.011583	-0.078100	-0.032159	-0.223629	0.113370
1	PTSD	-0.053695	-0.053695	0.012780	-0.079038	-0.028352	-0.222535	0.115145

All uncorrected p-values of the tracts:

tract	p_unc
0 UF_fd	0.339379
1 CB_fd	0.308395
2 ST_fd	0.267468
3 UF_fc	0.924301
4 CB_fc	0.468343
5 ST_fc	0.936035

Uncorrected p-values for FD Covariates

tract	mri_age_demeaned
0 UF_fd	2.137453e-08
1 CB_fd	4.947984e-01
2 ST_fd	1.239794e-05

Uncorrected p-values for FC Covariates

tract	mri_age_demeaned	t1_volume_demeaned
0 UF_fc	6.716191e-01	5.290667e-28
1 CB_fc	8.939124e-02	1.203092e-29
2 ST_fc	2.712867e-07	2.747524e-09

=====
=====
Group difference of FD and log FC -- With total CTQ as additional covariate
=====
=====

Loading FD data...

=====
UF_fd
=====

Descriptive statistics (raw):

	mean	std	count
group			
PTSD	0.403997	0.016449	51
control	0.408739	0.018337	57

Assumption checks

group*mri_age_demeaned interaction not significant (p=0.4913). Homogeneity of slopes assumed.
group*CTQ_total_score_demeaned interaction not significant (p=0.5461). Homogeneity of slopes assumed.
Levene W=0.281, p=0.597 — homogeneity of variances assumed.
Breusch-Pagan p=0.5524 — homoscedasticity assumed.
Shapiro-Wilk W=0.985, p=0.273 — normality of residuals assumed.

ANCOVA results (HC3-robust, Type II) — UF_fd

	sum_sq	df	F	PR(>F)	np2
C(group)	1.034947e-04	1.0	0.436907	5.100806e-01	0.004183
mri_age_demeaned	8.646691e-03	1.0	36.502382	2.400232e-08	0.259799
CTQ_total_score_demeaned	6.106571e-07	1.0	0.002578	9.596037e-01	0.000025
Residual	2.463554e-02	104.0	NaN	NaN	0.500000
group effect not significant (p=0.5101)					

OLS coefficient table (HC3 SEs):

	coef	std err	t	P> t	[0.025	0.975]
Intercept	0.4050	0.003	151.849	0.000	0.400	0.410
C(group)[T.control]	0.0028	0.004	0.661	0.510	-0.006	0.011
mri_age_demeaned	-0.0008	0.000	-6.042	0.000	-0.001	-0.001
CTQ_total_score_demeaned	-5.233e-06	0.000	-0.051	0.960	-0.000	0.000

Adjusted group means (all covariates at their mean):

	group	adjusted_mean	mean	mean_se	mean_ci_lower	mean_ci_upper	obs_ci_lower	obs_ci_upper
0	control	0.407807	0.407807	0.002495	0.402860	0.412754	0.376888	0.438726
1	PTSD	0.405039	0.405039	0.002667	0.399749	0.410328	0.374063	0.436014

=====
CB_fd
=====

Descriptive statistics (raw):

	mean	std	count
group			
PTSD	0.372557	0.012107	51
control	0.375593	0.015093	57

Assumption checks

group*mri_age_demeaned interaction not significant (p=0.8451). Homogeneity of slopes assumed.
group*CTQ_total_score_demeaned interaction not significant (p=0.7243). Homogeneity of slopes assumed.
Levene W=1.274, p=0.262 — homogeneity of variances assumed.
Breusch-Pagan p=0.3398 — homoscedasticity assumed.
Shapiro-Wilk W=0.988, p=0.461 — normality of residuals assumed.

ANCOVA results (HC3-robust, Type II) — CB_fd

	sum_sq	df	F	PR(>F)	np2
C(group)	0.000021	1.0	0.111365	0.739267	0.001070
mri_age_demeaned	0.000072	1.0	0.375055	0.541598	0.003593
CTQ_total_score_demeaned	0.000115	1.0	0.602335	0.439450	0.005758
Residual	0.019850	104.0	NaN	NaN	0.500000
group effect not significant (p=0.7393)					

OLS coefficient table (HC3 SEs):

	coef	std err	t	P> t	[0.025	0.975]
Intercept	0.3735	0.002	183.640	0.000	0.370	0.378
C(group)[T.control]	0.0012	0.004	0.334	0.739	-0.006	0.008
mri_age_demeaned	-9.403e-05	0.000	-0.612	0.542	-0.000	0.000
CTQ_total_score_demeaned	-6.444e-05	8.3e-05	-0.776	0.439	-0.000	0.000

Adjusted group means (all covariates at their mean):

	group	adjusted_mean	mean	mean_se	mean_ci_lower	mean_ci_upper	obs_ci_lower	obs_ci_upper
0	control	0.374714	0.374714	0.002372	0.370010	0.379417	0.346916	0.402511
1	PTSD	0.373540	0.373540	0.002034	0.369507	0.377574	0.345848	0.401232

=====
ST_fd

Descriptive statistics (raw):

	mean	std	count
group			
PTSD	0.443102	0.037062	51
control	0.453445	0.032223	57

Assumption checks

group*mri_age_demeaned interaction not significant (p=0.9365). Homogeneity of slopes assumed.
group*CTQ_total_score_demeaned interaction not significant (p=0.4380). Homogeneity of slopes assumed.
Levene W=0.500, p=0.481 — homogeneity of variances assumed.
Breusch-Pagan p=0.0578 — homoscedasticity assumed.
Shapiro-Wilk W=0.995, p=0.964 — normality of residuals assumed.

ANCOVA results (HC3-robust, Type II) — ST_fd

	sum_sq	df	F	PR(>F)	np2
C(group)	0.000329	1.0	0.373246	0.542573	0.003576
mri_age_demeaned	0.017761	1.0	20.136993	0.000019	0.162216
CTQ_total_score_demeaned	0.004933	1.0	5.593119	0.019886	0.051035
Residual	0.091731	104.0	NaN	NaN	0.500000
group effect not significant (p=0.5426)					

OLS coefficient table (HC3 SEs):

	coef	std err	t	P> t	[0.025	0.975]
Intercept	0.4512	0.005	83.518	0.000	0.440	0.462
C(group)[T.control]	-0.0050	0.008	-0.611	0.543	-0.021	0.011
mri_age_demeaned	-0.0015	0.000	-4.487	0.000	-0.002	-0.001
CTQ_total_score_demeaned	-0.0005	0.000	-2.365	0.020	-0.001	-7.54e-05

Adjusted group means (all covariates at their mean):

	group	adjusted_mean	mean	mean_se	mean_ci_lower	mean_ci_upper	obs_ci_lower	obs_ci_upper
0	control	0.446188	0.446188	0.004697	0.436875	0.455502	0.386562	0.505814
1	PTSD	0.451213	0.451213	0.005403	0.440500	0.461927	0.391352	0.511074

Loading FC data...

UF_fc

Descriptive statistics (raw):

	mean	std	count
group			
PTSD	0.054032	0.069023	51
control	0.084932	0.077414	57

Assumption checks

group*mri_age_demeaned interaction not significant (p=0.8625). Homogeneity of slopes assumed.
group*t1_volume_demeaned interaction not significant (p=0.6763). Homogeneity of slopes assumed.
group*CTQ_total_score_demeaned interaction not significant (p=0.4770). Homogeneity of slopes assumed.
Levene W=1.288, p=0.259 — homogeneity of variances assumed.
Breusch-Pagan p=0.3494 — homoscedasticity assumed.
Shapiro-Wilk W=0.989, p=0.560 — normality of residuals assumed.

ANCOVA results (HC3-robust, Type II) — UF_fc

	sum_sq	df	F	PR(>F)	np2
C(group)	0.000403	1.0	0.246875	6.203427e-01	0.002391
mri_age_demeaned	0.000402	1.0	0.246202	6.208193e-01	0.002385
t1_volume_demeaned	0.369379	1.0	226.030941	9.944010e-28	0.686960
CTQ_total_score_demeaned	0.000740	1.0	0.453076	5.023851e-01	0.004380
Residual	0.168322	103.0	NaN	NaN	0.500000
group effect not significant (p=0.6203)					

OLS coefficient table (HC3 SEs):

	coef	std err	t	P> t	[0.025	0.975]
Intercept	0.0672	0.007	8.954	0.000	0.052	0.082
C(group)[T.control]	0.0060	0.012	0.497	0.620	-0.018	0.030
mri_age_demeaned	-0.0002	0.000	-0.496	0.621	-0.001	0.001
t1_volume_demeaned	4.832e-07	3.21e-08	15.034	0.000	4.19e-07	5.47e-07
CTQ_total_score_demeaned	0.0002	0.000	0.673	0.502	-0.000	0.001

Adjusted group means (all covariates at their mean):

	group	adjusted_mean	mean	mean_se	mean_ci_lower	mean_ci_upper	obs_ci_lower	obs_ci_upper
0	control	0.073194	0.073194	0.007071	0.059170	0.087219	-0.008197	0.154586
1	PTSD	0.067151	0.067151	0.007500	0.052276	0.082025	-0.014391	0.148693

CB_fc

Descriptive statistics (raw):

	mean	std	count
group			
PTSD	0.017148	0.075854	51
control	0.043794	0.084570	57

Assumption checks

group*mri_age_demeaned interaction not significant (p=0.5255). Homogeneity of slopes assumed.
group*t1_volume_demeaned interaction not significant (p=0.5010). Homogeneity of slopes assumed.
group*CTQ_total_score_demeaned interaction not significant (p=0.7573). Homogeneity of slopes assumed.
Levene W=0.953, p=0.331 — homogeneity of variances assumed.
Breusch-Pagan p=0.3915 — homoscedasticity assumed.
Shapiro-Wilk W=0.996, p=0.984 — normality of residuals assumed.

ANCOVA results (HC3-robust, Type II) — CB_fc

	sum_sq	df	F	PR(>F)	np2
C(group)	0.001267	1.0	0.813841	3.690919e-01	0.007839
mri_age_demeaned	0.004748	1.0	3.048940	8.377138e-02	0.028750
t1_volume_demeaned	0.373956	1.0	240.121123	1.137083e-28	0.699814
CTQ_total_score_demeaned	0.000534	1.0	0.342708	5.595506e-01	0.003316
Residual	0.160409	103.0	NaN	NaN	0.500000
group effect not significant (p=0.3691)					

OLS coefficient table (HC3 SEs):

	coef	std err	t	P> t	[0.025	0.975]
Intercept	0.0366	0.008	4.723	0.000	0.021	0.052
C(group)[T.control]	-0.0102	0.011	-0.902	0.369	-0.032	0.012
mri_age_demeaned	0.0006	0.000	1.746	0.084	-8.36e-05	0.001
t1_volume_demeaned	5.53e-07	3.57e-08	15.496	0.000	4.82e-07	6.24e-07
CTQ_total_score_demeaned	-0.0002	0.000	-0.585	0.560	-0.001	0.000

Adjusted group means (all covariates at their mean):

	group	adjusted_mean	mean	mean_se	mean_ci_lower	mean_ci_upper	obs_ci_lower	obs_ci_upper
0	control	0.026414	0.026414	0.005924	0.014665	0.038164	-0.052729	0.105558
1	PTSD	0.036573	0.036573	0.007743	0.021217	0.051929	-0.043186	0.116331

ST_fc

Descriptive statistics (raw):

	mean	std	count
group			
PTSD	-0.063659	0.117743	51
control	-0.046215	0.110523	57

Assumption checks

group*mri_age_demeaned interaction not significant (p=0.4158). Homogeneity of slopes assumed.
group*t1_volume_demeaned interaction not significant (p=0.3327). Homogeneity of slopes assumed.
group*CTQ_total_score_demeaned interaction not significant (p=0.8782). Homogeneity of slopes assumed.
Levene W=0.478, p=0.491 — homogeneity of variances assumed.
Breusch-Pagan p=0.9326 — homoscedasticity assumed.
Shapiro-Wilk W=0.983, p=0.176 — normality of residuals assumed.

/home/amygdala/Desktop/Mannheim/repository/code/group_difference.py:135: RuntimeWarning: More than 20 figures have been opened. Figures created through the pyplot interface (matplotlib.pyplot.figure) are retained until explicitly closed and may consume too much memory. (To control this warning, see the rcParam `figure.max\_open\_warning`). Consider using `matplotlib.pyplot.close()`.
plt.figure(figsize=(7, 5))

ANCOVA results (HC3-robust, Type II) — ST_fc

	sum_sq	df	F	PR(>F)	np2
C(group)	0.000762	1.0	0.106931	7.443295e-01	0.001037
mri_age_demeaned	0.209193	1.0	29.372949	3.947981e-07	0.221895
t1_volume_demeaned	0.281773	1.0	39.563963	7.802889e-09	0.277517
CTQ_total_score_demeaned	0.002888	1.0	0.405502	5.256744e-01	0.003921
Residual	0.733562	103.0	NaN	NaN	0.500000
group effect not significant (p=0.7443)					

OLS coefficient table (HC3 SEs):

	coef	std err	t	P> t	[0.025	0.975]
Intercept	-0.0587	0.016	-3.730	0.000	-0.090	-0.027
C(group)[T.control]	0.0080	0.024	0.327	0.744	-0.041	0.057
mri_age_demeaned	0.0046	0.001	5.420	0.000	0.003	0.006
t1_volume_demeaned	4.803e-07	7.64e-08	6.290	0.000	3.29e-07	6.32e-07
CTQ_total_score_demeaned	0.0004	0.001	0.637	0.526	-0.001	0.002

Adjusted group means (all covariates at their mean):

	group	adjusted_mean	mean	mean_se	mean_ci_lower	mean_ci_upper	obs_ci_lower	obs_ci_upper
0	control	-0.050669	-0.050669	0.013940	-0.078317	-0.023022	-0.220309	0.118970
1	PTSD	-0.058680	-0.058680	0.015733	-0.089883	-0.027476	-0.228935	0.111575

All uncorrected p-values of the tracts:

tract	p_unc
0 UF_fd	0.510081
1 CB_fd	0.739267

2 ST_fd 0.542573
3 UF_fc 0.620343
4 CB_fc 0.369092
5 ST_fc 0.744329

Uncorrected p-values for FD Covariates

tract mri_age_demeaned CTQ_total_score_demeaned

0	UF_fd	2.400232e-08	0.959604
1	CB_fd	5.415983e-01	0.439450
2	ST_fd	1.866890e-05	0.019886

Uncorrected p-values for FC Covariates

tract mri_age_demeaned t1_volume_demeaned CTQ_total_score_demeaned

0	UF_fc	6.208193e-01	9.944010e-28	0.502385
1	CB_fc	8.377138e-02	1.137083e-28	0.559551
2	ST_fc	3.947981e-07	7.802889e-09	0.525674