

# stochastics and probability

## Lecture 3

Dr. Johannes Pahlke

# accept-reject sampling

# accept-reject sampling

want:

# accept-reject sampling

want:

$X$  with PDF  $f_X(x)$

# accept-reject sampling

want:

$X$  with PDF  $f_X(x)$

use:

# accept-reject sampling

want:

$X$  with PDF  $f_X(x)$

use:

$$f_X(x) \leq \alpha$$

# accept-reject sampling

want:

$X$  with PDF  $f_X(x)$

use:

$$f_X(x) \leq \alpha \quad \text{Range}(X) \subseteq [\beta_1, \beta_2]$$


# accept-reject sampling

want:

$X$  with PDF  $f_X(x)$

use:

$$f_X(x) \leq \alpha \quad \text{Range}(X) \subseteq [\beta_1, \beta_2]$$





# accept-reject sampling


want:

$X$  with PDF  $f_X(x)$

use:

$$f_X(x) \leq \alpha \quad \text{Range}(X) \subseteq [\beta_1, \beta_2]$$

algorithm:



# accept-reject sampling

want:


$X$  with PDF  $f_X(x)$

use:

$$f_X(x) \leq \alpha \quad \text{Range}(X) \subseteq [\beta_1, \beta_2]$$

algorithm:

1.) draw  $U_1 \sim \mathcal{U}(\beta_1, \beta_2)$ ,  $U_2 \sim \mathcal{U}(0, \alpha)$



# accept-reject sampling

want:


$X$  with PDF  $f_X(x)$

use:

$$f_X(x) \leq \alpha \quad \text{Range}(X) \subseteq [\beta_1, \beta_2]$$

algorithm:

- 1.) draw  $U_1 \sim \mathcal{U}(\beta_1, \beta_2)$ ,  $U_2 \sim \mathcal{U}(0, \alpha)$
- 2.) if  $U_2 \leq f_X(U_1)$ , then  $X = U_1$ , else go to 1.)



# accept-reject sampling

want:



$X$  with PDF  $f_X(x)$

use:

$$f_X(x) \leq \alpha \quad \text{Range}(X) \subseteq [\beta_1, \beta_2]$$

algorithm:

- 1.) draw  $U_1 \sim \mathcal{U}(\beta_1, \beta_2)$ ,  $U_2 \sim \mathcal{U}(0, \alpha)$
- 2.) if  $U_2 \leq f_X(U_1)$ , then  $X = U_1$ , else go to 1.)



# accept-reject sampling

want:



$X$  with PDF  $f_X(x)$

use:

$$f_X(x) \leq \alpha \quad \text{Range}(X) \subseteq [\beta_1, \beta_2]$$

algorithm:

- 1.) draw  $U_1 \sim \mathcal{U}(\beta_1, \beta_2)$ ,  $U_2 \sim \mathcal{U}(0, \alpha)$
- 2.) if  $U_2 \leq f_X(U_1)$ , then  $X = U_1$ , else go to 1.)



inefficient:

# accept-reject sampling

want:



$X$  with PDF  $f_X(x)$

use:

$$f_X(x) \leq \alpha \quad \text{Range}(X) \subseteq [\beta_1, \beta_2]$$

algorithm:

- 1.) draw  $U_1 \sim \mathcal{U}(\beta_1, \beta_2)$ ,  $U_2 \sim \mathcal{U}(0, \alpha)$
- 2.) if  $U_2 \leq f_X(U_1)$ , then  $X = U_1$ , else go to 1.)



inefficient:

- peaky densities

# accept-reject sampling

want:



$X$  with PDF  $f_X(x)$

use:

$$f_X(x) \leq \alpha \quad \text{Range}(X) \subseteq [\beta_1, \beta_2]$$

algorithm:

- 1.) draw  $U_1 \sim \mathcal{U}(\beta_1, \beta_2)$ ,  $U_2 \sim \mathcal{U}(0, \alpha)$
- 2.) if  $U_2 \leq f_X(U_1)$ , then  $X = U_1$ , else go to 1.)



inefficient:

- peaky densities
- high dimensions

# accept-reject sampling

want:



$X$  with PDF  $f_X(x)$

use:

$$f_X(x) \leq \alpha \quad \text{Range}(X) \subseteq [\beta_1, \beta_2]$$

algorithm:

- 1.) draw  $U_1 \sim \mathcal{U}(\beta_1, \beta_2)$ ,  $U_2 \sim \mathcal{U}(0, \alpha)$
- 2.) if  $U_2 \leq f_X(U_1)$ , then  $X = U_1$ , else go to 1.)



inefficient:

- peaky densities
- high dimensions
- long tails/infinite Range



# accept-reject sampling

# accept-reject sampling

want:

# accept-reject sampling

want:

$X$  with PDF  $f_X(x)$

# accept-reject sampling

want:

$X$  with PDF  $f_X(x)$

use:

# accept-reject sampling

want:

$X$  with PDF  $f_X(x)$

use:

$G \sim \mathcal{G}$  with PDF  $g_G(x)$

# accept-reject sampling

want:

$X$  with PDF  $f_X(x)$

use:

$G \sim \mathcal{G}$  with PDF  $g_G(x)$  and  $f_X(x) \leq c g_G(x)$

# accept-reject sampling

want:

$X$  with PDF  $f_X(x)$

use:

$G \sim \mathcal{G}$  with PDF  $g_G(x)$  and  $f_X(x) \leq c g_G(x)$  for all  $x$


# accept-reject sampling

want:

$X$  with PDF  $f_X(x)$

use:

$G \sim \mathcal{G}$  with PDF  $g_G(x)$  and  $f_X(x) \leq c g_G(x)$  for all  $x$





# accept-reject sampling


want:

$X$  with PDF  $f_X(x)$

use:

$G \sim \mathcal{G}$  with PDF  $g_G(x)$  and  $f_X(x) \leq c g_G(x)$  for all  $x$

algorithm:



# accept-reject sampling

want:


$X$  with PDF  $f_X(x)$

use:

$G \sim \mathcal{G}$  with PDF  $g_G(x)$  and  $f_X(x) \leq c g_G(x)$  for all  $x$

algorithm:

1.) draw  $G$  from  $\mathcal{G}$  and  $U$  from  $\mathcal{U}(0,1)$



# accept-reject sampling

want:


$X$  with PDF  $f_X(x)$

use:

$G \sim \mathcal{G}$  with PDF  $g_G(x)$  and  $f_X(x) \leq c g_G(x)$  for all  $x$

algorithm:

- 1.) draw  $G$  from  $\mathcal{G}$  and  $U$  from  $\mathcal{U}(0,1)$
- 2.) if  $U \leq \frac{f_X(G)}{c g_G(G)}$ , then  $X = G$ , else go to 1.)



# discrete-time stochastic processes

# discrete-time stochastic processes

$$n \in \mathbb{N}$$

# discrete-time stochastic processes

$n \in \mathbb{N}$       discrete Time

# discrete-time stochastic processes

$n \in \mathbb{N}$       discrete Time

$X_0, X_1, X_2, \dots$

# discrete-time stochastic processes

$n \in \mathbb{N}$       discrete Time

$X_0, X_1, X_2, \dots$       "sequence of random variables"



# discrete-time stochastic processes

$n \in \mathbb{N}$       discrete Time

$X_0, X_1, X_2, \dots$       "sequence of random variables"

$\neq x_0, x_1, x_2, \dots$

# discrete-time stochastic processes

$n \in \mathbb{N}$       discrete Time

$X_0, X_1, X_2, \dots$       "sequence of random variables"

$\neq x_0, x_1, x_2, \dots$       "sequence of random numbers"

# discrete-time stochastic processes

$n \in \mathbb{N}$       discrete Time

$X_0, X_1, X_2, \dots$       "sequence of random variables"

$\neq x_0, x_1, x_2, \dots$       "sequence of random numbers"

$\{X_n : n \in \mathbb{N}\}$

# discrete-time stochastic processes

$n \in \mathbb{N}$       discrete Time

$X_0, X_1, X_2, \dots$       "sequence of random variables"

$\neq x_0, x_1, x_2, \dots$       "sequence of random numbers"

$\{X_n : n \in \mathbb{N}\}$       discrete-time stochastic process

# discrete-time stochastic processes

$n \in \mathbb{N}$       discrete Time

$X_0, X_1, X_2, \dots$       "sequence of random variables"

$\neq x_0, x_1, x_2, \dots$       "sequence of random numbers"

$\{X_n : n \in \mathbb{N}\}$       discrete-time stochastic process

$x_n$

# discrete-time stochastic processes

$n \in \mathbb{N}$       discrete Time

$X_0, X_1, X_2, \dots$       "sequence of random variables"

$\neq x_0, x_1, x_2, \dots$       "sequence of random numbers"

$\{X_n : n \in \mathbb{N}\}$       discrete-time stochastic process

$x_n$       value of random variable  $X_n$

# discrete-time stochastic processes

$n \in \mathbb{N}$       discrete Time

$X_0, X_1, X_2, \dots$       "sequence of random variables"

$\neq x_0, x_1, x_2, \dots$       "sequence of random numbers"

$\{X_n : n \in \mathbb{N}\}$       discrete-time stochastic process

$x_n$       value of random variable  $X_n$

$X_n : \Omega \rightarrow \mathbb{S}$

# discrete-time stochastic processes

$n \in \mathbb{N}$       discrete Time

$X_0, X_1, X_2, \dots$       "sequence of random variables"

$\neq x_0, x_1, x_2, \dots$       "sequence of random numbers"

$\{X_n : n \in \mathbb{N}\}$       discrete-time stochastic process

$x_n$       value of random variable  $X_n$

$X_n : \Omega \rightarrow \mathbb{S}$

$x_n \in \mathbb{S}$



# discrete-time stochastic processes

$n \in \mathbb{N}$       discrete Time

$X_0, X_1, X_2, \dots$       "sequence of random variables"

$\neq x_0, x_1, x_2, \dots$       "sequence of random numbers"

$\{X_n : n \in \mathbb{N}\}$       discrete-time stochastic process

$x_n$       value of random variable  $X_n$

$X_n : \Omega \rightarrow \mathbb{S}$       state space  $\mathbb{S}$   
 $x_n \in \mathbb{S}$       (discrete or continuous)

# discrete-time stochastic processes

$n \in \mathbb{N}$       discrete Time

$X_0, X_1, X_2, \dots$       "sequence of random variables"

$\neq x_0, x_1, x_2, \dots$       "sequence of random numbers"

$\{X_n : n \in \mathbb{N}\}$       discrete-time stochastic process

$x_n$       value of random variable  $X_n$

$X_n : \Omega \rightarrow \mathbb{S}$       state space  $\mathbb{S}$   
 $x_n \in \mathbb{S}$       (discrete or continuous)

Examples:

# discrete-time stochastic processes

$n \in \mathbb{N}$       discrete Time

$X_0, X_1, X_2, \dots$       "sequence of random variables"

$\neq x_0, x_1, x_2, \dots$       "sequence of random numbers"

$\{X_n : n \in \mathbb{N}\}$       discrete-time stochastic process

$x_n$       value of random variable  $X_n$

$X_n : \Omega \rightarrow \mathbb{S}$       state space  $\mathbb{S}$   
 $x_n \in \mathbb{S}$       (discrete or continuous)

Examples:

population

# discrete-time stochastic processes

$n \in \mathbb{N}$       discrete Time

$X_0, X_1, X_2, \dots$       "sequence of random variables"


$\neq x_0, x_1, x_2, \dots$       "sequence of random numbers"

$\{X_n : n \in \mathbb{N}\}$       discrete-time stochastic process

$x_n$       value of random variable  $X_n$

$X_n : \Omega \rightarrow \mathbb{S}$       state space  $\mathbb{S}$   
 $x_n \in \mathbb{S}$       (discrete or continuous)

Examples:



# discrete-time stochastic processes

$n \in \mathbb{N}$  discrete Time

$X_0, X_1, X_2, \dots$  "sequence of random variables"

$\neq x_0, x_1, x_2, \dots$  "sequence of random numbers"


$\{X_n : n \in \mathbb{N}\}$  discrete-time stochastic process

$x_n$  value of random variable  $X_n$

$X_n : \Omega \rightarrow \mathbb{S}$  state space  $\mathbb{S}$   
 $x_n \in \mathbb{S}$  (discrete or continuous)

Examples:

population



# discrete-time stochastic processes

$n \in \mathbb{N}$  discrete Time

$X_0, X_1, X_2, \dots$  "sequence of random variables"

$\neq x_0, x_1, x_2, \dots$  "sequence of random numbers"


$\{X_n : n \in \mathbb{N}\}$  discrete-time stochastic process

$x_n$  value of random variable  $X_n$

$X_n : \Omega \rightarrow \mathbb{S}$  state space  $\mathbb{S}$   
 $x_n \in \mathbb{S}$  (discrete or continuous)

Examples:

population



rolling dice

# discrete-time stochastic processes

$n \in \mathbb{N}$  discrete Time

$X_0, X_1, X_2, \dots$  "sequence of random variables"

$\neq x_0, x_1, x_2, \dots$  "sequence of random numbers"


$\{X_n : n \in \mathbb{N}\}$  discrete-time stochastic process

$x_n$  value of random variable  $X_n$


$X_n : \Omega \rightarrow \mathbb{S}$  state space  $\mathbb{S}$   
 $x_n \in \mathbb{S}$  (discrete or continuous)

Examples:

population



rolling dice



1st



# discrete-time stochastic processes

$n \in \mathbb{N}$  discrete Time

$X_0, X_1, X_2, \dots$  "sequence of random variables"

$\neq x_0, x_1, x_2, \dots$  "sequence of random numbers"


$\{X_n : n \in \mathbb{N}\}$  discrete-time stochastic process

$x_n$  value of random variable  $X_n$


$X_n : \Omega \rightarrow \mathbb{S}$  state space  $\mathbb{S}$   
 $x_n \in \mathbb{S}$  (discrete or continuous)

Examples:

population



rolling dice



1st

2nd



# discrete-time stochastic processes

$n \in \mathbb{N}$  discrete Time

$X_0, X_1, X_2, \dots$  "sequence of random variables"

$\neq x_0, x_1, x_2, \dots$  "sequence of random numbers"


$\{X_n : n \in \mathbb{N}\}$  discrete-time stochastic process

$x_n$  value of random variable  $X_n$


$X_n : \Omega \rightarrow \mathbb{S}$  state space  $\mathbb{S}$   
 $x_n \in \mathbb{S}$  (discrete or continuous)

Examples:

population



rolling dice



**independent identical distributed (i.i.d.) process**

# independent identical distributed (i.i.d.) process

$$\{X_n : n \in \mathbb{N}\}$$

# independent identical distributed (i.i.d.) process

$\{X_n : n \in \mathbb{N}\}$       stochastic process

# independent identical distributed (i.i.d.) process

$\{X_n : n \in \mathbb{N}\}$       stochastic process

$$F_{X_0}(x) = F_{X_1}(x) = F_{X_2}(x) = \dots$$

# independent identical distributed (i.i.d.) process

$\{X_n : n \in \mathbb{N}\}$       stochastic process

$F_{X_0}(x) = F_{X_1}(x) = F_{X_2}(x) = \dots$       identical distributed

# independent identical distributed (i.i.d.) process

$\{X_n : n \in \mathbb{N}\}$       stochastic process

$F_{X_0}(x) = F_{X_1}(x) = F_{X_2}(x) = \dots$       identical distributed

$$F_{X_0, \dots, X_n}(x_0, \dots, x_n) = F_{X_0}(x_0) \cdot \dots \cdot F_{X_n}(x_n)$$

# independent identical distributed (i.i.d.) process

$\{X_n : n \in \mathbb{N}\}$  stochastic process

$F_{X_0}(x) = F_{X_1}(x) = F_{X_2}(x) = \dots$  identical distributed

$F_{X_0, \dots, X_n}(x_0, \dots, x_n) = F_{X_0}(x_0) \cdot \dots \cdot F_{X_n}(x_n)$  independent



# independent identical distributed (i.i.d.) process

$\{X_n : n \in \mathbb{N}\}$  stochastic process

$F_{X_0}(x) = F_{X_1}(x) = F_{X_2}(x) = \dots$  identical distributed

$F_{X_0, \dots, X_n}(x_0, \dots, x_n) = F_{X_0}(x_0) \cdot \dots \cdot F_{X_n}(x_n)$  independent

exercise:

# independent identical distributed (i.i.d.) process

$\{X_n : n \in \mathbb{N}\}$  stochastic process

$F_{X_0}(x) = F_{X_1}(x) = F_{X_2}(x) = \dots$  identical distributed

$F_{X_0, \dots, X_n}(x_0, \dots, x_n) = F_{X_0}(x_0) \cdot \dots \cdot F_{X_n}(x_n)$  independent

## exercise:

Which system can be captured by an i.i.d. process?

# independent identical distributed (i.i.d.) process

$\{X_n : n \in \mathbb{N}\}$  stochastic process


$F_{X_0}(x) = F_{X_1}(x) = F_{X_2}(x) = \dots$  identical distributed

$F_{X_0, \dots, X_n}(x_0, \dots, x_n) = F_{X_0}(x_0) \cdot \dots \cdot F_{X_n}(x_n)$  independent

## exercise:

Which system can be captured by an i.i.d. process?

coin-flip



# independent identical distributed (i.i.d.) process

$\{X_n : n \in \mathbb{N}\}$  stochastic process


$F_{X_0}(x) = F_{X_1}(x) = F_{X_2}(x) = \dots$  identical distributed

$F_{X_0, \dots, X_n}(x_0, \dots, x_n) = F_{X_0}(x_0) \cdot \dots \cdot F_{X_n}(x_n)$  independent


## exercise:

Which system can be captured by an i.i.d. process?

coin-flip



population development



2022

2023

# independent identical distributed (i.i.d.) process

$\{X_n : n \in \mathbb{N}\}$  stochastic process


$F_{X_0}(x) = F_{X_1}(x) = F_{X_2}(x) = \dots$  identical distributed

$F_{X_0, \dots, X_n}(x_0, \dots, x_n) = F_{X_0}(x_0) \cdot \dots \cdot F_{X_n}(x_n)$  independent


## exercise:

Which system can be captured by an i.i.d. process?

coin-flip



population development




2022

2023

5

rolling dice



# independent identical distributed (i.i.d.) process

$\{X_n : n \in \mathbb{N}\}$  stochastic process


$F_{X_0}(x) = F_{X_1}(x) = F_{X_2}(x) = \dots$  identical distributed

$F_{X_0, \dots, X_n}(x_0, \dots, x_n) = F_{X_0}(x_0) \cdot \dots \cdot F_{X_n}(x_n)$  independent


## exercise:

Which system can be captured by an i.i.d. process?

coin-flip



population development




2022

2023

5

rolling dice



# independent identical distributed (i.i.d.) process

$\{X_n : n \in \mathbb{N}\}$  stochastic process


$F_{X_0}(x) = F_{X_1}(x) = F_{X_2}(x) = \dots$  identical distributed

$F_{X_0, \dots, X_n}(x_0, \dots, x_n) = F_{X_0}(x_0) \cdot \dots \cdot F_{X_n}(x_n)$  independent


## exercise:

Which system can be captured by an i.i.d. process?

coin-flip




population development



5

rolling dice



# Markov chains



# Markov chains

$$\{X_n : n \in \mathbb{N}\}$$

# Markov chains

$$\{X_n : n \in \mathbb{N}\}$$

stochastic process

# Markov chains

$$\{X_n : n \in \mathbb{N}\}$$

stochastic process

$$P(X_{n+1} = x_j | X_n = x_i, X_{n-1} = x_{i-1}, \dots, X_0 = x_0)$$

# Markov chains

$$\{X_n : n \in \mathbb{N}\}$$

stochastic process

$$\begin{aligned} P(X_{n+1} = x_j | X_n = x_i, X_{n-1} = x_{i-1}, \dots, X_0 = x_0) \\ = P(X_{n+1} = x_j | X_n = x_i) \end{aligned}$$

# Markov chains

$$\{X_n : n \in \mathbb{N}\}$$

stochastic process

$$\begin{aligned} P(X_{n+1} = x_j | X_n = x_i, X_{n-1} = x_{i-1}, \dots, X_0 = x_0) \\ = P(X_{n+1} = x_j | X_n = x_i) \end{aligned}$$

Markov property

( $X_{n+1}$  only depends on  $X_n$ )

# Markov chains

$$\{X_n : n \in \mathbb{N}\}$$

stochastic process

$$\begin{aligned} P(X_{n+1} = x_j | X_n = x_i, X_{n-1} = x_{i-1}, \dots, X_0 = x_0) \\ = P(X_{n+1} = x_j | X_n = x_i) =: P_{ij} \end{aligned}$$

Markov property

( $X_{n+1}$  only depends on  $X_n$ )

# Markov chains

$$\{X_n : n \in \mathbb{N}\}$$

stochastic process

$$\begin{aligned} P(X_{n+1} = x_j | X_n = x_i, X_{n-1} = x_{i-1}, \dots, X_0 = x_0) \\ = P(X_{n+1} = x_j | X_n = x_i) =: P_{ij} \\ \uparrow \\ \text{(one-step transition probability)} \end{aligned}$$

Markov property

( $X_{n+1}$  only depends on  $X_n$ )

# Markov chains

$$\{X_n : n \in \mathbb{N}\}$$

stochastic process

$$\begin{aligned} P(X_{n+1} = x_j | X_n = x_i, X_{n-1} = x_{i-1}, \dots, X_0 = x_0) \\ = P(X_{n+1} = x_j | X_n = x_i) =: P_{ij} \\ \uparrow \\ \text{(one-step transition probability)} \end{aligned}$$

Markov property

( $X_{n+1}$  only depends on  $X_n$ )

$$P_{ij}^{(n)} := P(X_{n+m} = x_j | X_m = x_i)$$



# Markov chains

$$\{X_n : n \in \mathbb{N}\}$$

stochastic process

$$\begin{aligned} P(X_{n+1} = x_j | X_n = x_i, X_{n-1} = x_{i-1}, \dots, X_0 = x_0) \\ = P(X_{n+1} = x_j | X_n = x_i) =: P_{ij} \end{aligned}$$

↑  
(one-step transition probability)

Markov property  
( $X_{n+1}$  only depends on  $X_n$ )

$$P_{ij}^{(n)} := P(X_{n+m} = x_j | X_m = x_i)$$

( $n$ -step transition probability)

# Markov chains

$$\{X_n : n \in \mathbb{N}\}$$

stochastic process

$$\begin{aligned} P(X_{n+1} = x_j | X_n = x_i, X_{n-1} = x_{i-1}, \dots, X_0 = x_0) \\ = P(X_{n+1} = x_j | X_n = x_i) =: P_{ij} \end{aligned}$$

↑  
(one-step transition probability)

Markov property  
( $X_{n+1}$  only depends on  $X_n$ )

$$P_{ij}^{(n)} := P(X_{n+m} = x_j | X_m = x_i)$$

( $n$ -step transition probability)

$$\underline{\underline{P}} := (P_{ij})$$

# Markov chains

$$\{X_n : n \in \mathbb{N}\}$$

stochastic process

$$\begin{aligned} P(X_{n+1} = x_j | X_n = x_i, X_{n-1} = x_{i-1}, \dots, X_0 = x_0) \\ = P(X_{n+1} = x_j | X_n = x_i) =: P_{ij} \end{aligned}$$

↑  
(one-step transition probability)

Markov property  
( $X_{n+1}$  only depends on  $X_n$ )

$$P_{ij}^{(n)} := P(X_{n+m} = x_j | X_m = x_i)$$

( $n$ -step transition probability)

$$\underline{\underline{P}} := (P_{ij})$$

(one-step transition matrix)

# Markov chains

$$\{X_n : n \in \mathbb{N}\}$$

stochastic process

$$\begin{aligned} P(X_{n+1} = x_j | X_n = x_i, X_{n-1} = x_{i-1}, \dots, X_0 = x_0) \\ = P(X_{n+1} = x_j | X_n = x_i) =: P_{ij} \end{aligned}$$

(one-step transition probability)

Markov property

( $X_{n+1}$  only depends on  $X_n$ )

$$P_{ij}^{(n)} := P(X_{n+m} = x_j | X_m = x_i)$$

( $n$ -step transition probability)

$$\underline{\underline{P}} := (P_{ij}) = \begin{pmatrix} P_{00} & \dots & P_{0n} \\ \vdots & \ddots & \vdots \\ P_{n0} & \dots & P_{nn} \end{pmatrix}$$

(one-step transition matrix)

# Markov chains

$$\{X_n : n \in \mathbb{N}\}$$

stochastic process

$$\begin{aligned} P(X_{n+1} = x_j | X_n = x_i, X_{n-1} = x_{i-1}, \dots, X_0 = x_0) \\ = P(X_{n+1} = x_j | X_n = x_i) =: P_{ij} \end{aligned}$$

(one-step transition probability)

Markov property  
( $X_{n+1}$  only depends on  $X_n$ )

$$P_{ij}^{(n)} := P(X_{n+m} = x_j | X_m = x_i)$$

( $n$ -step transition probability)

$$\underline{\underline{P}} := (P_{ij}) = \begin{pmatrix} P_{00} & \dots & P_{0n} \\ \vdots & \ddots & \vdots \\ P_{n0} & \dots & P_{nn} \end{pmatrix} \text{ for } |\mathbb{S}| < \infty$$

(one-step transition matrix)

# Markov chains

$$\{X_n : n \in \mathbb{N}\}$$

stochastic process

$$\begin{aligned} P(X_{n+1} = x_j | X_n = x_i, X_{n-1} = x_{i-1}, \dots, X_0 = x_0) \\ = P(X_{n+1} = x_j | X_n = x_i) =: P_{ij} \end{aligned}$$

(one-step transition probability)

Markov property

( $X_{n+1}$  only depends on  $X_n$ )

$$P_{ij}^{(n)} := P(X_{n+m} = x_j | X_m = x_i)$$

( $n$ -step transition probability)

example:

$$\underline{\underline{P}} := (P_{ij}) = \begin{pmatrix} P_{00} & \dots & P_{0n} \\ \vdots & \ddots & \vdots \\ P_{n0} & \dots & P_{nn} \end{pmatrix} \text{ for } |\mathbb{S}| < \infty$$

(one-step transition matrix)

# Markov chains

$$\{X_n : n \in \mathbb{N}\}$$

stochastic process

$$\begin{aligned} P(X_{n+1} = x_j | X_n = x_i, X_{n-1} = x_{i-1}, \dots, X_0 = x_0) \\ = P(X_{n+1} = x_j | X_n = x_i) =: P_{ij} \end{aligned}$$

↑  
(one-step transition probability)

Markov property  
( $X_{n+1}$  only depends on  $X_n$ )


$$P_{ij}^{(n)} := P(X_{n+m} = x_j | X_m = x_i)$$

( $n$ -step transition probability)

$$\underline{P} := (P_{ij}) = \begin{pmatrix} P_{00} & \dots & P_{0n} \\ \vdots & \ddots & \vdots \\ P_{n0} & \dots & P_{nn} \end{pmatrix} \text{ for } |\mathcal{S}| < \infty$$

(one-step transition matrix)

example:



# Markov chains

$$\{X_n : n \in \mathbb{N}\}$$

stochastic process

$$\begin{aligned} P(X_{n+1} = x_j | X_n = x_i, X_{n-1} = x_{i-1}, \dots, X_0 = x_0) \\ = P(X_{n+1} = x_j | X_n = x_i) =: P_{ij} \end{aligned}$$

↑  
(one-step transition probability)

Markov property  
( $X_{n+1}$  only depends on  $X_n$ )


$$P_{ij}^{(n)} := P(X_{n+m} = x_j | X_m = x_i)$$

( $n$ -step transition probability)

$$\underline{\underline{P}} := (P_{ij}) = \begin{pmatrix} P_{00} & \dots & P_{0n} \\ \vdots & \ddots & \vdots \\ P_{n0} & \dots & P_{nn} \end{pmatrix} \text{ for } |\mathbb{S}| < \infty$$

(one-step transition matrix)

example:



$$\underline{\underline{P}} = \begin{pmatrix} 0.75 & 0.12 & 0.13 \\ 0.21 & 0.56 & 0.23 \\ 0.31 & 0.32 & 0.37 \end{pmatrix}$$



# Markov chains as recursions

# Markov chains as recursions

theorem 1:

# Markov chains as recursions

theorem 1:

Let

# Markov chains as recursions

theorem 1:

Let  $g(x, u) \in \mathbb{R}$

# Markov chains as recursions

theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and

# Markov chains as recursions

theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$

# Markov chains as recursions

theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,

# Markov chains as recursions

theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,  
then the recursion



# Markov chains as recursions

theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,

then the recursion  $X_{n+1} = g(X_n, U_n)$

# Markov chains as recursions

theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,  
then the recursion  $X_{n+1} = g(X_n, U_n)$  is a Markov chain.

# Markov chains as recursions

theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,

then the recursion  $X_{n+1} = g(X_n, U_n)$  is a Markov chain.

theorem 2:

# Markov chains as recursions

theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,  
then the recursion  $X_{n+1} = g(X_n, U_n)$  is a Markov chain.

theorem 2:

Every Markov chain can be represented as recursion

# Markov chains as recursions

theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,  
then the recursion  $X_{n+1} = g(X_n, U_n)$  is a Markov chain.

theorem 2:

Every Markov chain can be represented as recursion  $X_{n+1} = g(X_n, U_n)$

# Markov chains as recursions

theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,  
then the recursion  $X_{n+1} = g(X_n, U_n)$  is a Markov chain.

theorem 2:

Every Markov chain can be represented as recursion  $X_{n+1} = g(X_n, U_n)$   
where

# Markov chains as recursions

theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,  
then the recursion  $X_{n+1} = g(X_n, U_n)$  is a Markov chain.

theorem 2:

Every Markov chain can be represented as recursion  $X_{n+1} = g(X_n, U_n)$

where  $\{U_n : n \in \mathbb{N}\}$

# Markov chains as recursions

theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,  
then the recursion  $X_{n+1} = g(X_n, U_n)$  is a Markov chain.

theorem 2:

Every Markov chain can be represented as recursion  $X_{n+1} = g(X_n, U_n)$   
where  $\{U_n : n \in \mathbb{N}\}$  is an i.i.d. process with



# Markov chains as recursions

theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,  
then the recursion  $X_{n+1} = g(X_n, U_n)$  is a Markov chain.

theorem 2:

Every Markov chain can be represented as recursion  $X_{n+1} = g(X_n, U_n)$   
where  $\{U_n : n \in \mathbb{N}\}$  is an i.i.d. process with  $U_n \sim \mathcal{U}(0,1)$ .

# Markov chains as recursions

theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,  
then the recursion  $X_{n+1} = g(X_n, U_n)$  is a Markov chain.

theorem 2:

Every Markov chain can be represented as recursion  $X_{n+1} = g(X_n, U_n)$   
where  $\{U_n : n \in \mathbb{N}\}$  is an i.i.d. process with  $U_n \sim \mathcal{U}(0,1)$ .

theorem 3:

# Markov chains as recursions

theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,  
then the recursion  $X_{n+1} = g(X_n, U_n)$  is a Markov chain.

theorem 2:

Every Markov chain can be represented as recursion  $X_{n+1} = g(X_n, U_n)$   
where  $\{U_n : n \in \mathbb{N}\}$  is an i.i.d. process with  $U_n \sim \mathcal{U}(0,1)$ .

theorem 3:

Given a Markov chain

# Markov chains as recursions

## theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,  
then the recursion  $X_{n+1} = g(X_n, U_n)$  is a Markov chain.

## theorem 2:

Every Markov chain can be represented as recursion  $X_{n+1} = g(X_n, U_n)$   
where  $\{U_n : n \in \mathbb{N}\}$  is an i.i.d. process with  $U_n \sim \mathcal{U}(0,1)$ .

## theorem 3:

Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$

# Markov chains as recursions

## theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,  
then the recursion  $X_{n+1} = g(X_n, U_n)$  is a Markov chain.

## theorem 2:

Every Markov chain can be represented as recursion  $X_{n+1} = g(X_n, U_n)$   
where  $\{U_n : n \in \mathbb{N}\}$  is an i.i.d. process with  $U_n \sim \mathcal{U}(0,1)$ .

## theorem 3:

Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  and its one-step transition matrix

# Markov chains as recursions

## theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,  
then the recursion  $X_{n+1} = g(X_n, U_n)$  is a Markov chain.

## theorem 2:

Every Markov chain can be represented as recursion  $X_{n+1} = g(X_n, U_n)$   
where  $\{U_n : n \in \mathbb{N}\}$  is an i.i.d. process with  $U_n \sim \mathcal{U}(0,1)$ .

## theorem 3:

Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  and its one-step transition matrix  $\underline{\underline{P}}$ ,

# Markov chains as recursions

## theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,  
then the recursion  $X_{n+1} = g(X_n, U_n)$  is a Markov chain.

## theorem 2:

Every Markov chain can be represented as recursion  $X_{n+1} = g(X_n, U_n)$   
where  $\{U_n : n \in \mathbb{N}\}$  is an i.i.d. process with  $U_n \sim \mathcal{U}(0,1)$ .

## theorem 3:

Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  and its one-step transition matrix  $\underline{\underline{P}}$ ,  
then its  $n$ -step transition matrix

# Markov chains as recursions

## theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,  
then the recursion  $X_{n+1} = g(X_n, U_n)$  is a Markov chain.

## theorem 2:

Every Markov chain can be represented as recursion  $X_{n+1} = g(X_n, U_n)$   
where  $\{U_n : n \in \mathbb{N}\}$  is an i.i.d. process with  $U_n \sim \mathcal{U}(0,1)$ .

## theorem 3:

Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  and its one-step transition matrix  $\underline{\underline{P}}$ ,  
then its  $n$ -step transition matrix  $\underline{\underline{P}}^{(n)}$



# Markov chains as recursions

## theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,  
then the recursion  $X_{n+1} = g(X_n, U_n)$  is a Markov chain.

## theorem 2:

Every Markov chain can be represented as recursion  $X_{n+1} = g(X_n, U_n)$   
where  $\{U_n : n \in \mathbb{N}\}$  is an i.i.d. process with  $U_n \sim \mathcal{U}(0,1)$ .

## theorem 3:

Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  and its one-step transition matrix  $\underline{\underline{P}}$ ,  
then its  $n$ -step transition matrix  $\underline{\underline{P}}^{(n)} = \underline{\underline{P}}^n$

# Markov chains as recursions

## theorem 1:

Let  $g(x, u) \in \mathbb{R}$  be a function and  $\{U_n : n \in \mathbb{N}\}$  an i.i.d. process,  
then the recursion  $X_{n+1} = g(X_n, U_n)$  is a Markov chain.

## theorem 2:

Every Markov chain can be represented as recursion  $X_{n+1} = g(X_n, U_n)$   
where  $\{U_n : n \in \mathbb{N}\}$  is an i.i.d. process with  $U_n \sim \mathcal{U}(0,1)$ .

## theorem 3:

Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  and its one-step transition matrix  $\underline{\underline{P}}$ ,  
then its  $n$ -step transition matrix  $\underline{\underline{P}}^{(n)} = \underline{\underline{P}}^n = \underline{\underline{P}} \cdot \dots \cdot \underline{\underline{P}}$ .

# properties of Markov chains

# properties of Markov chains

def.:

# properties of Markov chains

def.: Given a Markov chain

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .



# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Let

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Let  $\mathbb{C} \subseteq \mathbb{S}$ ,

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Let  $C \subseteq \mathbb{S}$ , then

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Let  $\mathbb{C} \subseteq \mathbb{S}$ , then  $\mathbb{C}$

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Let  $\mathbb{C} \subseteq \mathbb{S}$ , then  $\mathbb{C}$  is **closed** iff

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Let  $\mathbb{C} \subseteq \mathbb{S}$ , then  $\mathbb{C}$  is **closed** iff  $\forall x_i \in \mathbb{C} \forall x_j \in \mathbb{S} \setminus \mathbb{C} : P_{ij} = 0$ .

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Let  $\mathbb{C} \subseteq \mathbb{S}$ , then  $\mathbb{C}$  is **closed** iff  $\forall x_i \in \mathbb{C} \forall x_j \in \mathbb{S} \setminus \mathbb{C} : P_{ij} = 0$ .

(from a state of  $\mathbb{C}$  no state outside of  $\mathbb{C}$  can be reached)

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Let  $\mathbb{C} \subseteq \mathbb{S}$ , then  $\mathbb{C}$  is **closed** iff  $\forall x_i \in \mathbb{C} \forall x_j \in \mathbb{S} \setminus \mathbb{C} : P_{ij} = 0$ .

(from a state of  $\mathbb{C}$  no state outside of  $\mathbb{C}$  can be reached)

If



# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Let  $\mathbb{C} \subseteq \mathbb{S}$ , then  $\mathbb{C}$  is **closed** iff  $\forall x_i \in \mathbb{C} \forall x_j \in \mathbb{S} \setminus \mathbb{C} : P_{ij} = 0$ .

(from a state of  $\mathbb{C}$  no state outside of  $\mathbb{C}$  can be reached)

If  $\{x_i\} = \mathbb{C}$ ,

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Let  $\mathbb{C} \subseteq \mathbb{S}$ , then  $\mathbb{C}$  is **closed** iff  $\forall x_i \in \mathbb{C} \forall x_j \in \mathbb{S} \setminus \mathbb{C} : P_{ij} = 0$ .

(from a state of  $\mathbb{C}$  no state outside of  $\mathbb{C}$  can be reached)

If  $\{x_i\} = \mathbb{C}$ , then

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Let  $\mathbb{C} \subseteq \mathbb{S}$ , then  $\mathbb{C}$  is **closed** iff  $\forall x_i \in \mathbb{C} \forall x_j \in \mathbb{S} \setminus \mathbb{C} : P_{ij} = 0$ .

(from a state of  $\mathbb{C}$  no state outside of  $\mathbb{C}$  can be reached)

If  $\{x_i\} = \mathbb{C}$ , then  $x_i$

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Let  $\mathbb{C} \subseteq \mathbb{S}$ , then  $\mathbb{C}$  is **closed** iff  $\forall x_i \in \mathbb{C} \forall x_j \in \mathbb{S} \setminus \mathbb{C} : P_{ij} = 0$ .

(from a state of  $\mathbb{C}$  no state outside of  $\mathbb{C}$  can be reached)

If  $\{x_i\} = \mathbb{C}$ , then  $x_i$  is **absorbing state**.

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Let  $\mathbb{C} \subseteq \mathbb{S}$ , then  $\mathbb{C}$  is **closed** iff  $\forall x_i \in \mathbb{C} \forall x_j \in \mathbb{S} \setminus \mathbb{C} : P_{ij} = 0$ .

(from a state of  $\mathbb{C}$  no state outside of  $\mathbb{C}$  can be reached)

If  $\{x_i\} = \mathbb{C}$ , then  $x_i$  is **absorbing state**.  
 $\longrightarrow P_{ii} = 1$

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Let  $\mathbb{C} \subseteq \mathbb{S}$ , then  $\mathbb{C}$  is **closed** iff  $\forall x_i \in \mathbb{C} \forall x_j \in \mathbb{S} \setminus \mathbb{C} : P_{ij} = 0$ .

(from a state of  $\mathbb{C}$  no state outside of  $\mathbb{C}$  can be reached)

If  $\{x_i\} = \mathbb{C}$ , then  $x_i$  is **absorbing state**.  
 $\longrightarrow P_{ii} = 1$

exercise:

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Let  $\mathbb{C} \subseteq \mathbb{S}$ , then  $\mathbb{C}$  is **closed** iff  $\forall x_i \in \mathbb{C} \forall x_j \in \mathbb{S} \setminus \mathbb{C} : P_{ij} = 0$ .

(from a state of  $\mathbb{C}$  no state outside of  $\mathbb{C}$  can be reached)

If  $\{x_i\} = \mathbb{C}$ , then  $x_i$  is **absorbing state**.  
 $\longrightarrow P_{ii} = 1$

exercise:

consier:

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .


Let  $\mathbb{C} \subseteq \mathbb{S}$ , then  $\mathbb{C}$  is **closed** iff  $\forall x_i \in \mathbb{C} \forall x_j \in \mathbb{S} \setminus \mathbb{C} : P_{ij} = 0$ .

(from a state of  $\mathbb{C}$  no state outside of  $\mathbb{C}$  can be reached)

If  $\{x_i\} = \mathbb{C}$ , then  $x_i$  is **absorbing state**.  
 $\longrightarrow P_{ii} = 1$

exercise:

consier:





# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Let  $\mathbb{C} \subseteq \mathbb{S}$ , then  $\mathbb{C}$  is **closed** iff  $\forall x_i \in \mathbb{C} \forall x_j \in \mathbb{S} \setminus \mathbb{C} : P_{ij} = 0$ .


(from a state of  $\mathbb{C}$  no state outside of  $\mathbb{C}$  can be reached)

If  $\{x_i\} = \mathbb{C}$ , then  $x_i$  is **absorbing state**.  
 $\longrightarrow P_{ii} = 1$

exercise:

What are the closed sets?

consier:



# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .


Let  $\mathbb{C} \subseteq \mathbb{S}$ , then  $\mathbb{C}$  is **closed** iff  $\forall x_i \in \mathbb{C} \forall x_j \in \mathbb{S} \setminus \mathbb{C} : P_{ij} = 0$ .

(from a state of  $\mathbb{C}$  no state outside of  $\mathbb{C}$  can be reached)

If  $\{x_i\} = \mathbb{C}$ , then  $x_i$  is **absorbing state**.  
 $\longrightarrow P_{ii} = 1$

exercise:

consier:



What are the closed sets?

What are the absorbing states?

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .


Let  $\mathbb{C} \subseteq \mathbb{S}$ , then  $\mathbb{C}$  is **closed** iff  $\forall x_i \in \mathbb{C} \forall x_j \in \mathbb{S} \setminus \mathbb{C} : P_{ij} = 0$ .

(from a state of  $\mathbb{C}$  no state outside of  $\mathbb{C}$  can be reached)

If  $\{x_i\} = \mathbb{C}$ , then  $x_i$  is **absorbing state**.  
 $\longrightarrow P_{ii} = 1$

exercise:

consier:



What are the closed sets?  $\mathbb{C}' = \{x_0, x_1\}$ ,

What are the absorbing states?

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .


Let  $\mathbb{C} \subseteq \mathbb{S}$ , then  $\mathbb{C}$  is **closed** iff  $\forall x_i \in \mathbb{C} \forall x_j \in \mathbb{S} \setminus \mathbb{C} : P_{ij} = 0$ .

(from a state of  $\mathbb{C}$  no state outside of  $\mathbb{C}$  can be reached)

If  $\{x_i\} = \mathbb{C}$ , then  $x_i$  is **absorbing state**.  
 $\longrightarrow P_{ii} = 1$

exercise:

consier:



What are the closed sets?  $\mathbb{C}' = \{x_0, x_1\}$ ,  $\mathbb{C}'' = \{x_1\}$ ,

What are the absorbing states?

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .


Let  $\mathbb{C} \subseteq \mathbb{S}$ , then  $\mathbb{C}$  is **closed** iff  $\forall x_i \in \mathbb{C} \forall x_j \in \mathbb{S} \setminus \mathbb{C} : P_{ij} = 0$ .

(from a state of  $\mathbb{C}$  no state outside of  $\mathbb{C}$  can be reached)

If  $\{x_i\} = \mathbb{C}$ , then  $x_i$  is **absorbing state**.  
 $\longrightarrow P_{ii} = 1$

exercise:

consier:



What are the closed sets?  $\mathbb{C}' = \{x_0, x_1\}$ ,  $\mathbb{C}'' = \{x_1\}$ ,  $\mathbb{C}''' = \mathbb{S}$

What are the absorbing states?

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .


Let  $\mathbb{C} \subseteq \mathbb{S}$ , then  $\mathbb{C}$  is **closed** iff  $\forall x_i \in \mathbb{C} \forall x_j \in \mathbb{S} \setminus \mathbb{C} : P_{ij} = 0$ .

(from a state of  $\mathbb{C}$  no state outside of  $\mathbb{C}$  can be reached)

If  $\{x_i\} = \mathbb{C}$ , then  $x_i$  is **absorbing state**.  
 $\longrightarrow P_{ii} = 1$

exercise:

consier:



What are the closed sets?  $\mathbb{C}' = \{x_0, x_1\}$ ,  $\mathbb{C}'' = \{x_1\}$ ,  $\mathbb{C}''' = \mathbb{S}$

What are the absorbing states?  $x_1$

# properties of Markov chains

# properties of Markov chains

def.:



# properties of Markov chains

def.: A Markov chain

# properties of Markov chains

def.: A Markov chain  $\{X_n : n \in \mathbb{N}\}$

# properties of Markov chains

def.: A Markov chain  $\{X_n : n \in \mathbb{N}\}$  with

# properties of Markov chains

def.: A Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$

# properties of Markov chains

def.: A Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$   
is **irreducible** iff

# properties of Markov chains

def.: A Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$   
is **irreducible** iff  $\nexists \mathbb{C} \subset \mathbb{S} : \mathbb{C} \text{ is closed}$

# properties of Markov chains

def.: A Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$   
is **irreducible** iff  $\nexists \mathbb{C} \subset \mathbb{S} : \mathbb{C} \text{ is closed}$  (  $\mathbb{S}$  is the only closed set )

# properties of Markov chains

def.: A Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$   
is **irreducible** iff  $\nexists \mathbb{C} \subset \mathbb{S} : \mathbb{C} \text{ is closed}$  (  $\mathbb{S}$  is the only closed set )

exercise:



# properties of Markov chains


def.: A Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$   
is **irreducible** iff  $\nexists \mathbb{C} \subset \mathbb{S} : \mathbb{C} \text{ is closed}$  ( $\mathbb{S}$  is the only closed set)

**exercise:** Which Markov chain is irreducible?

# properties of Markov chains

def.: A Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$   
is **irreducible** iff  $\nexists \mathbb{C} \subset \mathbb{S} : \mathbb{C} \text{ is closed}$  ( $\mathbb{S}$  is the only closed set)



**exercise:** Which Markov chain is irreducible?



# properties of Markov chains

def.: A Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$   
is **irreducible** iff  $\nexists \mathbb{C} \subset \mathbb{S} : \mathbb{C} \text{ is closed}$  ( $\mathbb{S}$  is the only closed set)



**exercise:** Which Markov chain is irreducible?



# properties of Markov chains

def.: A Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$   
is **irreducible** iff  $\nexists \mathbb{C} \subset \mathbb{S} : \mathbb{C} \text{ is closed}$  ( $\mathbb{S}$  is the only closed set)



exercise: Which Markov chain is irreducible?



# properties of Markov chains

def.: A Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$   
is **irreducible** iff  $\nexists \mathbb{C} \subset \mathbb{S} : \mathbb{C} \text{ is closed}$  ( $\mathbb{S}$  is the only closed set)

exercise: Which Markov chain is irreducible?



# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$



# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,  
then

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ ,

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then



# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then  $x_i$

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if  $d_i = 1$ ,

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if  $d_i = 1$ , then

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if  $d_i = 1$ , then  $x_i$

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if  $d_i = 1$ , then  $x_i$  is **aperiodic**.

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if  $d_i = 1$ , then  $x_i$  is **aperiodic**.

If



# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if  $d_i = 1$ , then  $x_i$  is **aperiodic**.

If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if  $d_i = 1$ , then  $x_i$  is **aperiodic**.

If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,

then

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if  $d_i = 1$ , then  $x_i$  is **aperiodic**.

If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,

then  $\{X_n : n \in \mathbb{N}\}$

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if  $d_i = 1$ , then  $x_i$  is **aperiodic**.

If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,

then  $\{X_n : n \in \mathbb{N}\}$  is **periodic**,

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if  $d_i = 1$ , then  $x_i$  is **aperiodic**.

If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,

then  $\{X_n : n \in \mathbb{N}\}$  is **periodic**,

(The Markov chain is **periodic** if all states are periodic with the same period.)

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if  $d_i = 1$ , then  $x_i$  is **aperiodic**.

If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,

then  $\{X_n : n \in \mathbb{N}\}$  is **periodic**,

(The Markov chain is **periodic** if all states are periodic with the same period.)

if

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if  $d_i = 1$ , then  $x_i$  is **aperiodic**.

If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,

then  $\{X_n : n \in \mathbb{N}\}$  is **periodic**,

(The Markov chain is **periodic** if all states are periodic with the same period.)

if  $\forall x_i \in \mathbb{S} : 1 = d_i$ ,

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if  $d_i = 1$ , then  $x_i$  is **aperiodic**.

If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,

then  $\{X_n : n \in \mathbb{N}\}$  is **periodic**,

(The Markov chain is **periodic** if all states are periodic with the same period.)

if  $\forall x_i \in \mathbb{S} : 1 = d_i$ , then



# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if  $d_i = 1$ , then  $x_i$  is **aperiodic**.

If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,

then  $\{X_n : n \in \mathbb{N}\}$  is **periodic**,

(The Markov chain is **periodic** if all states are periodic with the same period.)

if  $\forall x_i \in \mathbb{S} : 1 = d_i$ , then  $\{X_n : n \in \mathbb{N}\}$

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if  $d_i = 1$ , then  $x_i$  is **aperiodic**.

If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,

then  $\{X_n : n \in \mathbb{N}\}$  is **periodic**,

(The Markov chain is **periodic** if all states are periodic with the same period.)

if  $\forall x_i \in \mathbb{S} : 1 = d_i$ , then  $\{X_n : n \in \mathbb{N}\}$  is **aperiodic**.

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if  $d_i = 1$ , then  $x_i$  is **aperiodic**.

If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,

then  $\{X_n : n \in \mathbb{N}\}$  is **periodic**,

(The Markov chain is **periodic** if all states are periodic with the same period.)

if  $\forall x_i \in \mathbb{S} : 1 = d_i$ , then  $\{X_n : n \in \mathbb{N}\}$  is **aperiodic**.

(The Markov chain is **aperiodic** if all states are aperiodic.)

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

**exercise:**

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if  $d_i = 1$ , then  $x_i$  is **aperiodic**.

If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,

then  $\{X_n : n \in \mathbb{N}\}$  is **periodic**,

(The Markov chain is **periodic** if all states are periodic with the same period.)

if  $\forall x_i \in \mathbb{S} : 1 = d_i$ , then  $\{X_n : n \in \mathbb{N}\}$  is **aperiodic**.

(The Markov chain is **aperiodic** if all states are aperiodic.)

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

**exercise:**

Which states/ Markov chains are periodic?

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if  $d_i = 1$ , then  $x_i$  is **aperiodic**.

If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,

then  $\{X_n : n \in \mathbb{N}\}$  is **periodic**,

(The Markov chain is **periodic** if all states are periodic with the same period.)

if  $\forall x_i \in \mathbb{S} : 1 = d_i$ , then  $\{X_n : n \in \mathbb{N}\}$  is **aperiodic**.

(The Markov chain is **aperiodic** if all states are aperiodic.)

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

**exercise:**

Which states/ Markov chains are periodic?

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,

if  $d_i = 1$ , then  $x_i$  is **aperiodic**.


If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,

then  $\{X_n : n \in \mathbb{N}\}$  is **periodic**,

(The Markov chain is **periodic** if all states are periodic with the same period.)

if  $\forall x_i \in \mathbb{S} : 1 = d_i$ , then  $\{X_n : n \in \mathbb{N}\}$  is **aperiodic**.

(The Markov chain is **aperiodic** if all states are aperiodic.)





# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

**exercise:**

Which states/ Markov chains are periodic?



If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,  
if  $d_i = 1$ , then  $x_i$  is **aperiodic**.

If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,  
then  $\{X_n : n \in \mathbb{N}\}$  is **periodic**,

(The Markov chain is **periodic** if all states are periodic with the same period.)

if  $\forall x_i \in \mathbb{S} : 1 = d_i$ , then  $\{X_n : n \in \mathbb{N}\}$  is **aperiodic**.

(The Markov chain is **aperiodic** if all states are aperiodic.)



# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

**exercise:**




Which states/ Markov chains are periodic?

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,  
if  $d_i = 1$ , then  $x_i$  is **aperiodic**.

If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,  
then  $\{X_n : n \in \mathbb{N}\}$  is **periodic**,

(The Markov chain is **periodic** if all states are periodic with the same period.)

if  $\forall x_i \in \mathbb{S} : 1 = d_i$ , then  $\{X_n : n \in \mathbb{N}\}$  is **aperiodic**.  
(The Markov chain is **aperiodic** if all states are aperiodic.)





# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

**exercise:**


Which states/ Markov chains are periodic?

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,  
if  $d_i = 1$ , then  $x_i$  is **aperiodic**.



If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,  
then  $\{X_n : n \in \mathbb{N}\}$  is **periodic**,

(The Markov chain is **periodic** if all states are periodic with the same period.)

if  $\forall x_i \in \mathbb{S} : 1 = d_i$ , then  $\{X_n : n \in \mathbb{N}\}$  is **aperiodic**.  
(The Markov chain is **aperiodic** if all states are aperiodic.)



$x_0, x_2$  are periodic



# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

**exercise:**


Which states/ Markov chains are periodic?

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,  
if  $d_i = 1$ , then  $x_i$  is **aperiodic**.



If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,  
then  $\{X_n : n \in \mathbb{N}\}$  is **periodic**,

(The Markov chain is **periodic** if all states are periodic with the same period.)

if  $\forall x_i \in \mathbb{S} : 1 = d_i$ , then  $\{X_n : n \in \mathbb{N}\}$  is **aperiodic**.  
(The Markov chain is **aperiodic** if all states are aperiodic.)



$x_0, x_2$  are periodic  
 $d_i = 2$



# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

**exercise:**


Which states/ Markov chains are periodic?

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,  
if  $d_i = 1$ , then  $x_i$  is **aperiodic**.



If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,  
then  $\{X_n : n \in \mathbb{N}\}$  is **periodic**,

(The Markov chain is **periodic** if all states are periodic with the same period.)

if  $\forall x_i \in \mathbb{S} : 1 = d_i$ , then  $\{X_n : n \in \mathbb{N}\}$  is **aperiodic**.  
(The Markov chain is **aperiodic** if all states are aperiodic.)



$x_0, x_2$  are periodic  
 $d_i = 2$   
 $x_1$  is aperiodic



# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

**exercise:**


Which states/ Markov chains are periodic?

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,  
if  $d_i = 1$ , then  $x_i$  is **aperiodic**.

If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,  
then  $\{X_n : n \in \mathbb{N}\}$  is **periodic**,

(The Markov chain is **periodic** if all states are periodic with the same period.)


if  $\forall x_i \in \mathbb{S} : 1 = d_i$ , then  $\{X_n : n \in \mathbb{N}\}$  is **aperiodic**.  
(The Markov chain is **aperiodic** if all states are aperiodic.)




$x_0, x_2$  are periodic

$d_i = 2$

$x_1$  is aperiodic



all  $x_i$  are aperiodic





# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

**exercise:**


Which states/ Markov chains are periodic?

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,  
if  $d_i = 1$ , then  $x_i$  is **aperiodic**.

If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,  
then  $\{X_n : n \in \mathbb{N}\}$  is **periodic**,

(The Markov chain is **periodic** if all states are periodic with the same period.)


if  $\forall x_i \in \mathbb{S} : 1 = d_i$ , then  $\{X_n : n \in \mathbb{N}\}$  is **aperiodic**.  
(The Markov chain is **aperiodic** if all states are aperiodic.)




$x_0, x_2$  are periodic

$d_i = 2$

$x_1$  is aperiodic



all  $x_i$  are aperiodic



all  $x_i$  are periodic

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $N_i := \{n \in \mathbb{N}_{>0} : P_{ii}^{(n)} > 0\}$  set of number of transitions to reach  $x_i$  from  $x_i$ ,

then  $d_i := \gcd(N_i)$  **period** of  $x_i$ .

(greatest common divisor)

**exercise:**


Which states/ Markov chains are periodic?

If  $1 < d_i < \infty$ , then  $x_i$  is **periodic**,  
if  $d_i = 1$ , then  $x_i$  is **aperiodic**.


If  $\forall x_i, x_j \in \mathbb{S} : 1 < d_i = d_j < \infty$ ,  
then  $\{X_n : n \in \mathbb{N}\}$  is **periodic**,

(The Markov chain is **periodic** if all states are periodic with the same period.)


if  $\forall x_i \in \mathbb{S} : 1 = d_i$ , then  $\{X_n : n \in \mathbb{N}\}$  is **aperiodic**.  
(The Markov chain is **aperiodic** if all states are aperiodic.)



$x_0, x_2$  are periodic  
 $d_i = 2$   
 $x_1$  is aperiodic



all  $x_i$  are aperiodic



all  $x_i$  are periodic  
 $d_i = 4$

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

The Markov chain

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

If

The Markov chain



# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

$$\text{If } \sum_{n=1}^{\infty} P_{ii}^{(n)} < \infty$$

The Markov chain

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

If  $\sum_{n=1}^{\infty} P_{ii}^{(n)} < \infty$  then

The Markov chain

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

If  $\sum_{n=1}^{\infty} P_{ii}^{(n)} < \infty$  then  $x_i$

The Markov chain

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

If  $\sum_{n=1}^{\infty} P_{ii}^{(n)} < \infty$  then  $x_i$  is **transient**,

The Markov chain

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

If  $\sum_{n=1}^{\infty} P_{ii}^{(n)} < \infty$  then  $x_i$  is **transient**,

if

The Markov chain

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

If  $\sum_{n=1}^{\infty} P_{ii}^{(n)} < \infty$  then  $x_i$  is **transient**,

if  $\sum_{n=1}^{\infty} P_{ii}^{(n)} = \infty$

The Markov chain

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

If  $\sum_{n=1}^{\infty} P_{ii}^{(n)} < \infty$  then  $x_i$  is **transient**,

if  $\sum_{n=1}^{\infty} P_{ii}^{(n)} = \infty$  then

The Markov chain

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

If  $\sum_{n=1}^{\infty} P_{ii}^{(n)} < \infty$  then  $x_i$  is **transient**,

if  $\sum_{n=1}^{\infty} P_{ii}^{(n)} = \infty$  then  $x_i$

The Markov chain



# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

If  $\sum_{n=1}^{\infty} P_{ii}^{(n)} < \infty$  then  $x_i$  is **transient**,

if  $\sum_{n=1}^{\infty} P_{ii}^{(n)} = \infty$  then  $x_i$  is **recurrent**.

The Markov chain

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

If  $\sum_{n=1}^{\infty} P_{ii}^{(n)} < \infty$  then  $x_i$  is **transient**, (A state is **transient** iff there is a non-zero probability that it is never reached again and **recurrent** otherwise.)

if  $\sum_{n=1}^{\infty} P_{ii}^{(n)} = \infty$  then  $x_i$  is **recurrent**.

The Markov chain

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

If  $\sum_{n=1}^{\infty} P_{ii}^{(n)} < \infty$  then  $x_i$  is **transient**, (A state is **transient** iff there is a non-zero probability that it is never reached again and **recurrent** otherwise.)

if  $\sum_{n=1}^{\infty} P_{ii}^{(n)} = \infty$  then  $x_i$  is **recurrent**.

The Markov chain is **ergodic**,

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

If  $\sum_{n=1}^{\infty} P_{ii}^{(n)} < \infty$  then  $x_i$  is **transient**, (A state is **transient** iff there is a non-zero probability that it is never reached again and **recurrent** otherwise.)

if  $\sum_{n=1}^{\infty} P_{ii}^{(n)} = \infty$  then  $x_i$  is **recurrent**.

The Markov chain  $\{X_n : n \in \mathbb{N}\}$  is **ergodic**,

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

If  $\sum_{n=1}^{\infty} P_{ii}^{(n)} < \infty$  then  $x_i$  is **transient**, (A state is **transient** iff there is a non-zero probability that it is never reached again and **recurrent** otherwise.)

if  $\sum_{n=1}^{\infty} P_{ii}^{(n)} = \infty$  then  $x_i$  is **recurrent**.

The Markov chain  $\{X_n : n \in \mathbb{N}\}$  is **ergodic**, iff it is irreducible and aperiodic.

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

limiting distribution:

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

limiting distribution:

$$P_k := \lim_{n \rightarrow \infty} P_{jk}^{(n)}$$



# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

limiting distribution:  $P_k := \lim_{n \rightarrow \infty} P_{jk}^{(n)} \quad \forall x_j \in \mathbb{S}$

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

limiting distribution:

$$P_k := \lim_{n \rightarrow \infty} P_{jk}^{(n)} \quad \forall x_j \in \mathbb{S}$$

$$\sum_{x_i \in \mathbb{S}} P_i = 1$$

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

limiting distribution:

$$P_k := \lim_{n \rightarrow \infty} P_{jk}^{(n)} \quad \forall x_j \in \mathbb{S}$$

$$\sum_{x_i \in \mathbb{S}} P_i = 1$$

theorem 4:

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

limiting distribution:

$$P_k := \lim_{n \rightarrow \infty} P_{jk}^{(n)} \quad \forall x_j \in \mathbb{S}$$

$$\sum_{x_i \in \mathbb{S}} P_i = 1$$

theorem 4:

A Markov chain has a limiting distribution iff it is ergodic.

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

limiting distribution:  $P_k := \lim_{n \rightarrow \infty} P_{jk}^{(n)} \quad \forall x_j \in \mathbb{S}$

$$\sum_{x_i \in \mathbb{S}} P_i = 1$$

theorem 4:

A Markov chain has a limiting distribution iff it is ergodic.

example:

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

limiting distribution:  $P_k := \lim_{n \rightarrow \infty} P_{jk}^{(n)} \quad \forall x_j \in \mathbb{S}$

$$\sum_{x_i \in \mathbb{S}} P_i = 1$$

theorem 4:

A Markov chain has a limiting distribution iff it is ergodic.

example:

Every i.i.d. process is a Markov chain with

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

limiting distribution:  $P_k := \lim_{n \rightarrow \infty} P_{jk}^{(n)} \quad \forall x_j \in \mathbb{S}$

$$\sum_{x_i \in \mathbb{S}} P_i = 1$$

theorem 4:

A Markov chain has a limiting distribution iff it is ergodic.

example:

Every i.i.d. process is a Markov chain with  $g(x, U) := U$

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

limiting distribution:  $P_k := \lim_{n \rightarrow \infty} P_{jk}^{(n)} \quad \forall x_j \in \mathbb{S}$

$$\sum_{x_i \in \mathbb{S}} P_i = 1$$

theorem 4:

A Markov chain has a limiting distribution iff it is ergodic.

example:

Every i.i.d. process is a Markov chain with  $g(x, U) := U \quad U \sim \mathcal{U}(0,1)$



# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

limiting distribution:  $P_k := \lim_{n \rightarrow \infty} P_{jk}^{(n)} \quad \forall x_j \in \mathbb{S}$

$$\sum_{x_i \in \mathbb{S}} P_i = 1$$

theorem 4:

A Markov chain has a limiting distribution iff it is ergodic.

example:

Every i.i.d. process is a Markov chain with  $g(x, U) := U \quad U \sim \mathcal{U}(0,1)$

and the limiting distribution is

# properties of Markov chains

def.: Given a Markov chain  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

limiting distribution:  $P_k := \lim_{n \rightarrow \infty} P_{jk}^{(n)} \quad \forall x_j \in \mathbb{S}$

$$\sum_{x_i \in \mathbb{S}} P_i = 1$$

theorem 4:

A Markov chain has a limiting distribution iff it is ergodic.

example:

Every i.i.d. process is a Markov chain with  $g(x, U) := U \quad U \sim \mathcal{U}(0,1)$

and the limiting distribution is  $\mathcal{U}(0,1)$ .

# Markov chains for random number generation

Markov chain

$$\{X_n : n \in \mathbb{N}\} \quad X_n : \Omega \rightarrow \mathbb{S}$$

limiting distribution

$$P_k := \lim_{n \rightarrow \infty} P_{jk}^{(n)} \quad \forall x_j \in \mathbb{S}$$

# Markov chains for random number generation

Markov chain

$$\{X_n : n \in \mathbb{N}\} \quad X_n : \Omega \rightarrow \mathbb{S}$$

limiting distribution

$$P_k := \lim_{n \rightarrow \infty} P_{jk}^{(n)} \quad \forall x_j \in \mathbb{S}$$

praxis:

# Markov chains for random number generation

Markov chain

$$\{X_n : n \in \mathbb{N}\} \quad X_n : \Omega \rightarrow \mathbb{S}$$

limiting distribution

$$P_k := \lim_{n \rightarrow \infty} P_{jk}^{(n)} \quad \forall x_j \in \mathbb{S}$$

praxis:  $n \rightarrow \infty$

# Markov chains for random number generation

Markov chain

$$\{X_n : n \in \mathbb{N}\} \quad X_n : \Omega \rightarrow \mathbb{S}$$

limiting distribution

$$P_k := \lim_{n \rightarrow \infty} P_{jk}^{(n)} \quad \forall x_j \in \mathbb{S}$$

praxis:

~~$n \rightarrow \infty$~~

# Markov chains for random number generation

Markov chain

$$\{X_n : n \in \mathbb{N}\} \quad X_n : \Omega \rightarrow \mathbb{S}$$

limiting distribution

$$P_k := \lim_{n \rightarrow \infty} P_{jk}^{(n)} \quad \forall x_j \in \mathbb{S}$$

praxis:  ~~$n \rightarrow \infty$~~   $n \gg 1$

# Markov chains for random number generation

Markov chain

$$\{X_n : n \in \mathbb{N}\} \quad X_n : \Omega \rightarrow \mathbb{S}$$

limiting distribution

$$P_k := \lim_{n \rightarrow \infty} P_{jk}^{(n)} \quad \forall x_j \in \mathbb{S}$$

praxis:

~~$n \rightarrow \infty$~~

$n \gg 1$



# Markov chains for random number generation

Markov chain

$$\{X_n : n \in \mathbb{N}\} \quad X_n : \Omega \rightarrow \mathbb{S}$$

limiting distribution

$$P_k := \lim_{n \rightarrow \infty} P_{jk}^{(n)} \quad \forall x_j \in \mathbb{S}$$

praxis:

~~$n \rightarrow \infty$~~

$n \gg 1$

→ "mixing time"

# Markov chains for random number generation

Markov chain

$$\{X_n : n \in \mathbb{N}\} \quad X_n : \Omega \rightarrow \mathbb{S}$$

limiting distribution

$$P_k := \lim_{n \rightarrow \infty} P_{jk}^{(n)} \quad \forall x_j \in \mathbb{S}$$

praxis:

~~$n \rightarrow \infty$~~

$n \gg 1$

→ "mixing time"

(In praxis, statistical tests decide if the limiting distribution is reached.)

end