

# stochastics and probability

## Lecture 4

Dr. Johannes Pahlke

# moments

def.:

# moments

def.: Be

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$

# moments

def.: Be  $X: \Omega \rightarrow \mathbb{S}$  a discrete random variable and

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$

# moments

def.: Be  $X: \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i)$$



# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value}$$

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value}$$

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7),$$

# moments

def.: Be  $X: \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7},$$

# moments

def.: Be  $X: \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:  $x \in \mathbb{S} := (1, 2, \dots, 7)$ ,  $p_X(x) := \frac{1}{7}$ ,  $g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$

# moments

def.: Be  $X: \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\mathbb{E}[X] =$$



# moments

def.: Be  $X: \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\mathbb{E}[X] =$$

$$\mathbb{E}[g(X)] =$$

# moments

def.: Be  $X: \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\mathbb{E}[X] = 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7)$$

$$\mathbb{E}[g(X)] =$$

# moments

def.: Be  $X: \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} \end{aligned}$$

$$\mathbb{E}[g(X)] =$$

# moments

def.: Be  $X: \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} \end{aligned}$$

$$\mathbb{E}[g(X)] =$$

# moments

def.: Be  $X: \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\mathbb{E}[X] = 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7)$$

$$= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4$$

$$\mathbb{E}[g(X)] =$$

# moments

def.: Be  $X: \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\mathbb{E}[X] = 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7)$$

$$= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4$$

$$\mathbb{E}[g(X)] = g(1) \cdot p_X(1) + \dots + g(7) \cdot p_X(7)$$

# moments

def.: Be  $X: \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4 \end{aligned}$$

$$\begin{aligned} \mathbb{E}[g(X)] &= g(1) \cdot p_X(1) + \dots + g(7) \cdot p_X(7) \\ &= (0 + 0 + 0 + 0 + 0 + 1 + 1) \cdot \frac{1}{7} \end{aligned}$$

# moments

def.: Be  $X: \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4 \end{aligned}$$

$$\begin{aligned} \mathbb{E}[g(X)] &= g(1) \cdot p_X(1) + \dots + g(7) \cdot p_X(7) \\ &= (0 + 0 + 0 + 0 + 0 + 1 + 1) \cdot \frac{1}{7} = \frac{2}{7} \end{aligned}$$



# moments

def.: Be  $X: \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4 \end{aligned}$$

$$\begin{aligned} \mathbb{E}[g(X)] &= g(1) \cdot p_X(1) + \dots + g(7) \cdot p_X(7) \\ &= (0 + 0 + 0 + 0 + 0 + 1 + 1) \cdot \frac{1}{7} = \frac{2}{7} \end{aligned}$$

Be

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4 \end{aligned}$$

$$\begin{aligned} \mathbb{E}[g(X)] &= g(1) \cdot p_X(1) + \dots + g(7) \cdot p_X(7) \\ &= (0 + 0 + 0 + 0 + 0 + 1 + 1) \cdot \frac{1}{7} = \frac{2}{7} \end{aligned}$$

Be  $X : \Omega \rightarrow \mathbb{S}$

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4 \end{aligned}$$

$$\begin{aligned} \mathbb{E}[g(X)] &= g(1) \cdot p_X(1) + \dots + g(7) \cdot p_X(7) \\ &= (0 + 0 + 0 + 0 + 0 + 1 + 1) \cdot \frac{1}{7} = \frac{2}{7} \end{aligned}$$

Be  $X : \Omega \rightarrow \mathbb{S}$  a continuous random variable and

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4 \end{aligned}$$

$$\begin{aligned} \mathbb{E}[g(X)] &= g(1) \cdot p_X(1) + \dots + g(7) \cdot p_X(7) \\ &= (0 + 0 + 0 + 0 + 0 + 1 + 1) \cdot \frac{1}{7} = \frac{2}{7} \end{aligned}$$

Be  $X : \Omega \rightarrow \mathbb{S}$  a continuous random variable and  $f_X(x)$

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4 \end{aligned}$$

$$\begin{aligned} \mathbb{E}[g(X)] &= g(1) \cdot p_X(1) + \dots + g(7) \cdot p_X(7) \\ &= (0 + 0 + 0 + 0 + 0 + 1 + 1) \cdot \frac{1}{7} = \frac{2}{7} \end{aligned}$$

Be  $X : \Omega \rightarrow \mathbb{S}$  a continuous random variable and  $f_X(x)$  its probability density function (PDF).

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4 \end{aligned}$$

$$\begin{aligned} \mathbb{E}[g(X)] &= g(1) \cdot p_X(1) + \dots + g(7) \cdot p_X(7) \\ &= (0 + 0 + 0 + 0 + 0 + 1 + 1) \cdot \frac{1}{7} = \frac{2}{7} \end{aligned}$$

Be  $X : \Omega \rightarrow \mathbb{S}$  a continuous random variable and  $f_X(x)$  its probability density function (PDF).

$$\mathbb{E}[X] := \int_{\mathbb{S}} x f_X(x) dx$$

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4 \end{aligned}$$

$$\begin{aligned} \mathbb{E}[g(X)] &= g(1) \cdot p_X(1) + \dots + g(7) \cdot p_X(7) \\ &= (0 + 0 + 0 + 0 + 0 + 1 + 1) \cdot \frac{1}{7} = \frac{2}{7} \end{aligned}$$

Be  $X : \Omega \rightarrow \mathbb{S}$  a continuous random variable and  $f_X(x)$  its probability density function (PDF).

$$\mathbb{E}[X] := \int_{\mathbb{S}} x f_X(x) dx \quad \text{expectation/ expected value}$$

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4 \end{aligned}$$

$$\begin{aligned} \mathbb{E}[g(X)] &= g(1) \cdot p_X(1) + \dots + g(7) \cdot p_X(7) \\ &= (0 + 0 + 0 + 0 + 0 + 1 + 1) \cdot \frac{1}{7} = \frac{2}{7} \end{aligned}$$

Be  $X : \Omega \rightarrow \mathbb{S}$  a continuous random variable and  $f_X(x)$  its probability density function (PDF).

$$\mu := \mathbb{E}[X] := \int_{\mathbb{S}} x f_X(x) dx \quad \text{expectation/ expected value}$$



# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4 \end{aligned}$$

$$\begin{aligned} \mathbb{E}[g(X)] &= g(1) \cdot p_X(1) + \dots + g(7) \cdot p_X(7) \\ &= (0 + 0 + 0 + 0 + 0 + 1 + 1) \cdot \frac{1}{7} = \frac{2}{7} \end{aligned}$$

Be  $X : \Omega \rightarrow \mathbb{S}$  a continuous random variable and  $f_X(x)$  its probability density function (PDF).

$$\mu := \mathbb{E}[X] := \int_{\mathbb{S}} x f_X(x) dx \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx \quad \right]$$

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4 \end{aligned}$$

$$\begin{aligned} \mathbb{E}[g(X)] &= g(1) \cdot p_X(1) + \dots + g(7) \cdot p_X(7) \\ &= (0 + 0 + 0 + 0 + 0 + 1 + 1) \cdot \frac{1}{7} = \frac{2}{7} \end{aligned}$$

Be  $X : \Omega \rightarrow \mathbb{S}$  a continuous random variable and  $f_X(x)$  its probability density function (PDF).

$$\mu := \mathbb{E}[X] := \int_{\mathbb{S}} x f_X(x) dx \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx \quad \right]$$

$$\mu_k := \mathbb{E}[(X - \mu)^k]$$

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4 \end{aligned}$$

$$\begin{aligned} \mathbb{E}[g(X)] &= g(1) \cdot p_X(1) + \dots + g(7) \cdot p_X(7) \\ &= (0 + 0 + 0 + 0 + 0 + 1 + 1) \cdot \frac{1}{7} = \frac{2}{7} \end{aligned}$$

Be  $X : \Omega \rightarrow \mathbb{S}$  a continuous random variable and  $f_X(x)$  its probability density function (PDF).

$$\mu := \mathbb{E}[X] := \int_{\mathbb{S}} x f_X(x) dx \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx \quad \right]$$

$$\mu_k := \mathbb{E}[(X - \mu)^k] \quad \text{higher-order central moments}$$

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4 \end{aligned}$$

$$\begin{aligned} \mathbb{E}[g(X)] &= g(1) \cdot p_X(1) + \dots + g(7) \cdot p_X(7) \\ &= (0 + 0 + 0 + 0 + 0 + 1 + 1) \cdot \frac{1}{7} = \frac{2}{7} \end{aligned}$$

Be  $X : \Omega \rightarrow \mathbb{S}$  a continuous random variable and  $f_X(x)$  its probability density function (PDF).

$$\mu := \mathbb{E}[X] := \int_{\mathbb{S}} x f_X(x) dx \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx \quad \right]$$

$$\mu_k := \mathbb{E}[(X - \mu)^k] \quad \text{higher-order central moments}$$

$$\text{Var}(X) := \mu_2$$

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4 \end{aligned}$$

$$\begin{aligned} \mathbb{E}[g(X)] &= g(1) \cdot p_X(1) + \dots + g(7) \cdot p_X(7) \\ &= (0 + 0 + 0 + 0 + 0 + 1 + 1) \cdot \frac{1}{7} = \frac{2}{7} \end{aligned}$$

Be  $X : \Omega \rightarrow \mathbb{S}$  a continuous random variable and  $f_X(x)$  its probability density function (PDF).

$$\mu := \mathbb{E}[X] := \int_{\mathbb{S}} x f_X(x) dx \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx \quad \right]$$

$$\mu_k := \mathbb{E}[(X - \mu)^k] \quad \text{higher-order central moments}$$

$$\text{Var}(X) := \mu_2 \quad \text{variance (2nd central moments)}$$

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4 \end{aligned}$$

$$\begin{aligned} \mathbb{E}[g(X)] &= g(1) \cdot p_X(1) + \dots + g(7) \cdot p_X(7) \\ &= (0 + 0 + 0 + 0 + 0 + 1 + 1) \cdot \frac{1}{7} = \frac{2}{7} \end{aligned}$$

Be  $X : \Omega \rightarrow \mathbb{S}$  a continuous random variable and  $f_X(x)$  its probability density function (PDF).

$$\mu := \mathbb{E}[X] := \int_{\mathbb{S}} x f_X(x) dx \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx \quad \right]$$

$$\mu_k := \mathbb{E}[(X - \mu)^k] \quad \text{higher-order central moments}$$

$$\text{Var}(X) := \mu_2 = \mathbb{E}[(X - \mu)^2] \quad \text{variance (2nd central moments)}$$



# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4 \end{aligned}$$

$$\begin{aligned} \mathbb{E}[g(X)] &= g(1) \cdot p_X(1) + \dots + g(7) \cdot p_X(7) \\ &= (0 + 0 + 0 + 0 + 0 + 1 + 1) \cdot \frac{1}{7} = \frac{2}{7} \end{aligned}$$

Be  $X : \Omega \rightarrow \mathbb{S}$  a continuous random variable and  $f_X(x)$  its probability density function (PDF).

$$\mu := \mathbb{E}[X] := \int_{\mathbb{S}} x f_X(x) dx \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx \quad \right]$$

$$\mu_k := \mathbb{E}[(X - \mu)^k] \quad \text{higher-order central moments}$$

$$\text{Var}(X) := \mu_2 = \mathbb{E}[(X - \mu)^2] = \mathbb{E}[X^2 - 2\mu X + \mu^2] \quad \text{variance (2nd central moments)}$$

# moments

def.: Be  $X : \Omega \rightarrow \mathbb{S}$  a discrete random variable and  $p_X(x)$  its probability mass function (PMF).

$$\mu := \mathbb{E}[X] := \sum_{x_i \in \mathbb{S}} x_i p_X(x_i) \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \sum_{x_i \in \mathbb{S}} g(x_i) p_X(x_i) \quad \right]$$

exercise:

$$x \in \mathbb{S} := (1, 2, \dots, 7), \quad p_X(x) := \frac{1}{7}, \quad g(x) := \begin{cases} 0 & \text{if } x \in \{1, \dots, 5\} \\ 1 & \text{if } x \in \{6, 7\} \end{cases}$$

$$\begin{aligned} \mathbb{E}[X] &= 1 \cdot p_X(1) + \dots + 7 \cdot p_X(7) \\ &= (1 + \dots + 7) \cdot \frac{1}{7} = \frac{28}{7} = 4 \end{aligned}$$

$$\begin{aligned} \mathbb{E}[g(X)] &= g(1) \cdot p_X(1) + \dots + g(7) \cdot p_X(7) \\ &= (0 + 0 + 0 + 0 + 0 + 1 + 1) \cdot \frac{1}{7} = \frac{2}{7} \end{aligned}$$

Be  $X : \Omega \rightarrow \mathbb{S}$  a continuous random variable and  $f_X(x)$  its probability density function (PDF).

$$\mu := \mathbb{E}[X] := \int_{\mathbb{S}} x f_X(x) dx \quad \text{expectation/ expected value} \quad \left[ \quad \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx \quad \right]$$

$$\mu_k := \mathbb{E}[(X - \mu)^k] \quad \text{higher-order central moments}$$

$$\text{Var}(X) := \mu_2 = \mathbb{E}[(X - \mu)^2] = \mathbb{E}[X^2 - 2\mu X + \mu^2] = \mathbb{E}[X^2] - \mu^2 \quad \text{variance (2nd central moments)}$$



# laws of large numbers

def.:

# laws of large numbers

def.: Given an a stochastic process

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$ .

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and



# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,  
(WLLN)

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,  $\lim_{n \rightarrow \infty} P\left(-\epsilon < (\bar{X}_n - \mu) < \epsilon\right) = 1$   
(WLLN)

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,  $\lim_{n \rightarrow \infty} P\left(-\epsilon < (\bar{X}_n - \mu) < \epsilon\right) = 1 \quad \forall \epsilon > 0$ .  
(WLLN)

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,  $\lim_{n \rightarrow \infty} P\left(-\epsilon < (\bar{X}_n - \mu) < \epsilon\right) = 1 \quad \forall \epsilon > 0$ .  
(WLLN) (convergence of probabilities)

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,  $\lim_{n \rightarrow \infty} P\left(-\epsilon < (\bar{X}_n - \mu) < \epsilon\right) = 1 \quad \forall \epsilon > 0$ .  
(WLLN) (convergence of probabilities)

$\{X_n : n \in \mathbb{N}\}$



# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,  $\lim_{n \rightarrow \infty} P\left(-\epsilon < (\bar{X}_n - \mu) < \epsilon\right) = 1 \quad \forall \epsilon > 0$ .  
(WLLN) (convergence of probabilities)

$\{X_n : n \in \mathbb{N}\}$  follows the **strong law of large numbers** iff,

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,  $\lim_{n \rightarrow \infty} P\left(-\epsilon < (\bar{X}_n - \mu) < \epsilon\right) = 1 \quad \forall \epsilon > 0$ .  
(WLLN) (convergence of probabilities)

$\{X_n : n \in \mathbb{N}\}$  follows the **strong law of large numbers** iff,  
(SLLN)

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,  $\lim_{n \rightarrow \infty} P\left(-\epsilon < (\bar{X}_n - \mu) < \epsilon\right) = 1 \quad \forall \epsilon > 0$ .  
(WLLN) (convergence of probabilities)

$\{X_n : n \in \mathbb{N}\}$  follows the **strong law of large numbers** iff,  $P\left(\lim_{n \rightarrow \infty} \bar{X}_n = \mu\right) = 1$ .  
(SLLN)

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,  $\lim_{n \rightarrow \infty} P\left(-\epsilon < (\bar{X}_n - \mu) < \epsilon\right) = 1 \quad \forall \epsilon > 0$ .  
(WLLN) (convergence of probabilities)

$\{X_n : n \in \mathbb{N}\}$  follows the **strong law of large numbers** iff,  $P\left(\lim_{n \rightarrow \infty} \bar{X}_n = \mu\right) = 1$ .  
(SLLN) (behavior at the limit)

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,  $\lim_{n \rightarrow \infty} P\left(-\epsilon < (\bar{X}_n - \mu) < \epsilon\right) = 1 \quad \forall \epsilon > 0$ .  
(WLLN) (convergence of probabilities)

$\{X_n : n \in \mathbb{N}\}$  follows the **strong law of large numbers** iff,  $P\left(\lim_{n \rightarrow \infty} \bar{X}_n = \mu\right) = 1$ .  
(SLLN) (behavior at the limit)

theorem:

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,  $\lim_{n \rightarrow \infty} P\left(-\epsilon < (\bar{X}_n - \mu) < \epsilon\right) = 1 \quad \forall \epsilon > 0$ .  
(WLLN) (convergence of probabilities)

$\{X_n : n \in \mathbb{N}\}$  follows the **strong law of large numbers** iff,  $P\left(\lim_{n \rightarrow \infty} \bar{X}_n = \mu\right) = 1$ .  
(SLLN) (behavior at the limit)

**theorem:**

The **strong law of large numbers** implies the **weak law of large numbers**.

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,  $\lim_{n \rightarrow \infty} P\left(-\epsilon < (\bar{X}_n - \mu) < \epsilon\right) = 1 \quad \forall \epsilon > 0$ .  
(WLLN) (convergence of probabilities)

$\{X_n : n \in \mathbb{N}\}$  follows the **strong law of large numbers** iff,  $P\left(\lim_{n \rightarrow \infty} \bar{X}_n = \mu\right) = 1$ .  
(SLLN) (behavior at the limit)

**theorem:**

The **strong law of large numbers** implies the **weak law of large numbers**. (SLLN  $\rightarrow$  WLLN)

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,  $\lim_{n \rightarrow \infty} P\left(-\epsilon < (\bar{X}_n - \mu) < \epsilon\right) = 1 \quad \forall \epsilon > 0$ .  
(WLLN) (convergence of probabilities)

$\{X_n : n \in \mathbb{N}\}$  follows the **strong law of large numbers** iff,  $P\left(\lim_{n \rightarrow \infty} \bar{X}_n = \mu\right) = 1$ .  
(SLLN) (behavior at the limit)

**theorem:**

The **strong law of large numbers** implies the **weak law of large numbers**. (SLLN  $\rightarrow$  WLLN)

**theorem:**



# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,  $\lim_{n \rightarrow \infty} P\left(-\epsilon < (\bar{X}_n - \mu) < \epsilon\right) = 1 \quad \forall \epsilon > 0$ .  
(WLLN) (convergence of probabilities)

$\{X_n : n \in \mathbb{N}\}$  follows the **strong law of large numbers** iff,  $P\left(\lim_{n \rightarrow \infty} \bar{X}_n = \mu\right) = 1$ .  
(SLLN) (behavior at the limit)

**theorem:**

The **strong law of large numbers** implies the **weak law of large numbers**. (SLLN  $\rightarrow$  WLLN)

**theorem:**

If

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,  $\lim_{n \rightarrow \infty} P\left(-\epsilon < (\bar{X}_n - \mu) < \epsilon\right) = 1 \quad \forall \epsilon > 0$ .  
(WLLN) (convergence of probabilities)

$\{X_n : n \in \mathbb{N}\}$  follows the **strong law of large numbers** iff,  $P\left(\lim_{n \rightarrow \infty} \bar{X}_n = \mu\right) = 1$ .  
(SLLN) (behavior at the limit)

**theorem:**

The **strong law of large numbers** implies the **weak law of large numbers**. (SLLN  $\rightarrow$  WLLN)

**theorem:**

If  $\{X_n : n \in \mathbb{N}\}$

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,  $\lim_{n \rightarrow \infty} P\left(-\epsilon < (\bar{X}_n - \mu) < \epsilon\right) = 1 \quad \forall \epsilon > 0$ .  
(WLLN) (convergence of probabilities)

$\{X_n : n \in \mathbb{N}\}$  follows the **strong law of large numbers** iff,  $P\left(\lim_{n \rightarrow \infty} \bar{X}_n = \mu\right) = 1$ .  
(SLLN) (behavior at the limit)

**theorem:**

The **strong law of large numbers** implies the **weak law of large numbers**. (SLLN  $\rightarrow$  WLLN)

**theorem:**

If  $\{X_n : n \in \mathbb{N}\}$  is an i.i.d. process and

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,  $\lim_{n \rightarrow \infty} P\left(-\epsilon < (\bar{X}_n - \mu) < \epsilon\right) = 1 \quad \forall \epsilon > 0$ .  
(WLLN) (convergence of probabilities)

$\{X_n : n \in \mathbb{N}\}$  follows the **strong law of large numbers** iff,  $P\left(\lim_{n \rightarrow \infty} \bar{X}_n = \mu\right) = 1$ .  
(SLLN) (behavior at the limit)

**theorem:**

The **strong law of large numbers** implies the **weak law of large numbers**. (SLLN  $\rightarrow$  WLLN)

**theorem:**

If  $\{X_n : n \in \mathbb{N}\}$  is an i.i.d. process and  $\mathbb{E}(\bar{X}_n)$

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,  $\lim_{n \rightarrow \infty} P\left(-\epsilon < (\bar{X}_n - \mu) < \epsilon\right) = 1 \quad \forall \epsilon > 0$ .  
(WLLN) (convergence of probabilities)

$\{X_n : n \in \mathbb{N}\}$  follows the **strong law of large numbers** iff,  $P\left(\lim_{n \rightarrow \infty} \bar{X}_n = \mu\right) = 1$ .  
(SLLN) (behavior at the limit)

**theorem:**

The **strong law of large numbers** implies the **weak law of large numbers**. (SLLN  $\rightarrow$  WLLN)

**theorem:**

If  $\{X_n : n \in \mathbb{N}\}$  is an i.i.d. process and  $\mathbb{E}(\bar{X}_n)$  exists then

# laws of large numbers

def.: Given an a stochastic process  $\{X_n : n \in \mathbb{N}\}$  with  $X_n : \Omega \rightarrow \mathbb{S}$ .

Be  $\mu = \mathbb{E}[X_0] = \mathbb{E}[X_1] = \dots$  and  $\bar{X}_n := \frac{1}{n} \sum_{i=0}^{n-1} X_i$ .

$\{X_n : n \in \mathbb{N}\}$  follows the **weak law of large numbers** iff,  $\lim_{n \rightarrow \infty} P\left(-\epsilon < (\bar{X}_n - \mu) < \epsilon\right) = 1 \quad \forall \epsilon > 0$ .  
(WLLN) (convergence of probabilities)

$\{X_n : n \in \mathbb{N}\}$  follows the **strong law of large numbers** iff,  $P\left(\lim_{n \rightarrow \infty} \bar{X}_n = \mu\right) = 1$ .  
(SLLN) (behavior at the limit)

**theorem:**

The **strong law of large numbers** implies the **weak law of large numbers**. (SLLN  $\rightarrow$  WLLN)

**theorem:**

If  $\{X_n : n \in \mathbb{N}\}$  is an i.i.d. process and  $\mathbb{E}(\bar{X}_n)$  exists then **strong law of large numbers** holds.


# Monte Carlo integration (intuition)

# Monte Carlo integration (intuition)

$$\int_a^b g(x) dx$$




# Monte Carlo integration (intuition)




$$\int_a^b g(x) dx$$


# Monte Carlo integration (intuition)




# Monte Carlo integration (intuition)



# Monte Carlo integration (intuition)




$$\int_a^b g(x) dx$$




$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$

# Monte Carlo integration (intuition)




$$\int_a^b g(x) dx$$





$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$

$$\Delta x = \frac{b-a}{n}$$

# Monte Carlo integration (intuition)




$$\int_a^b g(x) dx$$




$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$

$$\Delta x = \frac{b-a}{n}$$

# Monte Carlo integration (intuition)




$$\int_a^b g(x) dx$$




$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$

$$\Delta x = \frac{b-a}{n}$$




$$= \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} g(x_i) \frac{b-a}{n}$$

# Monte Carlo integration (intuition)




$$\int_a^b g(x) dx$$



$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$


$$\Delta x = \frac{b-a}{n}$$




$$= \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} g(x_i) \frac{b-a}{n}$$



# Monte Carlo integration (intuition)




$$\int_a^b g(x) dx$$




$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$

$$\Delta x = \frac{b-a}{n}$$




$$= \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} g(x_i) \frac{b-a}{n}$$

# Monte Carlo integration (intuition)




$$\int_a^b g(x) dx$$



$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$


$$\Delta x = \frac{b-a}{n}$$




$$= \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} g(x_i) \frac{b-a}{n}$$

$$= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(x_i)$$

# Monte Carlo integration (intuition)




$$\int_a^b g(x) dx$$



$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$


$$\Delta x = \frac{b-a}{n}$$




$$= \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} g(x_i) \frac{b-a}{n}$$

$$= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(x_i)$$

# Monte Carlo integration (intuition)




$$\int_a^b g(x) dx$$




$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$

$$\Delta x = \frac{b-a}{n}$$




$$= \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} g(x_i) \frac{b-a}{n}$$


$$= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(x_i)$$



# Monte Carlo integration (intuition)




$$\int_a^b g(x) dx$$




$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$

$$\Delta x = \frac{b-a}{n}$$




$$= \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} g(x_i) \frac{b-a}{n}$$

$$= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(x_i)$$



# Monte Carlo integration (intuition)



$$\int_a^b g(x) dx$$

$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$


$$\Delta x = \frac{b-a}{n}$$

$$= \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} g(x_i) \frac{b-a}{n}$$

$$= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(x_i)$$



# Monte Carlo integration (intuition)



$$\int_a^b g(x) dx$$

$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$


$$\Delta x = \frac{b-a}{n}$$

$$= \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} g(x_i) \frac{b-a}{n}$$

$$= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(x_i)$$

(SLLN)

# Monte Carlo integration (intuition)



$$\int_a^b g(x) dx$$

$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$

$$\Delta x = \frac{b-a}{n}$$


$$= \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} g(x_i) \frac{b-a}{n}$$

$$= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(x_i)$$


(SLLN)  $1 = P \left( \lim_{n \rightarrow \infty} \bar{Y}_n = \mu \right)$



# Monte Carlo integration (intuition)




$$\int_a^b g(x) dx$$




$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$

$$\Delta x = \frac{b-a}{n}$$



$$= \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} g(x_i) \frac{b-a}{n}$$





$$= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(x_i)$$



$$\bar{Y}_n := \frac{1}{n} \sum_{i=1}^n Y_i$$

(SLLN)  $1 = P \left( \lim_{n \rightarrow \infty} \bar{Y}_n = \mu \right)$

# Monte Carlo integration (intuition)



$$\int_a^b g(x) dx$$

$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$


$$\Delta x = \frac{b-a}{n}$$

$$= \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} g(x_i) \frac{b-a}{n}$$

$$= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(x_i)$$

$$\begin{aligned} \bar{Y}_n &:= \frac{1}{n} \sum_{i=1}^n Y_i \\ \text{(SLLN)} \quad 1 &= P \left( \lim_{n \rightarrow \infty} \bar{Y}_n = \mu \right) = P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} Y_i = \mu \right) \end{aligned}$$

# Monte Carlo integration (intuition)



$$\int_a^b g(x) dx$$

$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$

$$\Delta x = \frac{b-a}{n}$$

$$= \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} g(x_i) \frac{b-a}{n}$$

$$= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(x_i)$$


(SLLN)

$$1 = P \left( \lim_{n \rightarrow \infty} \bar{Y}_n = \mu \right)$$

$$= P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} Y_i = \mu \right)$$

$\bar{Y}_n := \frac{1}{n} \sum_{i=1}^n Y_i$  $Y := g(X)$

# Monte Carlo integration (intuition)



$$\int_a^b g(x) dx$$

$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$

$$\Delta x = \frac{b-a}{n}$$

$$= \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} g(x_i) \frac{b-a}{n}$$

$$= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(x_i)$$


$\bar{Y}_n := \frac{1}{n} \sum_{i=1}^n Y_i$

$Y := g(X) \quad \mu = \mathbb{E}[Y] = \mathbb{E}[g(X)]$

(SLLN)

$1 = P \left( \lim_{n \rightarrow \infty} \bar{Y}_n = \mu \right) = P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} Y_i = \mu \right)$

# Monte Carlo integration (intuition)



$$\int_a^b g(x) dx$$

$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$

$$\Delta x = \frac{b-a}{n}$$

$$= \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} g(x_i) \frac{b-a}{n}$$

$$= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(x_i)$$

$$\bar{Y}_n := \frac{1}{n} \sum_{i=1}^n Y_i$$

$$Y := g(X) \quad \mu = \mathbb{E}[Y] = \mathbb{E}[g(X)]$$

$$X \sim \mathcal{U}(a, b)$$

(SLLN)


1 = P

$$\left( \lim_{n \rightarrow \infty} \bar{Y}_n = \mu \right)$$

= P

$$\left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} Y_i = \mu \right)$$

# Monte Carlo integration (intuition)



$$\int_a^b g(x) dx$$

$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$

$$\Delta x = \frac{b-a}{n}$$

$$= \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} g(x_i) \frac{b-a}{n}$$

$$= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(x_i)$$

$\bar{Y}_n := \frac{1}{n} \sum_{i=1}^n Y_i$


$Y := g(X) \quad \mu = \mathbb{E}[Y] = \mathbb{E}[g(X)]$

$X \sim \mathcal{U}(a, b)$

(SLLN)

$$1 = P \left( \lim_{n \rightarrow \infty} \bar{Y}_n = \mu \right) = P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} Y_i = \mu \right) = P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right)$$

# Monte Carlo integration (intuition)



$$\bar{Y}_n := \frac{1}{n} \sum_{i=1}^n Y_i$$


$$Y := g(X) \quad \mu = \mathbb{E}[Y] = \mathbb{E}[g(X)] \quad X \sim \mathcal{U}(a, b)$$

$$(SLLN) \quad 1 = P \left( \lim_{n \rightarrow \infty} \bar{Y}_n = \mu \right)$$

$$= P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} Y_i = \mu \right)$$

$$= P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right)$$

# Monte Carlo integration (intuition)



$(SLLN)$

$1 = P\left(\lim_{n \rightarrow \infty} \bar{Y}_n = \mu\right)$

$\bar{Y}_n := \frac{1}{n} \sum_{i=1}^n Y_i$

$= P\left(\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} Y_i = \mu\right)$

$Y := g(X)$





$= P\left(\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)]\right)$

$\mu = \mathbb{E}[Y] = \mathbb{E}[g(X)]$

$X \sim \mathcal{U}(a, b)$



# Monte Carlo integration (intuition)



$$\int_a^b g(x) dx$$

$$= \lim_{\Delta x \rightarrow 0} \sum_{i=0}^{n-1} g(x_i) \Delta x$$

$$\Delta x = \frac{b-a}{n}$$

$$= \lim_{n \rightarrow \infty} \sum_{i=0}^{n-1} g(x_i) \frac{b-a}{n}$$

$$= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(x_i)$$

$$= (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i)$$

$$\approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i)$$

$$\begin{aligned} \bar{Y}_n &:= \frac{1}{n} \sum_{i=1}^n Y_i & Y &:= g(X) & \mu &= \mathbb{E}[Y] = \mathbb{E}[g(X)] & X &\sim \mathcal{U}(a, b) \\ \text{(SLLN)} \quad 1 &= P \left( \lim_{n \rightarrow \infty} \bar{Y}_n = \mu \right) & &= P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} Y_i = \mu \right) & &= P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) \end{aligned}$$

# Monte Carlo integration (derivation)

# Monte Carlo integration (derivation)

$$\int_a^b g(x) dx$$

# Monte Carlo integration (derivation)

$$\int_a^b g(x) dx = \int_a^b g(x) 1 dx$$

# Monte Carlo integration (derivation)

$$\int_a^b g(x) dx = \int_a^b g(x) 1 dx = \int_a^b g(x) \frac{b-a}{b-a} dx$$

# Monte Carlo integration (derivation)

$$\int_a^b g(x) dx = \int_a^b g(x) 1 dx = \int_a^b g(x) \frac{b-a}{b-a} dx = (b-a) \int_a^b g(x) \frac{1}{b-a} dx$$

# Monte Carlo integration (derivation)

$$\int_a^b g(x) dx = \int_a^b g(x) 1 dx = \int_a^b g(x) \frac{b-a}{b-a} dx = (b-a) \int_a^b g(x) \overbrace{\frac{1}{b-a}}^{=f_X(x)} dx$$

# Monte Carlo integration (derivation)

$$\int_a^b g(x) dx = \int_a^b g(x) 1 dx = \int_a^b g(x) \frac{b-a}{b-a} dx = (b-a) \int_a^b g(x) \overbrace{\frac{1}{b-a}}^{=f_X(x) \quad X \sim \mathcal{U}(a,b)} dx$$



# Monte Carlo integration (derivation)

$$\int_a^b g(x) dx = \int_a^b g(x) 1 dx = \int_a^b g(x) \frac{b-a}{b-a} dx = (b-a) \int_a^b g(x) \overbrace{\frac{1}{b-a}}^{=f_X(x) \quad X \sim \mathcal{U}(a,b)} dx \quad \left[ \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx \right]$$

# Monte Carlo integration (derivation)

$$\begin{aligned}
 \int_a^b g(x) dx &= \int_a^b g(x) \cdot 1 dx = \int_a^b g(x) \frac{b-a}{b-a} dx = (b-a) \int_a^b g(x) \overbrace{\frac{1}{b-a}}^{=f_X(x) \quad X \sim \mathcal{U}(a,b)} dx & \left[ \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx \right] \\
 &= (b-a) \mathbb{E}[g(X)] & \text{SLLN: } P\left(\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)]\right) = 1
 \end{aligned}$$

# Monte Carlo integration (derivation)

$$\int_a^b g(x) \, dx = \int_a^b g(x) \, 1 \, dx = \int_a^b g(x) \frac{b-a}{b-a} \, dx = (b-a) \int_a^b g(x) \overbrace{\frac{1}{b-a}}^{= f_X(x) \quad X \sim \mathcal{U}(a,b)} \, dx \quad \left[ \quad \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) \, dx \quad \right]$$

$$= (b-a) \mathbb{E}[g(X)] \qquad \text{SLLN:} \quad P\left(\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)]\right) = 1$$

=  
almost  
always

# Monte Carlo integration (derivation)

$$\begin{aligned}
 \int_a^b g(x) dx &= \int_a^b g(x) 1 dx = \int_a^b g(x) \frac{b-a}{b-a} dx = (b-a) \int_a^b g(x) \overbrace{\frac{1}{b-a}}^{=f_X(x) \quad X \sim \mathcal{U}(a,b)} dx & \left[ \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx \right] \\
 &= (b-a) \mathbb{E}[g(X)] & \text{SLLN: } P\left(\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)]\right) = 1 \\
 &\stackrel{\text{almost always}}{=} (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i)
 \end{aligned}$$

# Monte Carlo integration (derivation)

$$\begin{aligned}
 \int_a^b g(x) dx &= \int_a^b g(x) 1 dx = \int_a^b g(x) \frac{b-a}{b-a} dx = (b-a) \int_a^b g(x) \overbrace{\frac{1}{b-a}}^{=f_X(x) \quad X \sim \mathcal{U}(a,b)} dx & \left[ \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx \right] \\
 &= (b-a) \mathbb{E}[g(X)] & \text{SLLN: } P\left(\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)]\right) = 1 \\
 &\stackrel{\text{almost always}}{=} (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) \\
 &\approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i)
 \end{aligned}$$

# Monte Carlo integration (derivation)

$$\begin{aligned}
 \int_a^b g(x) dx &= \int_a^b g(x) 1 dx = \int_a^b g(x) \frac{b-a}{b-a} dx = (b-a) \int_a^b g(x) \overbrace{\frac{1}{b-a}}^{=f_X(x) \quad X \sim \mathcal{U}(a,b)} dx & \left[ \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx \right] \\
 &= (b-a) \mathbb{E}[g(X)] & \text{SLLN: } P\left(\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)]\right) = 1 \\
 &\stackrel{\text{almost always}}{=} (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) \\
 &\approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n
 \end{aligned}$$

# Monte Carlo integration (derivation)

$$\begin{aligned}
 \int_a^b g(x) dx &= \int_a^b g(x) 1 dx = \int_a^b g(x) \frac{b-a}{b-a} dx = (b-a) \int_a^b g(x) \overbrace{\frac{1}{b-a}}^{= f_X(x) \quad X \sim \mathcal{U}(a,b)} dx & \left[ \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx \right] \\
 &= (b-a) \mathbb{E}[g(X)] & \text{SLLN: } P\left(\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)]\right) = 1 \\
 &\stackrel{\text{almost always}}{=} (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) \\
 &\approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n \quad \text{"Monte Carlo estimator"}
 \end{aligned}$$

# Monte Carlo integration (derivation)

$$\begin{aligned}
 \int_a^b g(x) dx &= \int_a^b g(x) 1 dx = \int_a^b g(x) \frac{b-a}{b-a} dx = (b-a) \int_a^b g(x) \overbrace{\frac{1}{b-a}}^{= f_X(x) \quad X \sim \mathcal{U}(a,b)} dx & \left[ \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx \right] \\
 &= (b-a) \mathbb{E}[g(X)] & \text{SLLN: } P\left(\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)]\right) = 1 \\
 &\stackrel{\text{almost always}}{=} (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) \\
 &\approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n \quad \text{"Monte Carlo estimator"}
 \end{aligned}$$

multi-dimensional



# Monte Carlo integration (derivation)

$$\begin{aligned}
 \int_a^b g(x) dx &= \int_a^b g(x) 1 dx = \int_a^b g(x) \frac{b-a}{b-a} dx = (b-a) \int_a^b g(x) \overbrace{\frac{1}{b-a}}^{=f_X(x)} dx && X \sim \mathcal{U}(a,b) \\
 &&& \left[ \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx \right] \\
 &= (b-a) \mathbb{E}[g(X)] && \text{SLLN: } P\left(\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)]\right) = 1 \\
 &\stackrel{\text{almost always}}{=} (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) \\
 &\approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n && \text{"Monte Carlo estimator"}
 \end{aligned}$$

multi-dimensional

$$\int_{\mathbb{S}} g(\underline{x}) d\underline{x}$$

# Monte Carlo integration (derivation)

$$\begin{aligned}
 \int_a^b g(x) dx &= \int_a^b g(x) 1 dx = \int_a^b g(x) \frac{b-a}{b-a} dx = (b-a) \int_a^b g(x) \overbrace{\frac{1}{b-a}}^{=f_X(x) \quad X \sim \mathcal{U}(a,b)} dx & \left[ \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx \right] \\
 &= (b-a) \mathbb{E}[g(X)] & \text{SLLN: } P\left(\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)]\right) = 1 \\
 &\stackrel{\text{almost always}}{=} (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) \\
 &\approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n \quad \text{"Monte Carlo estimator"}
 \end{aligned}$$

multi-dimensional

$$\int_{\mathbb{S}} g(\underline{x}) d\underline{x} \approx V \frac{1}{n} \sum_{i=0}^{n-1} g(\underline{X}_i)$$

# Monte Carlo integration (derivation)

$$\begin{aligned}
 \int_a^b g(x) dx &= \int_a^b g(x) 1 dx = \int_a^b g(x) \frac{b-a}{b-a} dx = (b-a) \int_a^b g(x) \overbrace{\frac{1}{b-a}}^{=f_X(x) \quad X \sim \mathcal{U}(a,b)} dx & \left[ \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx \right] \\
 &= (b-a) \mathbb{E}[g(X)] & \text{SLLN: } P\left(\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)]\right) = 1 \\
 &\stackrel{\text{almost always}}{=} (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) \\
 &\approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n \quad \text{"Monte Carlo estimator"}
 \end{aligned}$$

multi-dimensional

$$\int_{\mathbb{S}} g(\underline{x}) d\underline{x} \stackrel{\text{(volume)}}{\approx} V \frac{1}{n} \sum_{i=0}^{n-1} g(\underline{X}_i)$$

# Monte Carlo integration (derivation)


$$\begin{aligned}
 \int_a^b g(x) dx &= \int_a^b g(x) 1 dx = \int_a^b g(x) \frac{b-a}{b-a} dx = (b-a) \int_a^b g(x) \overbrace{\frac{1}{b-a}}^{=f_X(x) \quad X \sim \mathcal{U}(a,b)} dx & \left[ \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx \right] \\
 &= (b-a) \mathbb{E}[g(X)] & \text{SLLN: } P\left(\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)]\right) = 1 \\
 &\stackrel{\text{almost always}}{=} (b-a) \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) \\
 &\approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n \quad \text{"Monte Carlo estimator"}
 \end{aligned}$$

multi-dimensional

$$\int_{\mathbb{S}} g(\underline{x}) d\underline{x} \stackrel{\text{(volume)}}{\approx} V \frac{1}{n} \sum_{i=0}^{n-1} g(\underline{X}_i) =: \vartheta_n$$


# Monte Carlo integration (alternative like accept-reject sampling)

# Monte Carlo integration (alternative like accept-reject sampling)




# Monte Carlo integration (alternative like accept-reject sampling)

$x \in [a, b]$



# Monte Carlo integration (alternative like accept-reject sampling)

$$x \in [a, b]$$
$$g(x) \in [0, \beta]$$







# Monte Carlo integration (alternative like accept-reject sampling)

$$\begin{aligned} x &\in [a, b] \\ g(x) &\in [0, \beta] \end{aligned}$$

$$\mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$



# Monte Carlo integration (alternative like accept-reject sampling)



$$\begin{aligned} x &\in [a, b] \\ g(x) &\in [0, \beta] \end{aligned} \qquad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$


$$\int_a^b g(x) dx$$

# Monte Carlo integration (alternative like accept-reject sampling)

$$\begin{aligned}x &\in [a, b] \\ g(x) &\in [0, \beta]\end{aligned}$$

$$\mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$

$$\int_a^b g(x) dx = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) dx dy$$




# Monte Carlo integration (alternative like accept-reject sampling)

$$\begin{aligned}x &\in [a, b] \\ g(x) &\in [0, \beta]\end{aligned}$$

$$\mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$

$$\int_a^b g(x) dx = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) 1 dx dy$$




# Monte Carlo integration (alternative like accept-reject sampling)


$$\begin{aligned}x &\in [a, b] \\ g(x) &\in [0, \beta]\end{aligned}$$

$$\mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$

$$\int_a^b g(x) dx = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) 1 dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \frac{(\beta - 0)(b - a)}{(\beta - 0)(b - a)} dx dy$$



# Monte Carlo integration (alternative like accept-reject sampling)




$$\begin{aligned} x &\in [a, b] \\ g(x) &\in [0, \beta] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$
$$\int_a^b g(x) dx = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) 1 dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \frac{(\beta - 0)(b - a)}{(\beta - 0)(b - a)} dx dy$$

$$= \beta(b - a) \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \frac{1}{\beta(b - a)} dx dy$$

# Monte Carlo integration (alternative like accept-reject sampling)


$$\begin{aligned} x &\in [a, b] \\ g(x) &\in [0, \beta] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$



$$\int_a^b g(x) dx = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) 1 dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \frac{(\beta - 0)(b - a)}{(\beta - 0)(b - a)} dx dy$$

$$= \beta(b - a) \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \underbrace{\frac{1}{\beta(b - a)}}_{= f_{XY}(x, y)} dx dy$$

# Monte Carlo integration (alternative like accept-reject sampling)




$$\begin{aligned} x &\in [a, b] \\ g(x) &\in [0, \beta] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$

$$\int_a^b g(x) dx = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) 1 dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \frac{(\beta - 0)(b - a)}{(\beta - 0)(b - a)} dx dy$$

$$= \beta(b - a) \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \underbrace{\frac{1}{\beta(b - a)}}_{= f_{XY}(x, y)} dx dy = \beta(b - a) \mathbb{E}[\mathbb{I}_g(X, Y)]$$




# Monte Carlo integration (alternative like accept-reject sampling)



$$\begin{aligned} x &\in [a, b] \\ g(x) &\in [0, \beta] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$

$$\begin{aligned} \int_a^b g(x) dx &= \int_0^\beta \int_a^b \mathbb{I}_g(x, y) dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) 1 dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \frac{(\beta - 0)(b - a)}{(\beta - 0)(b - a)} dx dy \\ &= \beta(b - a) \int_0^\beta \int_a^b \underbrace{\mathbb{I}_g(x, y) \frac{1}{\beta(b - a)}}_{= f_{XY}(x, y)} dx dy = \beta(b - a) \mathbb{E}[\mathbb{I}_g(X, Y)] \approx \beta(b - a) \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{I}_g(X_i, Y_i) \end{aligned}$$

# Monte Carlo integration (alternative like accept-reject sampling)




$$\begin{aligned} x &\in [a, b] \\ g(x) &\in [0, \beta] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$

$$\int_a^b g(x) dx = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) 1 dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \frac{(\beta - 0)(b - a)}{(\beta - 0)(b - a)} dx dy$$

$$= \beta(b - a) \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \underbrace{\frac{1}{\beta(b - a)}}_{= f_{XY}(x, y)} dx dy = \beta(b - a) \mathbb{E}[\mathbb{I}_g(X, Y)] \approx \beta(b - a) \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{I}_g(X_i, Y_i) \quad X_i \sim \mathcal{U}(a, b)$$

# Monte Carlo integration (alternative like accept-reject sampling)




$$\begin{aligned} x &\in [a, b] \\ g(x) &\in [0, \beta] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$

$$\int_a^b g(x) dx = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) 1 dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \frac{(\beta - 0)(b - a)}{(\beta - 0)(b - a)} dx dy$$

$$= \beta(b - a) \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \underbrace{\frac{1}{\beta(b - a)}}_{= f_{XY}(x, y)} dx dy = \beta(b - a) \mathbb{E}[\mathbb{I}_g(X, Y)] \approx \beta(b - a) \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{I}_g(X_i, Y_i) \quad \begin{aligned} X_i &\sim \mathcal{U}(a, b) \\ Y_i &\sim \mathcal{U}(0, \beta) \end{aligned}$$

# Monte Carlo integration (alternative like accept-reject sampling)




$$\begin{aligned} x &\in [a, b] \\ g(x) &\in [0, \beta] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$

$$\int_a^b g(x) dx = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) 1 dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \frac{(\beta - 0)(b - a)}{(\beta - 0)(b - a)} dx dy$$

$$= \beta(b - a) \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \underbrace{\frac{1}{\beta(b - a)}}_{= f_{XY}(x, y)} dx dy = \beta(b - a) \mathbb{E}[\mathbb{I}_g(X, Y)] \approx \beta(b - a) \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{I}_g(X_i, Y_i) \quad \begin{aligned} X_i &\sim \mathcal{U}(a, b) \\ Y_i &\sim \mathcal{U}(0, \beta) \end{aligned}$$

example:

# Monte Carlo integration (alternative like accept-reject sampling)




$$\begin{aligned} x &\in [a, b] \\ g(x) &\in [0, \beta] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$


$$\int_a^b g(x) dx = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) 1 dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \frac{(\beta - 0)(b - a)}{(\beta - 0)(b - a)} dx dy$$

$$= \beta(b - a) \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \underbrace{\frac{1}{\beta(b - a)}}_{= f_{XY}(x, y)} dx dy = \beta(b - a) \mathbb{E}[\mathbb{I}_g(X, Y)] \approx \beta(b - a) \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{I}_g(X_i, Y_i) \quad \begin{aligned} X_i &\sim \mathcal{U}(a, b) \\ Y_i &\sim \mathcal{U}(0, \beta) \end{aligned}$$

example:




# Monte Carlo integration (alternative like accept-reject sampling)



$$\begin{aligned} x &\in [a, b] \\ g(x) &\in [0, \beta] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$


$$\begin{aligned} \int_a^b g(x) dx &= \int_0^\beta \int_a^b \mathbb{I}_g(x, y) dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) 1 dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \frac{(\beta - 0)(b - a)}{(\beta - 0)(b - a)} dx dy \\ &= \beta(b - a) \int_0^\beta \int_a^b \underbrace{\mathbb{I}_g(x, y) \frac{1}{\beta(b - a)}}_{= f_{XY}(x, y)} dx dy = \beta(b - a) \mathbb{E}[\mathbb{I}_g(X, Y)] \approx \beta(b - a) \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{I}_g(X_i, Y_i) \quad \begin{aligned} X_i &\sim \mathcal{U}(a, b) \\ Y_i &\sim \mathcal{U}(0, \beta) \end{aligned} \end{aligned}$$

example:



$$x \in [0, 1]$$

# Monte Carlo integration (alternative like accept-reject sampling)




$$\begin{aligned} x &\in [a, b] \\ g(x) &\in [0, \beta] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$

$$\int_a^b g(x) dx = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) 1 dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \frac{(\beta - 0)(b - a)}{(\beta - 0)(b - a)} dx dy$$

$$= \beta(b - a) \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \underbrace{\frac{1}{\beta(b - a)}}_{= f_{XY}(x, y)} dx dy = \beta(b - a) \mathbb{E}[\mathbb{I}_g(X, Y)] \approx \beta(b - a) \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{I}_g(X_i, Y_i) \quad \begin{aligned} X_i &\sim \mathcal{U}(a, b) \\ Y_i &\sim \mathcal{U}(0, \beta) \end{aligned}$$


example:



$$\begin{aligned} x &\in [0, 1] \\ g(x) &\in [0, 1] \end{aligned}$$



# Monte Carlo integration (alternative like accept-reject sampling)




$$\begin{aligned} x &\in [a, b] \\ g(x) &\in [0, \beta] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$

$$\int_a^b g(x) dx = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) 1 dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \frac{(\beta - 0)(b - a)}{(\beta - 0)(b - a)} dx dy$$

$$= \beta(b - a) \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \underbrace{\frac{1}{\beta(b - a)}}_{= f_{XY}(x, y)} dx dy = \beta(b - a) \mathbb{E}[\mathbb{I}_g(X, Y)] \approx \beta(b - a) \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{I}_g(X_i, Y_i) \quad \begin{aligned} X_i &\sim \mathcal{U}(a, b) \\ Y_i &\sim \mathcal{U}(0, \beta) \end{aligned}$$


example:



$$\begin{aligned} x &\in [0, 1] \\ g(x) &\in [0, 1] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } x^2 + y^2 \leq 1 \\ 0 & \text{else} \end{cases}$$



# Monte Carlo integration (alternative like accept-reject sampling)




$$\begin{aligned} x &\in [a, b] \\ g(x) &\in [0, \beta] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$

$$\int_a^b g(x) dx = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) 1 dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \frac{(\beta - 0)(b - a)}{(\beta - 0)(b - a)} dx dy$$

$$= \beta(b - a) \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \underbrace{\frac{1}{\beta(b - a)}}_{= f_{XY}(x, y)} dx dy = \beta(b - a) \mathbb{E}[\mathbb{I}_g(X, Y)] \approx \beta(b - a) \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{I}_g(X_i, Y_i) \quad \begin{aligned} X_i &\sim \mathcal{U}(a, b) \\ Y_i &\sim \mathcal{U}(0, \beta) \end{aligned}$$


example:



$$\begin{aligned} x &\in [0, 1] \\ g(x) &\in [0, 1] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } x^2 + y^2 \leq 1 \\ 0 & \text{else} \end{cases}$$

$$Y_i, X_i \sim \mathcal{U}(0, 1)$$

# Monte Carlo integration (alternative like accept-reject sampling)




$$\begin{aligned} x &\in [a, b] \\ g(x) &\in [0, \beta] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$

$$\int_a^b g(x) dx = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) 1 dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \frac{(\beta - 0)(b - a)}{(\beta - 0)(b - a)} dx dy$$

$$= \beta(b - a) \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \underbrace{\frac{1}{\beta(b - a)}}_{= f_{XY}(x, y)} dx dy = \beta(b - a) \mathbb{E}[\mathbb{I}_g(X, Y)] \approx \beta(b - a) \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{I}_g(X_i, Y_i) \quad \begin{aligned} X_i &\sim \mathcal{U}(a, b) \\ Y_i &\sim \mathcal{U}(0, \beta) \end{aligned}$$


example:



$$\begin{aligned} x &\in [0, 1] \\ g(x) &\in [0, 1] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } x^2 + y^2 \leq 1 \\ 0 & \text{else} \end{cases} \quad Y_i, X_i \sim \mathcal{U}(0, 1)$$

$$\int_0^1 g(x) dx$$

# Monte Carlo integration (alternative like accept-reject sampling)




$$\begin{aligned} x &\in [a, b] \\ g(x) &\in [0, \beta] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$

$$\int_a^b g(x) dx = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) 1 dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \frac{(\beta - 0)(b - a)}{(\beta - 0)(b - a)} dx dy$$

$$= \beta(b - a) \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \underbrace{\frac{1}{\beta(b - a)}}_{= f_{XY}(x, y)} dx dy = \beta(b - a) \mathbb{E}[\mathbb{I}_g(X, Y)] \approx \beta(b - a) \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{I}_g(X_i, Y_i) \quad \begin{aligned} X_i &\sim \mathcal{U}(a, b) \\ Y_i &\sim \mathcal{U}(0, \beta) \end{aligned}$$

example:




$$\begin{aligned} x &\in [0, 1] \\ g(x) &\in [0, 1] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } x^2 + y^2 \leq 1 \\ 0 & \text{else} \end{cases}$$

$$Y_i, X_i \sim \mathcal{U}(0, 1)$$

$$\int_0^1 g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{I}_g(X_i, Y_i)$$

# Monte Carlo integration (alternative like accept-reject sampling)




$$\begin{aligned} x &\in [a, b] \\ g(x) &\in [0, \beta] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } y \leq g(x) \\ 0 & \text{else} \end{cases}$$

$$\int_a^b g(x) dx = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) 1 dx dy = \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \frac{(\beta - 0)(b - a)}{(\beta - 0)(b - a)} dx dy$$

$$= \beta(b - a) \int_0^\beta \int_a^b \mathbb{I}_g(x, y) \underbrace{\frac{1}{\beta(b - a)}}_{= f_{XY}(x, y)} dx dy = \beta(b - a) \mathbb{E}[\mathbb{I}_g(X, Y)] \approx \beta(b - a) \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{I}_g(X_i, Y_i) \quad \begin{aligned} X_i &\sim \mathcal{U}(a, b) \\ Y_i &\sim \mathcal{U}(0, \beta) \end{aligned}$$

example:



$$\begin{aligned} x &\in [0, 1] \\ g(x) &\in [0, 1] \end{aligned} \quad \mathbb{I}_g(x, y) := \begin{cases} 1 & \text{if } x^2 + y^2 \leq 1 \\ 0 & \text{else} \end{cases}$$

$$Y_i, X_i \sim \mathcal{U}(0, 1)$$

$$\int_0^1 g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \mathbb{I}_g(X_i, Y_i) \approx \frac{\pi}{4}$$

# Monte Carlo integration (error)

# Monte Carlo integration (error)

$error_{\vartheta_n}$

# Monte Carlo integration (error)

$$error_{\vartheta_n} \approx \sqrt{Var(\vartheta_n)}$$

# Monte Carlo integration (error)

$$error_{\vartheta_n} \approx \sqrt{Var(\vartheta_n)} = \sqrt{Var\left((b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i)\right)}$$



# Monte Carlo integration (error)

$$\begin{aligned} error_{\vartheta_n} &\approx \sqrt{Var(\vartheta_n)} = \sqrt{Var\left((b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i)\right)} \\ &= \frac{b-a}{n} \sqrt{Var\left(\sum_{i=0}^{n-1} g(X_i)\right)} \end{aligned}$$

# Monte Carlo integration (error)

$$\begin{aligned} error_{\vartheta_n} &\approx \sqrt{Var(\vartheta_n)} = \sqrt{Var\left((b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i)\right)} \\ &= \frac{b-a}{n} \sqrt{Var\left(\sum_{i=0}^{n-1} g(X_i)\right)} \quad X_i \text{ i.i.d.} \end{aligned}$$

# Monte Carlo integration (error)

$$\begin{aligned} error_{\vartheta_n} &\approx \sqrt{Var(\vartheta_n)} = \sqrt{Var\left((b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i)\right)} \\ &= \frac{b-a}{n} \sqrt{Var\left(\sum_{i=0}^{n-1} g(X_i)\right)} \quad X_i \text{ i.i.d.} \\ &= \frac{b-a}{n} \sqrt{\sum_{i=0}^{n-1} Var(g(X_i))} \end{aligned}$$

# Monte Carlo integration (error)

$$\begin{aligned} error_{\vartheta_n} &\approx \sqrt{Var(\vartheta_n)} = \sqrt{Var\left((b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i)\right)} \\ &= \frac{b-a}{n} \sqrt{Var\left(\sum_{i=0}^{n-1} g(X_i)\right)} \quad X_i \text{ i.i.d.} \\ &= \frac{b-a}{n} \sqrt{\sum_{i=0}^{n-1} Var(g(X_i))} \\ &= \frac{b-a}{n} \sqrt{n Var(g(X))} \end{aligned}$$

# Monte Carlo integration (error)

$$\begin{aligned} error_{\vartheta_n} &\approx \sqrt{Var(\vartheta_n)} = \sqrt{Var\left((b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i)\right)} \\ &= \frac{b-a}{n} \sqrt{Var\left(\sum_{i=0}^{n-1} g(X_i)\right)} \quad X_i \text{ i.i.d.} \\ &= \frac{b-a}{n} \sqrt{\sum_{i=0}^{n-1} Var(g(X_i))} \\ &= \frac{b-a}{n} \sqrt{n Var(g(X))} \\ &= \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))} \end{aligned}$$

# Monte Carlo integration (error)

$$\begin{aligned} error_{\vartheta_n} &\approx \sqrt{Var(\vartheta_n)} = \sqrt{Var\left((b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i)\right)} \\ &= \frac{b-a}{n} \sqrt{Var\left(\sum_{i=0}^{n-1} g(X_i)\right)} \quad X_i \text{ i.i.d.} \\ &= \frac{b-a}{n} \sqrt{\sum_{i=0}^{n-1} Var(g(X_i))} \\ &= \frac{b-a}{n} \sqrt{n \, Var(g(X))} \\ &= \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))} \propto \frac{1}{\sqrt{n}} \end{aligned}$$

# Monte Carlo integration (error)

$$error_{\vartheta_n} \approx \sqrt{Var(\vartheta_n)} = \sqrt{Var\left((b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i)\right)}$$

$$= \frac{b-a}{n} \sqrt{Var\left(\sum_{i=0}^{n-1} g(X_i)\right)} \quad X_i \text{ i.i.d.}$$

$$= \frac{b-a}{n} \sqrt{\sum_{i=0}^{n-1} Var(g(X_i))}$$

$$= \frac{b-a}{n} \sqrt{n Var(g(X))}$$

higher dimension error

$$= \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))} \propto \frac{1}{\sqrt{n}}$$

# Monte Carlo integration (error)

$$error_{\vartheta_n} \approx \sqrt{Var(\vartheta_n)} = \sqrt{Var\left((b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i)\right)}$$

$$= \frac{b-a}{n} \sqrt{Var\left(\sum_{i=0}^{n-1} g(X_i)\right)} \quad X_i \text{ i.i.d.}$$

$$= \frac{b-a}{n} \sqrt{\sum_{i=0}^{n-1} Var(g(X_i))}$$

$$= \frac{b-a}{n} \sqrt{n Var(g(X))}$$

higher dimension error

$$= \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))} \propto \frac{1}{\sqrt{n}}$$

$error_{\vartheta_n}$



# Monte Carlo integration (error)

$$error_{\vartheta_n} \approx \sqrt{Var(\vartheta_n)} = \sqrt{Var\left((b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i)\right)}$$

$$= \frac{b-a}{n} \sqrt{Var\left(\sum_{i=0}^{n-1} g(X_i)\right)} \quad X_i \text{ i.i.d.}$$

$$= \frac{b-a}{n} \sqrt{\sum_{i=0}^{n-1} Var(g(X_i))}$$

$$= \frac{b-a}{n} \sqrt{n Var(g(X))}$$

$$= \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))} \propto \frac{1}{\sqrt{n}}$$

higher dimension error

(volume)

$error_{\vartheta_n}$

# Monte Carlo integration (error)

$$\begin{aligned}
 error_{\vartheta_n} &\approx \sqrt{Var(\vartheta_n)} = \sqrt{Var\left((b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i)\right)} \\
 &= \frac{b-a}{n} \sqrt{Var\left(\sum_{i=0}^{n-1} g(X_i)\right)} \quad X_i \text{ i.i.d.} \\
 &= \frac{b-a}{n} \sqrt{\sum_{i=0}^{n-1} Var(g(X_i))} \\
 &= \frac{b-a}{n} \sqrt{n \, Var(g(X))} \\
 &= \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))} \propto \frac{1}{\sqrt{n}}
 \end{aligned}$$

higher dimension error

$$error_{\vartheta_n} \approx \frac{\overset{\text{(volume)}}{V}}{\sqrt{n}} \sqrt{Var(g(X))}$$

# Monte Carlo integration (importance sampling)

# Monte Carlo integration (importance sampling)

intuition:

# Monte Carlo integration (importance sampling)


intuition:

$$\int_a^b g(x) dx$$

# Monte Carlo integration (importance sampling)

intuition:


$$\int_a^b g(x) dx$$



# Monte Carlo integration (importance sampling)

intuition:


$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i$$



# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i$$






# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i$$

$$\Delta x = \frac{b-a}{n}$$




# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i}$$




# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$




# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \frac{importance(x_i)}{b-a}}$$

$$\Delta x = \frac{b-a}{n} \quad importance(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$




# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{= f_X(x_i)}}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$




# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$




# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$




# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$



$$X \sim \mathcal{U}(a, b)$$




# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$




~~$X \sim \mathcal{U}(a, b)$~~

# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$


# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$


# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$

# Monte Carlo integration (importance sampling)


intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:

$$\int_{\mathbb{S}} g(x) dx$$



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$

# Monte Carlo integration (importance sampling)


intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:

$$\int_{\mathbb{S}} g(x) dx = \int_{\mathbb{S}} g(x) 1 dx$$



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$

# Monte Carlo integration (importance sampling)


intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:

$$\int_{\mathbb{S}} g(x) dx = \int_{\mathbb{S}} g(x) 1 dx = \int_{\mathbb{S}} g(x) \frac{f_X(x)}{f_X(x)} dx$$



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$

# Monte Carlo integration (importance sampling)


intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:

$$\int_{\mathbb{S}} g(x) dx = \int_{\mathbb{S}} g(x) 1 dx = \int_{\mathbb{S}} g(x) \frac{f_X(x)}{f_X(x)} dx = \int_{\mathbb{S}} \frac{g(x)}{f_X(x)} f_X(x) dx$$



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$



# Monte Carlo integration (importance sampling)


intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:

$$\int_{\mathbb{S}} g(x) dx = \int_{\mathbb{S}} g(x) 1 dx = \int_{\mathbb{S}} g(x) \frac{f_X(x)}{f_X(x)} dx = \int_{\mathbb{S}} \frac{g(x)}{f_X(x)} f_X(x) dx = \mathbb{E} \left[ \frac{g(X)}{f_X(X)} \right]$$



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$

# Monte Carlo integration (importance sampling)

intuition:


$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:

$$\int_{\mathbb{S}} g(x) dx = \int_{\mathbb{S}} g(x) 1 dx = \int_{\mathbb{S}} g(x) \frac{f_X(x)}{f_X(x)} dx = \int_{\mathbb{S}} \frac{g(x)}{f_X(x)} f_X(x) dx = \mathbb{E} \left[ \frac{g(X)}{f_X(X)} \right]$$

=  
almost  
always



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$

# Monte Carlo integration (importance sampling)

intuition:


$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:

$$\int_{\mathbb{S}} g(x) dx = \int_{\mathbb{S}} g(x) 1 dx = \int_{\mathbb{S}} g(x) \frac{f_X(x)}{f_X(x)} dx = \int_{\mathbb{S}} \frac{g(x)}{f_X(x)} f_X(x) dx = \mathbb{E} \left[ \frac{g(X)}{f_X(X)} \right]$$

$$\stackrel{\text{almost always}}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$

# Monte Carlo integration (importance sampling)


intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:

$$\begin{aligned} \int_{\mathbb{S}} g(x) dx &= \int_{\mathbb{S}} g(x) 1 dx = \int_{\mathbb{S}} g(x) \frac{f_X(x)}{f_X(x)} dx = \int_{\mathbb{S}} \frac{g(x)}{f_X(x)} f_X(x) dx = \mathbb{E} \left[ \frac{g(X)}{f_X(X)} \right] \\ &\stackrel{\text{almost always}}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)} \end{aligned}$$



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$

# Monte Carlo integration (importance sampling)

intuition:


$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:

$$\begin{aligned} \int_{\mathbb{S}} g(x) dx &= \int_{\mathbb{S}} g(x) 1 dx = \int_{\mathbb{S}} g(x) \frac{f_X(x)}{f_X(x)} dx = \int_{\mathbb{S}} \frac{g(x)}{f_X(x)} f_X(x) dx = \mathbb{E} \left[ \frac{g(X)}{f_X(X)} \right] \\ &\stackrel{\text{almost always}}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)} \end{aligned}$$

$error_{\vartheta_n}$



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$

# Monte Carlo integration (importance sampling)

intuition:


$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:

$$\begin{aligned} \int_{\mathbb{S}} g(x) dx &= \int_{\mathbb{S}} g(x) 1 dx = \int_{\mathbb{S}} g(x) \frac{f_X(x)}{f_X(x)} dx = \int_{\mathbb{S}} \frac{g(x)}{f_X(x)} f_X(x) dx = \mathbb{E} \left[ \frac{g(X)}{f_X(X)} \right] \\ &\stackrel{\text{almost always}}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)} \end{aligned}$$

$$\text{error}_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{\text{Var} \left( \frac{g(X)}{f_X(X)} \right)}$$



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$



# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$


$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:

$$\begin{aligned} \int_{\mathbb{S}} g(x) dx &= \int_{\mathbb{S}} g(x) 1 dx = \int_{\mathbb{S}} g(x) \frac{f_X(x)}{f_X(x)} dx = \int_{\mathbb{S}} \frac{g(x)}{f_X(x)} f_X(x) dx = \mathbb{E} \left[ \frac{g(X)}{f_X(X)} \right] \\ &\stackrel{\text{almost always}}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)} \end{aligned}$$

$$\text{error}_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{\text{Var} \left( \frac{g(X)}{f_X(X)} \right)}$$

choice of  $f_X(x)$ :



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$

# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:


$$\int_{\mathbb{S}} g(x) dx = \int_{\mathbb{S}} g(x) 1 dx = \int_{\mathbb{S}} g(x) \frac{f_X(x)}{f_X(x)} dx = \int_{\mathbb{S}} \frac{g(x)}{f_X(x)} f_X(x) dx = \mathbb{E} \left[ \frac{g(X)}{f_X(X)} \right]$$

$$\stackrel{\text{almost always}}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\text{error}_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{\text{Var} \left( \frac{g(X)}{f_X(X)} \right)}$$

choice of  $f_X(x)$ :

$$f_X(x) = g(x)$$



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$



# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:


$$\int_{\mathbb{S}} g(x) dx = \int_{\mathbb{S}} g(x) 1 dx = \int_{\mathbb{S}} g(x) \frac{f_X(x)}{f_X(x)} dx = \int_{\mathbb{S}} \frac{g(x)}{f_X(x)} f_X(x) dx = \mathbb{E} \left[ \frac{g(X)}{f_X(X)} \right]$$

$$\stackrel{\text{almost always}}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\text{error}_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{\text{Var} \left( \frac{g(X)}{f_X(X)} \right)}$$

choice of  $f_X(x)$ :

$$f_X(x) = g(x) \cdot k$$



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$

# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:


$$\int_{\mathbb{S}} g(x) dx = \int_{\mathbb{S}} g(x) 1 dx = \int_{\mathbb{S}} g(x) \frac{f_X(x)}{f_X(x)} dx = \int_{\mathbb{S}} \frac{g(x)}{f_X(x)} f_X(x) dx = \mathbb{E} \left[ \frac{g(X)}{f_X(X)} \right]$$

$$\stackrel{\substack{= \\ \text{almost} \\ \text{always}}}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\text{error}_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{\text{Var} \left( \frac{g(X)}{f_X(X)} \right)}$$

choice of  $f_X(x)$ :

$$f_X(x) = g(x) \cdot k \quad 1 = \int_{\mathbb{S}} f_X(x) dx$$



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$

# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:


$$\int_{\mathbb{S}} g(x) dx = \int_{\mathbb{S}} g(x) 1 dx = \int_{\mathbb{S}} g(x) \frac{f_X(x)}{f_X(x)} dx = \int_{\mathbb{S}} \frac{g(x)}{f_X(x)} f_X(x) dx = \mathbb{E} \left[ \frac{g(X)}{f_X(X)} \right]$$

$$\stackrel{\substack{= \\ \text{almost} \\ \text{always}}}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\text{error}_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{\text{Var} \left( \frac{g(X)}{f_X(X)} \right)}$$

choice of  $f_X(x)$ :

$$f_X(x) = g(x) \cdot k \quad 1 = \int_{\mathbb{S}} f_X(x) dx = \int_{\mathbb{S}} g(x) \cdot k dx$$



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$

# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:


$$\int_{\mathbb{S}} g(x) dx = \int_{\mathbb{S}} g(x) 1 dx = \int_{\mathbb{S}} g(x) \frac{f_X(x)}{f_X(x)} dx = \int_{\mathbb{S}} \frac{g(x)}{f_X(x)} f_X(x) dx = \mathbb{E} \left[ \frac{g(X)}{f_X(X)} \right]$$

$$\stackrel{\substack{= \\ \text{almost} \\ \text{always}}}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\text{error}_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{\text{Var} \left( \frac{g(X)}{f_X(X)} \right)}$$

choice of  $f_X(x)$ :

$$f_X(x) = g(x) \cdot k \quad 1 = \int_{\mathbb{S}} f_X(x) dx = \int_{\mathbb{S}} g(x) \cdot k dx \quad \frac{1}{k} = \int_{\mathbb{S}} g(x) dx$$



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$

# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:


$$\int_{\mathbb{S}} g(x) dx = \int_{\mathbb{S}} g(x) 1 dx = \int_{\mathbb{S}} g(x) \frac{f_X(x)}{f_X(x)} dx = \int_{\mathbb{S}} \frac{g(x)}{f_X(x)} f_X(x) dx = \mathbb{E} \left[ \frac{g(X)}{f_X(X)} \right]$$

$$\stackrel{\substack{= \\ \text{almost} \\ \text{always}}}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\text{error}_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{\text{Var} \left( \frac{g(X)}{f_X(X)} \right)}$$

choice of  $f_X(x)$ :

$$f_X(x) = g(x) \cdot k \quad 1 = \int_{\mathbb{S}} f_X(x) dx = \int_{\mathbb{S}} g(x) \cdot k dx \quad \frac{1}{k} = \int_{\mathbb{S}} g(x) dx \quad \rightarrow$$



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$



# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:


$$\int_{\mathbb{S}} g(x) dx = \int_{\mathbb{S}} g(x) 1 dx = \int_{\mathbb{S}} g(x) \frac{f_X(x)}{f_X(x)} dx = \int_{\mathbb{S}} \frac{g(x)}{f_X(x)} f_X(x) dx = \mathbb{E} \left[ \frac{g(X)}{f_X(X)} \right]$$

$$\stackrel{\substack{= \\ \text{almost} \\ \text{always}}}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\text{error}_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{\text{Var} \left( \frac{g(X)}{f_X(X)} \right)}$$

choice of  $f_X(x)$ :

$$f_X(x) = g(x) \cdot k \quad 1 = \int_{\mathbb{S}} f_X(x) dx = \int_{\mathbb{S}} g(x) \cdot k dx \quad \frac{1}{k} = \int_{\mathbb{S}} g(x) dx \quad \longrightarrow \quad f_X(x)$$



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$

# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:


$$\int_{\mathbb{S}} g(x) dx = \int_{\mathbb{S}} g(x) 1 dx = \int_{\mathbb{S}} g(x) \frac{f_X(x)}{f_X(x)} dx = \int_{\mathbb{S}} \frac{g(x)}{f_X(x)} f_X(x) dx = \mathbb{E} \left[ \frac{g(X)}{f_X(X)} \right]$$

$$\stackrel{\substack{= \\ \text{almost} \\ \text{always}}}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\text{error}_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{\text{Var} \left( \frac{g(X)}{f_X(X)} \right)}$$

choice of  $f_X(x)$ :

$$f_X(x) = g(x) \cdot k \quad 1 = \int_{\mathbb{S}} f_X(x) dx = \int_{\mathbb{S}} g(x) \cdot k dx \quad \frac{1}{k} = \int_{\mathbb{S}} g(x) dx \quad \longrightarrow \quad f_X(x) \text{ similar to}$$



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$

# Monte Carlo integration (importance sampling)

intuition:

$$\int_a^b g(x) dx \approx \sum_{i=0}^{n-1} g(x_i) \Delta x_i = \sum_{i=0}^{n-1} g(x_i) \frac{1}{n \cdot \underbrace{\frac{b-a}{\Delta x_i}}_{=f_X(x_i)}} = \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(x_i)}{f_X(x_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\Delta x = \frac{b-a}{n} \quad \text{importance}(x_i) := \frac{\Delta x}{\Delta x_i} = \frac{b-a}{n \cdot \Delta x_i}$$

derivation:


$$\int_{\mathbb{S}} g(x) dx = \int_{\mathbb{S}} g(x) 1 dx = \int_{\mathbb{S}} g(x) \frac{f_X(x)}{f_X(x)} dx = \int_{\mathbb{S}} \frac{g(x)}{f_X(x)} f_X(x) dx = \mathbb{E} \left[ \frac{g(X)}{f_X(X)} \right]$$

$$\stackrel{\substack{= \\ \text{almost} \\ \text{always}}}{=} \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)} \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\text{error}_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{\text{Var} \left( \frac{g(X)}{f_X(X)} \right)}$$

choice of  $f_X(x)$ :

$$f_X(x) = g(x) \cdot k \quad 1 = \int_{\mathbb{S}} f_X(x) dx = \int_{\mathbb{S}} g(x) \cdot k dx \quad \frac{1}{k} = \int_{\mathbb{S}} g(x) dx \quad \longrightarrow \quad f_X(x) \text{ similar to } g(x)$$



~~$X \sim \mathcal{U}(a, b)$~~

$X$  distributed with PDF  $f_X$

expectation

$$\mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

SLLN

$$P \left( \lim_{n \rightarrow \infty} \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) = \mathbb{E}[g(X)] \right) = 1$$



# Monte Carlo integration (example)

# Monte Carlo integration (example)

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$\text{Var}(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$\text{error}_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{\text{Var}(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\text{error}_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{\text{Var}\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$\text{Var}(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$\text{error}_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{\text{Var}(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\text{error}_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{\text{Var}\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$\text{Var}(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$\text{error}_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{\text{Var}(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\text{error}_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{\text{Var}\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$\text{Var}(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$\text{error}_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{\text{Var}(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\text{error}_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{\text{Var}\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$\text{Var}(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$\text{error}_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{\text{Var}(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\text{error}_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{\text{Var}\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$\text{Var}(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$\text{error}_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{\text{Var}(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$\text{error}_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{\text{Var}\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)}$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$



# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2}$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2}$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5}$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2}$$
$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2]$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$



# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}} \approx \frac{0.298}{\sqrt{n}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}} \approx \frac{0.298}{\sqrt{n}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

Monte Carlo integration with importance sampling

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}} \approx \frac{0.298}{\sqrt{n}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

Monte Carlo integration with importance sampling

$$\int_0^1 x^2 dx$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}} \approx \frac{0.298}{\sqrt{n}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

Monte Carlo integration with importance sampling

$$\int_0^1 x^2 dx \quad X_i \text{ is distributed with PDF } f_X(x) := 2x$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}} \approx \frac{0.298}{\sqrt{n}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

Monte Carlo integration with importance sampling

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{X_i^2}{2X_i} \quad X_i \text{ is distributed with PDF } f_X(x) := 2x$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}} \approx \frac{0.298}{\sqrt{n}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

Monte Carlo integration with importance sampling

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{X_i^2}{2X_i} = \vartheta'_n \quad X_i \text{ is distributed with PDF } f_X(x) := 2x$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}} \approx \frac{0.298}{\sqrt{n}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

Monte Carlo integration with importance sampling

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{X_i^2}{2X_i} = \vartheta'_n \quad X_i \text{ is distributed with PDF } f_X(x) := 2x$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{X^2}{2X}\right)}$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$



# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}} \approx \frac{0.298}{\sqrt{n}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

Monte Carlo integration with importance sampling

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{X_i^2}{2X_i} = \vartheta'_n \quad X_i \text{ is distributed with PDF } f_X(x) := 2x$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{X^2}{2X}\right)} = \frac{1}{2\sqrt{n}} \sqrt{Var(X)}$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$



# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}} \approx \frac{0.298}{\sqrt{n}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

Monte Carlo integration with importance sampling

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{X_i^2}{2X_i} = \vartheta'_n \quad X_i \text{ is distributed with PDF } f_X(x) := 2x$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{X^2}{2X}\right)} = \frac{1}{2\sqrt{n}} \sqrt{Var(X)} = \frac{1}{2\sqrt{n}} \sqrt{\mathbb{E}[X^2] - \mathbb{E}[X]^2}$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}} \approx \frac{0.298}{\sqrt{n}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

Monte Carlo integration with importance sampling

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{X_i^2}{2X_i} = \vartheta'_n \quad X_i \text{ is distributed with PDF } f_X(x) := 2x$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{X^2}{2X}\right)} = \frac{1}{2\sqrt{n}} \sqrt{Var(X)} = \frac{1}{2\sqrt{n}} \sqrt{\mathbb{E}[X^2] - \mathbb{E}[X]^2}$$

$$\mathbb{E}[X^2]$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}} \approx \frac{0.298}{\sqrt{n}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

Monte Carlo integration with importance sampling

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{X_i^2}{2X_i} = \vartheta'_n \quad X_i \text{ is distributed with PDF } f_X(x) := 2x$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{X^2}{2X}\right)} = \frac{1}{2\sqrt{n}} \sqrt{Var(X)} = \frac{1}{2\sqrt{n}} \sqrt{\mathbb{E}[X^2] - \mathbb{E}[X]^2}$$

$$\mathbb{E}[X^2] = \int_0^1 x^2 f_X(x) dx$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}} \approx \frac{0.298}{\sqrt{n}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

Monte Carlo integration with importance sampling

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{X_i^2}{2X_i} = \vartheta'_n \quad X_i \text{ is distributed with PDF } f_X(x) := 2x$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{X^2}{2X}\right)} = \frac{1}{2\sqrt{n}} \sqrt{Var(X)} = \frac{1}{2\sqrt{n}} \sqrt{\mathbb{E}[X^2] - \mathbb{E}[X]^2}$$

$$\mathbb{E}[X^2] = \int_0^1 x^2 f_X(x) dx = \int_0^1 2x^3 dx$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}} \approx \frac{0.298}{\sqrt{n}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

Monte Carlo integration with importance sampling

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{X_i^2}{2X_i} = \vartheta'_n \quad X_i \text{ is distributed with PDF } f_X(x) := 2x$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{X^2}{2X}\right)} = \frac{1}{2\sqrt{n}} \sqrt{Var(X)} = \frac{1}{2\sqrt{n}} \sqrt{\mathbb{E}[X^2] - \mathbb{E}[X]^2}$$

$$\mathbb{E}[X^2] = \int_0^1 x^2 f_X(x) dx = \int_0^1 2x^3 dx = \frac{1}{2}$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}} \approx \frac{0.298}{\sqrt{n}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

Monte Carlo integration with importance sampling

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{X_i^2}{2X_i} = \vartheta'_n \quad X_i \text{ is distributed with PDF } f_X(x) := 2x$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{X^2}{2X}\right)} = \frac{1}{2\sqrt{n}} \sqrt{Var(X)} = \frac{1}{2\sqrt{n}} \sqrt{\mathbb{E}[X^2] - \mathbb{E}[X]^2}$$

$$\mathbb{E}[X^2] = \int_0^1 x^2 f_X(x) dx = \int_0^1 2x^3 dx = \frac{1}{2} \quad \mathbb{E}[X]$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$



# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}} \approx \frac{0.298}{\sqrt{n}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

Monte Carlo integration with importance sampling

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{X_i^2}{2X_i} = \vartheta'_n \quad X_i \text{ is distributed with PDF } f_X(x) := 2x$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{X^2}{2X}\right)} = \frac{1}{2\sqrt{n}} \sqrt{Var(X)} = \frac{1}{2\sqrt{n}} \sqrt{\mathbb{E}[X^2] - \mathbb{E}[X]^2}$$

$$\mathbb{E}[X^2] = \int_0^1 x^2 f_X(x) dx = \int_0^1 2x^3 dx = \frac{1}{2} \quad \mathbb{E}[X] = \int_0^1 x f_X(x) dx$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}} \approx \frac{0.298}{\sqrt{n}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

Monte Carlo integration with importance sampling

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{X_i^2}{2X_i} = \vartheta'_n \quad X_i \text{ is distributed with PDF } f_X(x) := 2x$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{X^2}{2X}\right)} = \frac{1}{2\sqrt{n}} \sqrt{Var(X)} = \frac{1}{2\sqrt{n}} \sqrt{\mathbb{E}[X^2] - \mathbb{E}[X]^2}$$

$$\mathbb{E}[X^2] = \int_0^1 x^2 f_X(x) dx = \int_0^1 2x^3 dx = \frac{1}{2} \quad \mathbb{E}[X] = \int_0^1 x f_X(x) dx = \int_0^1 2x^2 dx$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$



# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}} \approx \frac{0.298}{\sqrt{n}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

Monte Carlo integration with importance sampling

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{X_i^2}{2X_i} = \vartheta'_n \quad X_i \text{ is distributed with PDF } f_X(x) := 2x$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{X^2}{2X}\right)} = \frac{1}{2\sqrt{n}} \sqrt{Var(X)} = \frac{1}{2\sqrt{n}} \sqrt{\mathbb{E}[X^2] - \mathbb{E}[X]^2}$$

$$\mathbb{E}[X^2] = \int_0^1 x^2 f_X(x) dx = \int_0^1 2x^3 dx = \frac{1}{2} \quad \mathbb{E}[X] = \int_0^1 x f_X(x) dx = \int_0^1 2x^2 dx = \frac{2}{3}$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}} \approx \frac{0.298}{\sqrt{n}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

Monte Carlo integration with importance sampling

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{X_i^2}{2X_i} = \vartheta'_n \quad X_i \text{ is distributed with PDF } f_X(x) := 2x$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{X^2}{2X}\right)} = \frac{1}{2\sqrt{n}} \sqrt{Var(X)} = \frac{1}{2\sqrt{n}} \sqrt{\mathbb{E}[X^2] - \mathbb{E}[X]^2} = \frac{1}{2\sqrt{n}} \sqrt{\frac{1}{2} - \frac{4}{9}}$$

$$\mathbb{E}[X^2] = \int_0^1 x^2 f_X(x) dx = \int_0^1 2x^3 dx = \frac{1}{2} \quad \mathbb{E}[X] = \int_0^1 x f_X(x) dx = \int_0^1 2x^2 dx = \frac{2}{3}$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

# Monte Carlo integration (example)

Monte Carlo integration

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} X_i^2 = \vartheta'_n \quad X_i \sim \mathcal{U}(0,1)$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var(X^2)} = \frac{1}{\sqrt{n}} \sqrt{\mathbb{E}[X^4] - \mathbb{E}[X^2]^2} = \frac{1}{\sqrt{n}} \sqrt{\frac{1}{5} - \frac{1}{9}} \approx \frac{0.298}{\sqrt{n}}$$

$$\mathbb{E}[X^4] = \int_0^1 x^4 dx = \frac{1}{5} \quad \mathbb{E}[X^2] = \int_0^1 x^2 dx = \frac{1}{3}$$

Monte Carlo integration with importance sampling

$$\int_0^1 x^2 dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{X_i^2}{2X_i} = \vartheta'_n \quad X_i \text{ is distributed with PDF } f_X(x) := 2x$$

$$error_{\vartheta'_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{X^2}{2X}\right)} = \frac{1}{2\sqrt{n}} \sqrt{Var(X)} = \frac{1}{2\sqrt{n}} \sqrt{\mathbb{E}[X^2] - \mathbb{E}[X]^2} = \frac{1}{2\sqrt{n}} \sqrt{\frac{1}{2} - \frac{4}{9}} \approx \frac{0.118}{\sqrt{n}}$$

$$\mathbb{E}[X^2] = \int_0^1 x^2 f_X(x) dx = \int_0^1 2x^3 dx = \frac{1}{2} \quad \mathbb{E}[X] = \int_0^1 x f_X(x) dx = \int_0^1 2x^2 dx = \frac{2}{3}$$

moments

$$\mu := \mathbb{E}[g(X)] := \int_{\mathbb{S}} g(x) f_X(x) dx$$

$$Var(X) = \mathbb{E}[X^2] - \mu^2$$

MC integration

$$\int_a^b g(x) dx \approx (b-a) \frac{1}{n} \sum_{i=0}^{n-1} g(X_i) =: \vartheta_n$$

$$error_{\vartheta_n} \approx \frac{b-a}{\sqrt{n}} \sqrt{Var(g(X))}$$

MC integration importance sampling

$$\int_{\mathbb{S}} g(x) dx \approx \frac{1}{n} \sum_{i=0}^{n-1} \frac{g(X_i)}{f_X(X_i)}$$

$$error_{\vartheta_n} \approx \frac{1}{\sqrt{n}} \sqrt{Var\left(\frac{g(X)}{f_X(X)}\right)}$$

end