

stochastics and probability

Lecture 1

Dr. Johannes Pahlke

indroduction

indroduction

x , t

indroduction

$$\underline{x}, t \mid \underline{x}_0, t_0$$

indroduction

$$P(\underline{x}, t | \underline{x}_0, t_0)$$

indroduction

stochastic model

$$P(\underline{x}, t | \underline{x}_0, t_0)$$

indroduction

stochastic model

$$P(\underline{x}, t | \underline{x}_0, t_0)$$

$$\underline{x}_0, \underline{x}_1, \underline{x}_2, \dots, \underline{x}_T$$

indroduction

stochastic model

$$P(\underline{x}, t | \underline{x}_0, t_0)$$

stochastic simulation

$$\underline{x}_0, \underline{x}_1, \underline{x}_2, \dots, \underline{x}_T$$

indroduction

stochastic model

$$P(\underline{x}, t \mid \underline{x}_0, t_0)$$

stochastic simulation

$$\underline{x}_0, \underline{x}_1, \underline{x}_2, \dots, \underline{x}_T$$

$$\underline{x}_t \sim P(\underline{x}, t \mid \underline{x}_0, t_0)$$

indroduction

stochastic model

$$P(\underline{x}, t \mid \underline{x}_0, t_0)$$

stochastic simulation

$$\underline{x}_0, \underline{x}_1, \underline{x}_2, \dots, \underline{x}_T$$

$$\forall t = 0, \dots, T : \quad \underline{x}_t \sim P(\underline{x}, t \mid \underline{x}_0, t_0)$$

problems

problems

1) $P(\cdot)$ not known in closed form

problems

- 1) $P(\cdot)$ not known in closed form
→ Master Equations

problems

1) $P(\cdot)$ not known in closed form

→ Master Equations

2) generating random numbers $\sim P(\cdot)$ is not easy/ possible

problems

1) $P(.)$ not known in closed form

→ Master Equations

2) generating random numbers $\sim P(.)$ is not easy/ possible

→ sampling

problems

1) $P(.)$ not known in closed form

→ Master Equations

2) generating random numbers $\sim P(.)$ is not easy/ possible

→ sampling

3) time is continuous $\underline{x}(t)$

problems

1) $P(\cdot)$ not known in closed form

→ Master Equations

2) generating random numbers $\sim P(\cdot)$ is not easy/ possible

→ sampling

3) time is continuous $\underline{x}(t)$

→ stochastic calculus

problems

- 1) $P(.)$ not known in closed form
→ Master Equations
- 2) generating random numbers $\sim P(.)$ is not easy/ possible
→ sampling
- 3) time is continuous $\underline{x}(t)$
→ stochastic calculus
- 4) $P(.)$ changes as a function of history

problems

- 1) $P(.)$ not known in closed form
→ Master Equations
- 2) generating random numbers $\sim P(.)$ is not easy/ possible
→ sampling
- 3) time is continuous $\underline{x}(t)$
→ stochastic calculus
- 4) $P(.)$ changes as a function of history
→ non-Markovian processes

elementary probability

elementary probability

definitions:

elementary probability

definitions:

population:

elementary probability

definitions:

population: a collection of objects

elementary probability

definitions:

population: a collection of objects

sample:

elementary probability

definitions:

population: a collection of objects

sample: a subset of a population

elementary probability

definitions:

population: a collection of objects

sample: a subset of a population

experiment:

elementary probability

definitions:

population: a collection of objects

sample: a subset of a population

experiment: measuring something of a sample

elementary probability

definitions:

population: a collection of objects

sample: a subset of a population

experiment: measuring something of a sample

event:

elementary probability

definitions:

population: a collection of objects

sample: a subset of a population

experiment: measuring something of a sample

event: a concrete set of outcomes of an experiment

elementary probability

definitions:

population: a collection of objects

sample: a subset of a population

experiment: measuring something of a sample

event: a concrete set of outcomes of an experiment

sample space:

elementary probability

definitions:

population: a collection of objects

sample: a subset of a population

experiment: measuring something of a sample

event: a concrete set of outcomes of an experiment

sample space: space of all possible events Ω

elementary probability

definitions:

population: a collection of objects

sample: a subset of a population

experiment: measuring something of a sample

event: a concrete set of outcomes of an experiment

sample space: space of all possible events Ω

probability:

elementary probability

definitions:

population: a collection of objects

sample: a subset of a population

experiment: measuring something of a sample

event: a concrete set of outcomes of an experiment

sample space: space of all possible events Ω

probability: likelihood of an event in a sample space

$P(A)$ for the event $A \subseteq \Omega$

example elementary probability

Population: a collection of objects

Sample: a subset of a population


Experiment: measuring something of a sample

Event: a concrete set of outcomes of an experiment

Sample space: space of all possible events Ω

Probability: likelihood of an event in a sample space
 $P(A)$ for the event $A \subseteq \Omega$

example elementary probability



Population: a collection of objects

Sample: a subset of a population

Experiment: measuring something of a sample


Event: a concrete set of outcomes of an experiment

Sample space: space of all possible events Ω

Probability: likelihood of an event in a sample space
 $P(A)$ for the event $A \subseteq \Omega$

example elementary probability

population



Population: a collection of objects

Sample: a subset of a population

Experiment: measuring something of a sample

Event: a concrete set of outcomes of an experiment



Sample space: space of all possible events Ω

Probability: likelihood of an event in a sample space
 $P(A)$ for the event $A \subseteq \Omega$

example elementary probability

population

sample



Population: a collection of objects

Sample: a subset of a population


Experiment: measuring something of a sample

Event: a concrete set of outcomes of an experiment

Sample space: space of all possible events Ω

Probability: likelihood of an event in a sample space
 $P(A)$ for the event $A \subseteq \Omega$

example elementary probability



Population: a collection of objects

Sample: a subset of a population


Experiment: measuring something of a sample

Event: a concrete set of outcomes of an experiment

Sample space: space of all possible events Ω

Probability: likelihood of an event in a sample space
 $P(A)$ for the event $A \subseteq \Omega$

example elementary probability




event

140
190
175
130

- Population: a collection of objects
- Sample: a subset of a population
- Experiment: measuring something of a sample
- Event: a concrete set of outcomes of an experiment
- Sample space: space of all possible events Ω
- Probability: likelihood of an event in a sample space $P(A)$ for the event $A \subseteq \Omega$

example elementary probability




- Population: a collection of objects
- Sample: a subset of a population
- Experiment: measuring something of a sample
- Event: a concrete set of outcomes of an experiment
- Sample space: space of all possible events Ω
- Probability: likelihood of an event in a sample space $P(A)$ for the event $A \subseteq \Omega$

event
140
190
175
130

sample space

$$\Omega = [0,300]$$

example elementary probability



event

140

190

175

130

Population: a collection of objects

Sample: a subset of a population

Experiment: measuring something of a sample

Event: a concrete set of outcomes of an experiment

Sample space: space of all possible events Ω

Probability: likelihood of an event in a sample space
 $P(A)$ for the event $A \subseteq \Omega$

sample space

$$\Omega = [0, 300]$$

probability

$$P(X \in [50, 200]) = 0.98$$

Kolmogorov axioms

Kolmogorov axioms

$$1) P(A) \geq 0 \quad \forall A \subseteq \Omega$$

Kolmogorov axioms

$$1) P(A) \geq 0 \quad \forall A \subseteq \Omega$$

$$2) P(\Omega) = 1$$

Kolmogorov axioms

$$1) P(A) \geq 0 \quad \forall A \subseteq \Omega$$

$$2) P(\Omega) = 1$$

$$3) P(A_1 \cup A_2 \cup \dots) = \sum P(A_i) \quad \text{if } A_i \cap A_j = \emptyset \text{ for all } i \neq j$$

implications

implications

$$P(\emptyset) = 0$$

implications

$$P(\emptyset) = 0$$

$$P(\overline{A}) = 1 - P(A)$$

implications

$$P(\emptyset) = 0$$

$$P(\overline{A}) = 1 - P(A)$$

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

exercice

Prove: $P(\bar{A}) = 1 - P(A)$

Axioms:

$$\forall A : P(A) \geq 0$$

$$P(\Omega) = 1$$

$$P(A \cup B) = P(A) + P(B) \quad \text{if } P(A \cap B) = 0$$

$$\bar{A} = A^c = A' = \neg A := \Omega \setminus A = \Omega - A$$

exercice

Prove: $P(\bar{A}) = 1 - P(A)$

Axioms:

$$\forall A : P(A) \geq 0$$

$$P(\Omega) = 1$$

$$P(A \cup B) = P(A) + P(B) \quad \text{if } P(A \cap B) = 0$$

$$\bar{A} = A^c = A' = \neg A := \Omega \setminus A = \Omega - A$$

Ω



exercice

Prove: $P(\bar{A}) = 1 - P(A)$

$$1 = P(\Omega)$$

Axioms:

$$\forall A : P(A) \geq 0$$

$$P(\Omega) = 1$$

$$P(A \cup B) = P(A) + P(B) \quad \text{if } P(A \cap B) = 0$$

$$\bar{A} = A^c = A' = \neg A := \Omega \setminus A = \Omega - A$$

Ω



exercice

Prove: $P(\bar{A}) = 1 - P(A)$

$$1 = P(\Omega)$$

Axioms:

$$\forall A : P(A) \geq 0$$

$$P(\Omega) = 1$$

$$P(A \cup B) = P(A) + P(B) \quad \text{if } P(A \cap B) = 0$$

$$\bar{A} = A^c = A' = \neg A := \Omega \setminus A = \Omega - A$$



exercice

Prove: $P(\bar{A}) = 1 - P(A)$

$$\begin{aligned} 1 &= P(\Omega) \\ &= P(\Omega \setminus A \cup A) \end{aligned}$$

Axioms:

$$\forall A : P(A) \geq 0$$

$$P(\Omega) = 1$$

$$P(A \cup B) = P(A) + P(B) \quad \text{if } P(A \cap B) = 0$$

$$\bar{A} = A^c = A' = \neg A := \Omega \setminus A = \Omega - A$$



exercice

Prove: $P(\bar{A}) = 1 - P(A)$

$$\begin{aligned} 1 &= P(\Omega) \\ &= P(\Omega \setminus A \cup A) \end{aligned}$$

Axioms:

$$\forall A : P(A) \geq 0$$

$$P(\Omega) = 1$$

$$P(A \cup B) = P(A) + P(B) \quad \text{if } P(A \cap B) = 0$$

$$\bar{A} = A^c = A' = \neg A := \Omega \setminus A = \Omega - A$$



exercice

Prove: $P(\bar{A}) = 1 - P(A)$

$$\begin{aligned} 1 &= P(\Omega) \\ &= P(\Omega \setminus A \cup A) \\ &= P(\bar{A} \cup A) \end{aligned}$$


Axioms:

$$\forall A : P(A) \geq 0$$

$$P(\Omega) = 1$$

$$P(A \cup B) = P(A) + P(B) \quad \text{if } P(A \cap B) = 0$$

$$\bar{A} = A^c = A' = \neg A := \Omega \setminus A = \Omega - A$$



exercice

Prove: $P(\bar{A}) = 1 - P(A)$

$$\begin{aligned} 1 &= P(\Omega) \\ &= P(\Omega \setminus A \cup A) \\ &= P(\bar{A} \cup A) \end{aligned}$$

Axioms:

$$\forall A : P(A) \geq 0$$

$$P(\Omega) = 1$$

$$P(A \cup B) = P(A) + P(B) \quad \text{if } P(A \cap B) = 0$$

$$\bar{A} = A^c = A' = \neg A := \Omega \setminus A = \Omega - A$$



$$\bar{A} \cap A = \emptyset$$

exercice

Prove: $P(\bar{A}) = 1 - P(A)$

$$\begin{aligned} 1 &= P(\Omega) \\ &= P(\Omega \setminus A \cup A) \\ &= P(\bar{A} \cup A) \\ &= P(\bar{A}) + P(A) \end{aligned}$$

Axioms:

$$\forall A : P(A) \geq 0$$

$$P(\Omega) = 1$$

$$P(A \cup B) = P(A) + P(B) \quad \text{if } P(A \cap B) = 0$$

$$\bar{A} = A^c = A' = \neg A := \Omega \setminus A = \Omega - A$$



$$\bar{A} \cap A = \emptyset$$

exercice

Prove: $P(\bar{A}) = 1 - P(A)$

$$\begin{aligned} 1 &= P(\Omega) \\ &= P(\Omega \setminus A \cup A) \\ &= P(\bar{A} \cup A) \\ &= P(\bar{A}) + P(A) \\ \rightarrow P(\bar{A}) &= 1 - P(A) \end{aligned}$$

Axioms:

$$\forall A : P(A) \geq 0$$

$$P(\Omega) = 1$$

$$P(A \cup B) = P(A) + P(B) \quad \text{if } P(A \cap B) = 0$$

$$\bar{A} = A^c = A' = \neg A := \Omega \setminus A = \Omega - A$$



$$\bar{A} \cap A = \emptyset$$

exercice

Prove: $P(\bar{A}) = 1 - P(A)$

$$\begin{aligned} 1 &= P(\Omega) \\ &= P(\Omega \setminus A \cup A) \\ &= P(\bar{A} \cup A) \\ &= P(\bar{A}) + P(A) \\ \rightarrow P(\bar{A}) &= 1 - P(A) \quad \square \end{aligned}$$


Axioms:

$$\forall A : P(A) \geq 0$$

$$P(\Omega) = 1$$

$$P(A \cup B) = P(A) + P(B) \quad \text{if } P(A \cap B) = 0$$

$$\bar{A} = A^c = A' = \neg A := \Omega \setminus A = \Omega - A$$



$$\bar{A} \cap A = \emptyset$$

conditional probabilities

conditional probabilities

definitions:

conditional probabilities

definitions:

conditional probabilities

conditional probabilities

definitions:

conditional probabilities

$$P(A \mid B) := \frac{P(A \cap B)}{P(B)}$$

conditional probabilities

definitions:

conditional probabilities

$$P(A \mid B) := \frac{P(A \cap B)}{P(B)}$$

"A given B"


example conditional probabilities

$$P(A | B) := \frac{P(A \cap B)}{P(B)}$$

$$P(A \cup B) = P(A) + P(B) \\ \text{if } P(A \cap B) = 0$$

example conditional probabilities

draw




$$P(A | B) := \frac{P(A \cap B)}{P(B)}$$

$$P(A \cup B) = P(A) + P(B) \\ \text{if } P(A \cap B) = 0$$

example conditional probabilities

draw



$$P(\text{blue rabbit}) = \frac{1}{3}$$

$$P(\text{pink rabbit}) = \frac{1}{3}$$


$$P(\text{blue duck}) = \frac{1}{3}$$

$$P(A | B) := \frac{P(A \cap B)}{P(B)}$$

$$P(A \cup B) = P(A) + P(B) \\ \text{if } P(A \cap B) = 0$$

example conditional probabilities

draw



$$P(\text{blue rabbit}) = \frac{1}{3} \quad P(\text{pink rabbit}) = \frac{1}{3} \quad P(\text{blue duck}) = \frac{1}{3}$$


$$P(\text{grey rabbit} \mid \text{blue circle})$$

$$P(A \mid B) := \frac{P(A \cap B)}{P(B)}$$

$$P(A \cup B) = P(A) + P(B) \\ \text{if } P(A \cap B) = 0$$

example conditional probabilities

draw



$$P(\text{blue rabbit}) = \frac{1}{3}$$

$$P(\text{pink rabbit}) = \frac{1}{3}$$

$$P(\text{blue duck}) = \frac{1}{3}$$


$$P(\text{pink rabbit} \mid \text{blue circle}) = \frac{P(\text{pink rabbit} \cap \text{blue circle})}{P(\text{blue circle})}$$

$$P(A \mid B) := \frac{P(A \cap B)}{P(B)}$$

$$P(A \cup B) = P(A) + P(B) \\ \text{if } P(A \cap B) = 0$$

example conditional probabilities

draw



$$P(\text{blue rabbit}) = \frac{1}{3} \quad P(\text{pink rabbit}) = \frac{1}{3} \quad P(\text{blue duck}) = \frac{1}{3}$$


$$P(\text{pink rabbit} \mid \text{blue circle}) = \frac{P(\text{pink rabbit} \cap \text{blue circle})}{P(\text{blue circle})} = \frac{P(\text{blue rabbit})}{P(\text{blue rabbit} \cup \text{blue duck})}$$

$$P(A \mid B) := \frac{P(A \cap B)}{P(B)}$$

$$P(A \cup B) = P(A) + P(B) \\ \text{if } P(A \cap B) = 0$$

example conditional probabilities

draw



$$P(\text{blue rabbit}) = \frac{1}{3} \quad P(\text{pink rabbit}) = \frac{1}{3} \quad P(\text{blue duck}) = \frac{1}{3}$$


$$P(\text{blue rabbit} \mid \text{blue}) = \frac{P(\text{blue rabbit} \cap \text{blue})}{P(\text{blue})} = \frac{P(\text{blue rabbit})}{P(\text{blue rabbit} \cup \text{blue duck})} = \frac{P(\text{blue rabbit})}{P(\text{blue rabbit}) + P(\text{blue duck})}$$

$$P(A \mid B) := \frac{P(A \cap B)}{P(B)}$$

$$P(A \cup B) = P(A) + P(B) \\ \text{if } P(A \cap B) = 0$$

example conditional probabilities

draw



$$P(\text{blue rabbit}) = \frac{1}{3} \quad P(\text{pink rabbit}) = \frac{1}{3} \quad P(\text{blue duck}) = \frac{1}{3}$$


$$\begin{aligned} P(\text{blue rabbit} \mid \text{blue}) &= \frac{P(\text{blue rabbit} \cap \text{blue})}{P(\text{blue})} = \frac{P(\text{blue rabbit})}{P(\text{blue rabbit} \cup \text{blue duck})} = \frac{P(\text{blue rabbit})}{P(\text{blue rabbit}) + P(\text{blue duck})} \\ &= \frac{\frac{1}{3}}{\frac{1}{3} + \frac{1}{3}} \end{aligned}$$

$$P(A \mid B) := \frac{P(A \cap B)}{P(B)}$$

$$P(A \cup B) = P(A) + P(B) \\ \text{if } P(A \cap B) = 0$$

example conditional probabilities

draw



$$P(\text{blue rabbit}) = \frac{1}{3} \quad P(\text{pink rabbit}) = \frac{1}{3} \quad P(\text{blue duck}) = \frac{1}{3}$$

$$\begin{aligned} P(\text{blue rabbit} \mid \text{blue}) &= \frac{P(\text{blue rabbit} \cap \text{blue})}{P(\text{blue})} = \frac{P(\text{blue rabbit})}{P(\text{blue rabbit} \cup \text{blue duck})} = \frac{P(\text{blue rabbit})}{P(\text{blue rabbit}) + P(\text{blue duck})} \\ &= \frac{\frac{1}{3}}{\frac{1}{3} + \frac{1}{3}} = \frac{1}{2} \end{aligned}$$

$$P(A \mid B) := \frac{P(A \cap B)}{P(B)}$$

$$P(A \cup B) = P(A) + P(B) \\ \text{if } P(A \cap B) = 0$$

conditional probabilities

conditional probabilities

definitions:

conditional probabilities

definitions:


An event A is independent of event B iff $P(A | B) = P(A)$.

conditional probabilities

definitions:

An event A is independent of event B iff $P(A | B) = P(A)$.

draw and flip




conditional probabilities

definitions:

An event A is independent of event B iff $P(A | B) = P(A)$.

draw and flip




$$P(\text{head}) = \frac{1}{2} \quad P(\text{2\$}) = \frac{1}{2}$$

conditional probabilities

definitions:

An event A is independent of event B iff $P(A | B) = P(A)$.

draw and flip



$$P(\text{head}) = \frac{1}{2} \quad P(2\$) = \frac{1}{2}$$


$$P(\text{head} | \text{pink})$$

conditional probabilities

definitions:

An event A is independent of event B iff $P(A | B) = P(A)$.

draw and flip



$$P(\text{head}) = \frac{1}{2} \quad P(2\$) = \frac{1}{2}$$


$$P(\text{head} | \text{pink}) = \frac{1}{2}$$

conditional probabilities

definitions:

An event A is independent of event B iff $P(A | B) = P(A)$.

draw and flip



$$P(\text{head}) = \frac{1}{2} \quad P(2\$) = \frac{1}{2}$$

$$P(\text{head} | \text{pink}) = \frac{1}{2} = P(\text{head})$$


conditional probabilities

definitions:

An event A is independent of event B iff $P(A | B) = P(A)$.

$$\longrightarrow P(A \cap B) = P(A) \cdot P(B)$$

draw and flip



$$P(\text{head}) = \frac{1}{2} \quad P(2\$) = \frac{1}{2}$$

$$P(\text{head} | \text{pink}) = \frac{1}{2} = P(\text{head})$$

exercice

Prove: $P(A \cap B) = P(A) \cdot P(B)$
for A, B independent

conditional probabilities:

$$P(A | B) := \frac{P(A \cap B)}{P(B)}$$

A independent of B :

$$P(A | B) = P(A)$$

exercice

Prove: $P(A \cap B) = P(A) \cdot P(B)$
for A, B independent

$$P(A \cap B) = P(A | B) \cdot P(B)$$

conditional probabilities:

$$P(A | B) := \frac{P(A \cap B)}{P(B)}$$

A independent of B :

$$P(A | B) = P(A)$$

exercice

Prove: $P(A \cap B) = P(A) \cdot P(B)$
for A, B independent

$$\begin{aligned} P(A \cap B) &= P(A | B) \cdot P(B) \\ &= P(A) \cdot P(B) \end{aligned}$$

conditional probabilities:

$$P(A | B) := \frac{P(A \cap B)}{P(B)}$$

A independent of B :

$$P(A | B) = P(A)$$

exercice

Prove: $P(A \cap B) = P(A) \cdot P(B)$
for A, B independent

$$\begin{aligned} P(A \cap B) &= P(A | B) \cdot P(B) \\ &= P(A) \cdot P(B) \end{aligned}$$

□

conditional probabilities:

$$P(A | B) := \frac{P(A \cap B)}{P(B)}$$

A independent of B :

$$P(A | B) = P(A)$$


Bayes' theorem

Bayes' theorem

$$P(A | B) = \frac{P(B | A) \cdot P(A)}{P(B)}$$

Bayes' theorem


posterior




$$P(A | B) = \frac{P(B | A) \cdot P(A)}{P(B)}$$

Bayes' theorem

posterior



prior



$$P(A | B) = \frac{P(B | A) \cdot P(A)}{P(B)}$$

Bayes' theorem


posterior

prior

$$P(A | B) = \frac{P(B | A) \cdot P(A)}{P(B)}$$

marginal

Bayes' theorem



A diagram illustrating Bayes' theorem. The equation $P(A | B) = \frac{P(B | A) \cdot P(A)}{P(B)}$ is centered. Four labels with arrows point to parts of the equation: 'posterior' points to $P(A | B)$, 'likelihood' points to $P(B | A)$, 'prior' points to $P(A)$, and 'marginal' points to $P(B)$.

posterior

likelihood

prior

$$P(A | B) = \frac{P(B | A) \cdot P(A)}{P(B)}$$

marginal

exercice

prove Bayes' theorem:

$$P(A | B) = \frac{P(B | A) \cdot P(A)}{P(B)}$$

conditional probabilities:

$$P(A | B) := \frac{P(A \cap B)}{P(B)}$$

exercice

prove Bayes' theorem:

$$P(A | B) = \frac{P(B | A) \cdot P(A)}{P(B)}$$

$$P(A \cap B) = P(A | B) \cdot P(B)$$

conditional probabilities:

$$P(A | B) := \frac{P(A \cap B)}{P(B)}$$

exercice

prove Bayes' theorem:

$$P(A | B) = \frac{P(B | A) \cdot P(A)}{P(B)}$$

$$P(A \cap B) = P(A | B) \cdot P(B)$$

$$P(B \cap A) = P(B | A) \cdot P(A)$$

conditional probabilities:

$$P(A | B) := \frac{P(A \cap B)}{P(B)}$$

exercice

prove Bayes' theorem:

$$P(A | B) = \frac{P(B | A) \cdot P(A)}{P(B)}$$

$$P(A \cap B) = P(A | B) \cdot P(B)$$

$$P(B \cap A) = P(B | A) \cdot P(A)$$

$$P(A \cap B) = P(B \cap A)$$

conditional probabilities:

$$P(A | B) := \frac{P(A \cap B)}{P(B)}$$

exercice

prove Bayes' theorem:

$$P(A | B) = \frac{P(B | A) \cdot P(A)}{P(B)}$$

$$P(A \cap B) = P(A | B) \cdot P(B)$$

$$P(B \cap A) = P(B | A) \cdot P(A)$$

$$P(A \cap B) = P(B \cap A)$$

$$P(A | B) \cdot P(B) = P(B | A) \cdot P(A)$$

conditional probabilities:

$$P(A | B) := \frac{P(A \cap B)}{P(B)}$$

exercice

prove Bayes' theorem:

$$P(A | B) = \frac{P(B | A) \cdot P(A)}{P(B)}$$

$$P(A \cap B) = P(A | B) \cdot P(B)$$

$$P(B \cap A) = P(B | A) \cdot P(A)$$

$$P(A \cap B) = P(B \cap A)$$

$$P(A | B) \cdot P(B) = P(B | A) \cdot P(A) \quad \longrightarrow \quad P(A | B) = \frac{P(B | A) \cdot P(A)}{P(B)}$$

conditional probabilities:

$$P(A | B) := \frac{P(A \cap B)}{P(B)}$$

exercice

prove Bayes' theorem:

$$P(A | B) = \frac{P(B | A) \cdot P(A)}{P(B)}$$

$$P(A \cap B) = P(A | B) \cdot P(B)$$

$$P(B \cap A) = P(B | A) \cdot P(A)$$

$$P(A \cap B) = P(B \cap A)$$

$$P(A | B) \cdot P(B) = P(B | A) \cdot P(A) \quad \longrightarrow \quad P(A | B) = \frac{P(B | A) \cdot P(A)}{P(B)}$$

conditional probabilities:

$$P(A | B) := \frac{P(A \cap B)}{P(B)}$$




Law of total probability

Law of total probability


Ω



Law of total probability




Law of total probability



$$\bigcup_{i=1}^n B_i = \Omega$$


Law of total probability



$$\bigcup_{i=1}^n B_i = \Omega$$

$$B_i \cap B_j = \emptyset$$

Law of total probability



$$\bigcup_{i=1}^n B_i = \Omega$$


$$B_i \cap B_j = \emptyset$$

Law of total probability

$$P(A) = \sum_{i=1}^n P(A | B_i) \cdot P(B_i)$$

$$\bigcup_{i=1}^n B_i = \Omega$$

$$B_i \cap B_j = \emptyset$$




Law of total probability

$$P(A) = \sum_{i=1}^n P(A | B_i) \cdot P(B_i)$$

$$\bigcup_{i=1}^n B_i = \Omega$$

$$B_i \cap B_j = \emptyset$$




prove

Law of total probability

$$P(A) = \sum_{i=1}^n P(A | B_i) \cdot P(B_i)$$

prove



$$\bigcup_{i=1}^n B_i = \Omega \quad B_i \cap B_j = \emptyset$$

$$P(A \cup B) = P(A) + P(B) \quad \text{if } P(A \cap B) = 0$$


$$P(A \cap B) = P(A | B) \cdot P(B)$$

Law of total probability

$$P(A) = \sum_{i=1}^n P(A | B_i) \cdot P(B_i)$$

prove

$$P(A) = P(A \cap \Omega)$$



$$\bigcup_{i=1}^n B_i = \Omega$$

$$B_i \cap B_j = \emptyset$$

$$P(A \cup B) = P(A) + P(B) \quad \text{if } P(A \cap B) = 0$$


$$P(A \cap B) = P(A | B) \cdot P(B)$$

Law of total probability

$$P(A) = \sum_{i=1}^n P(A | B_i) \cdot P(B_i)$$

prove

$$P(A) = P(A \cap \Omega) = P\left(A \cap \bigcup_{i=1}^n B_i\right)$$




$$\bigcup_{i=1}^n B_i = \Omega$$

$$B_i \cap B_j = \emptyset$$

$$P(A \cup B) = P(A) + P(B) \quad \text{if } P(A \cap B) = 0$$

$$P(A \cap B) = P(A | B) \cdot P(B)$$

Law of total probability



$$P(A) = \sum_{i=1}^n P(A | B_i) \cdot P(B_i)$$

$$\bigcup_{i=1}^n B_i = \Omega \quad B_i \cap B_j = \emptyset$$


prove

$$P(A \cup B) = P(A) + P(B) \quad \text{if } P(A \cap B) = 0$$

$$P(A \cap B) = P(A | B) \cdot P(B)$$

$$P(A) = P(A \cap \Omega) = P\left(A \cap \bigcup_{i=1}^n B_i\right) = P\left(\bigcup_{i=1}^n (A \cap B_i)\right)$$

Law of total probability



$$P(A) = \sum_{i=1}^n P(A | B_i) \cdot P(B_i)$$

$$\bigcup_{i=1}^n B_i = \Omega \quad B_i \cap B_j = \emptyset$$


prove

$$P(A \cup B) = P(A) + P(B) \quad \text{if } P(A \cap B) = 0$$

$$P(A \cap B) = P(A | B) \cdot P(B)$$

$$P(A) = P(A \cap \Omega) = P\left(A \cap \bigcup_{i=1}^n B_i\right) = P\left(\bigcup_{i=1}^n (A \cap B_i)\right) = \sum_{i=1}^n P(A \cap B_i)$$

Law of total probability



$$P(A) = \sum_{i=1}^n P(A | B_i) \cdot P(B_i)$$

$$\bigcup_{i=1}^n B_i = \Omega \quad B_i \cap B_j = \emptyset$$

prove

$$P(A \cup B) = P(A) + P(B) \quad \text{if } P(A \cap B) = 0$$

$$P(A \cap B) = P(A | B) \cdot P(B)$$

$$P(A) = P(A \cap \Omega) = P\left(A \cap \bigcup_{i=1}^n B_i\right) = P\left(\bigcup_{i=1}^n (A \cap B_i)\right) = \sum_{i=1}^n P(A \cap B_i) = \sum_{i=1}^n P(A | B_i) \cdot P(B_i)$$

probability expansion

probability expansion

$$P(A_1 \cap A_2 \cap \dots \cap A_n)$$

probability expansion

$$\begin{aligned} P(A_1 \cap A_2 \cap \dots \cap A_n) \\ = P(A_1) \cdot P(A_2 | A_1) \cdot P(A_3 | A_1 \cap A_2) \cdot \dots \cdot P(A_n | A_1 \cap \dots \cap A_{n-1}) \end{aligned}$$

probability expansion

$$\begin{aligned} P(A_1 \cap A_2 \cap \dots \cap A_n) \\ = P(A_1) \cdot P(A_2 | A_1) \cdot P(A_3 | A_1 \cap A_2) \cdot \dots \cdot P(A_n | A_1 \cap \dots \cap A_{n-1}) \end{aligned}$$

prove idea

$$P(A \cap B) = P(A | B) \cdot P(B)$$

probability expansion

$$\begin{aligned} P(A_1 \cap A_2 \cap \dots \cap A_n) \\ = P(A_1) \cdot P(A_2 | A_1) \cdot P(A_3 | A_1 \cap A_2) \cdot \dots \cdot P(A_n | A_1 \cap \dots \cap A_{n-1}) \end{aligned}$$

prove idea

$$B_{n-1} := A_1 \cap \dots \cap A_{n-1}$$

$$P(A \cap B) = P(A | B) \cdot P(B)$$

probability expansion

$$\begin{aligned} P(A_1 \cap A_2 \cap \dots \cap A_n) \\ = P(A_1) \cdot P(A_2 | A_1) \cdot P(A_3 | A_1 \cap A_2) \cdot \dots \cdot P(A_n | A_1 \cap \dots \cap A_{n-1}) \end{aligned}$$

prove idea

$$B_{n-1} := A_1 \cap \dots \cap A_{n-1}$$

$$P(A_1 \cap \dots \cap A_n)$$

$$P(A \cap B) = P(A | B) \cdot P(B)$$

probability expansion

$$\begin{aligned} P(A_1 \cap A_2 \cap \dots \cap A_n) \\ = P(A_1) \cdot P(A_2 | A_1) \cdot P(A_3 | A_1 \cap A_2) \cdot \dots \cdot P(A_n | A_1 \cap \dots \cap A_{n-1}) \end{aligned}$$

prove idea

$$B_{n-1} := A_1 \cap \dots \cap A_{n-1}$$

$$P(A_1 \cap \dots \cap A_n) = P(A_n \cap B_{n-1})$$

$$P(A \cap B) = P(A | B) \cdot P(B)$$

probability expansion

$$\begin{aligned} P(A_1 \cap A_2 \cap \dots \cap A_n) \\ = P(A_1) \cdot P(A_2 | A_1) \cdot P(A_3 | A_1 \cap A_2) \cdot \dots \cdot P(A_n | A_1 \cap \dots \cap A_{n-1}) \end{aligned}$$

prove idea

$$P(A \cap B) = P(A | B) \cdot P(B)$$

$$B_{n-1} := A_1 \cap \dots \cap A_{n-1}$$

$$P(A_1 \cap \dots \cap A_n) = P(A_n \cap B_{n-1}) = P(A_n | B_{n-1}) \cdot P(B_{n-1})$$

probability expansion

$$\begin{aligned} P(A_1 \cap A_2 \cap \dots \cap A_n) \\ = P(A_1) \cdot P(A_2 | A_1) \cdot P(A_3 | A_1 \cap A_2) \cdot \dots \cdot P(A_n | A_1 \cap \dots \cap A_{n-1}) \end{aligned}$$

prove idea

$$P(A \cap B) = P(A | B) \cdot P(B)$$

$$B_{n-1} := A_1 \cap \dots \cap A_{n-1}$$

$$\begin{aligned} P(A_1 \cap \dots \cap A_n) &= P(A_n \cap B_{n-1}) = P(A_n | B_{n-1}) \cdot P(B_{n-1}) \\ &= P(A_1 \cap \dots \cap A_{n-1}) \cdot P(A_n | A_1 \cap \dots \cap A_{n-1}) \end{aligned}$$

probability expansion

$$\begin{aligned} P(A_1 \cap A_2 \cap \dots \cap A_n) \\ = P(A_1) \cdot P(A_2 | A_1) \cdot P(A_3 | A_1 \cap A_2) \cdot \dots \cdot P(A_n | A_1 \cap \dots \cap A_{n-1}) \end{aligned}$$

prove idea

$$P(A \cap B) = P(A | B) \cdot P(B)$$

$$B_{n-1} := A_1 \cap \dots \cap A_{n-1}$$

$$\begin{aligned} P(A_1 \cap \dots \cap A_n) &= P(A_n \cap B_{n-1}) = P(A_n | B_{n-1}) \cdot P(B_{n-1}) \\ &= P(A_1 \cap \dots \cap A_{n-1}) \cdot P(A_n | A_1 \cap \dots \cap A_{n-1}) \\ &= \dots \end{aligned}$$

probability expansion example

probability expansion example

draw 4 cards from 52 cards deck

probability expansion example

draw 4 cards from 52 cards deck

What is the probability that all 4 aces are drawn?

probability expansion example

draw 4 cards from 52 cards deck

What is the probability that all 4 aces are drawn?

A_i : draw an aces in i -th draw

probability expansion example

draw 4 cards from 52 cards deck

What is the probability that all 4 aces are drawn?

A_i : draw an aces in i -th draw

$$P(A_1) = \frac{4}{52},$$

probability expansion example

draw 4 cards from 52 cards deck

What is the probability that all 4 aces are drawn?

A_i : draw an aces in i -th draw

$$P(A_1) = \frac{4}{52}, \quad P(A_2 | A_1) = \frac{3}{51},$$

probability expansion example

draw 4 cards from 52 cards deck

What is the probability that all 4 aces are drawn?

A_i : draw an aces in i -th draw

$$P(A_1) = \frac{4}{52}, \quad P(A_2 | A_1) = \frac{3}{51}, \quad P(A_3 | A_1 \cap A_2) = \frac{2}{50},$$

probability expansion example

draw 4 cards from 52 cards deck

What is the probability that all 4 aces are drawn?

A_i : draw an aces in i -th draw

$$P(A_1) = \frac{4}{52}, \quad P(A_2 | A_1) = \frac{3}{51}, \quad P(A_3 | A_1 \cap A_2) = \frac{2}{50}, \quad P(A_4 | A_1 \cap A_2 \cap A_3) = \frac{1}{49}$$

probability expansion example

draw 4 cards from 52 cards deck

What is the probability that all 4 aces are drawn?

A_i : draw an aces in i -th draw

$$P(A_1) = \frac{4}{52}, \quad P(A_2 | A_1) = \frac{3}{51}, \quad P(A_3 | A_1 \cap A_2) = \frac{2}{50}, \quad P(A_4 | A_1 \cap A_2 \cap A_3) = \frac{1}{49}$$

$$P(A_1 \cap A_2 \cap A_3 \cap A_4)$$

probability expansion example

draw 4 cards from 52 cards deck

What is the probability that all 4 aces are drawn?

A_i : draw an aces in i -th draw

$$P(A_1) = \frac{4}{52}, \quad P(A_2 | A_1) = \frac{3}{51}, \quad P(A_3 | A_1 \cap A_2) = \frac{2}{50}, \quad P(A_4 | A_1 \cap A_2 \cap A_3) = \frac{1}{49}$$

$$P(A_1 \cap A_2 \cap A_3 \cap A_4) = \frac{4}{52} \cdot \frac{3}{51} \cdot \frac{2}{50} \cdot \frac{1}{49}$$

random variable


random variable

A random variable is a number associated to an event.

random variable

A random variable is a number associated to an event.


events




random variable

A random variable is a number associated to an event.

events




random variable




random variable

A random variable is a number associated to an event.




random variable

A random variable is a number associated to an event.



random variable

A random variable is a number associated to an event.



Random variable

A random variable is a number associated to an event.

Random variable

A random variable is a number associated to an event.

binary random variable/ indicator variable/ Bernoulli random variables

Random variable

A random variable is a number associated to an event.

binary random variable/ indicator variable/ Bernoulli random variables

$$\mathbb{B} = \{0,1\}$$

Random variable

A random variable is a number associated to an event.

binary random variable/ indicator variable/ Bernoulli random variables

$$\mathbb{B} = \{0,1\}$$

$$I : \Omega \rightarrow \mathbb{B}$$

Random variable

A random variable is a number associated to an event.

binary random variable/ indicator variable/ Bernoulli random variables

$$\mathbb{B} = \{0,1\}$$

$$I : \Omega \rightarrow \mathbb{B}$$

$$I(\omega) = \begin{cases} 1 & \text{if } \omega = \tilde{\omega} \\ 0 & \text{else} \end{cases}$$

Random variable

A random variable is a number associated to an event.

binary random variable/ indicator variable/ Bernoulli random variables

$$\mathbb{B} = \{0,1\}$$

$$I : \Omega \rightarrow \mathbb{B}$$

$$I(\omega) = \begin{cases} 1 & \text{if } \omega = \tilde{\omega} \\ 0 & \text{else} \end{cases} \quad \tilde{\omega} \in \Omega$$

Random variable

A random variable is a number associated to an event.

binary random variable/ indicator variable/ Bernoulli random variables

$$\mathbb{B} = \{0,1\} \quad I : \Omega \rightarrow \mathbb{B} \quad I(\omega) = \begin{cases} 1 & \text{if } \omega = \tilde{\omega} \\ 0 & \text{else} \end{cases} \quad \tilde{\omega} \in \Omega$$

Random variable

A random variable is a number associated to an event.

binary random variable/ indicator variable/ Bernoulli random variables

$$\mathbb{B} = \{0,1\} \quad I : \Omega \rightarrow \mathbb{B} \quad I(\omega) = \begin{cases} 1 & \text{if } \omega = \tilde{\omega} \\ 0 & \text{else} \end{cases} \quad \tilde{\omega} \in \Omega$$

discrete random variable

Random variable

A random variable is a number associated to an event.

binary random variable/ indicator variable/ Bernoulli random variables

$$\mathbb{B} = \{0,1\} \quad I : \Omega \rightarrow \mathbb{B} \quad I(\omega) = \begin{cases} 1 & \text{if } \omega = \tilde{\omega} \\ 0 & \text{else} \end{cases} \quad \tilde{\omega} \in \Omega$$

discrete random variable

$$S := \{x_1, x_2, x_3, \dots\}$$

Random variable

A random variable is a number associated to an event.

binary random variable/ indicator variable/ Bernoulli random variables

$$\mathbb{B} = \{0,1\} \quad I : \Omega \rightarrow \mathbb{B} \quad I(\omega) = \begin{cases} 1 & \text{if } \omega = \tilde{\omega} \\ 0 & \text{else} \end{cases} \quad \tilde{\omega} \in \Omega$$

discrete random variable

$$S := \{x_1, x_2, x_3, \dots\} \quad \text{finite or countable infinite}$$

Random variable

A random variable is a number associated to an event.

binary random variable/ indicator variable/ Bernoulli random variables

$$\mathbb{B} = \{0,1\} \quad I : \Omega \rightarrow \mathbb{B} \quad I(\omega) = \begin{cases} 1 & \text{if } \omega = \tilde{\omega} \\ 0 & \text{else} \end{cases} \quad \tilde{\omega} \in \Omega$$

discrete random variable

$$S := \{x_1, x_2, x_3, \dots\} \quad \text{finite or countable infinite}$$

$$X : \Omega \rightarrow S$$

Random variable

A random variable is a number associated to an event.

binary random variable/ indicator variable/ Bernoulli random variables

$$\mathbb{B} = \{0,1\} \quad I : \Omega \rightarrow \mathbb{B} \quad I(\omega) = \begin{cases} 1 & \text{if } \omega = \tilde{\omega} \\ 0 & \text{else} \end{cases} \quad \tilde{\omega} \in \Omega$$

discrete random variable

$$S := \{x_1, x_2, x_3, \dots\} \quad \text{finite or countable infinite}$$

$$X : \Omega \rightarrow S \quad X(\omega) = x_\omega \in S$$

Random variable

A random variable is a number associated to an event.

binary random variable/ indicator variable/ Bernoulli random variables

$$\mathbb{B} = \{0,1\} \quad I : \Omega \rightarrow \mathbb{B} \quad I(\omega) = \begin{cases} 1 & \text{if } \omega = \tilde{\omega} \\ 0 & \text{else} \end{cases} \quad \tilde{\omega} \in \Omega$$

discrete random variable

$$S := \{x_1, x_2, x_3, \dots\} \quad \text{finite or countable infinite}$$

$$X : \Omega \rightarrow S \quad X(\omega) = x_\omega \in S$$

$$P(X = x_i) = ?$$

Random variable

A random variable is a number associated to an event.

binary random variable/ indicator variable/ Bernoulli random variables

$$\mathbb{B} = \{0,1\} \quad I : \Omega \rightarrow \mathbb{B} \quad I(\omega) = \begin{cases} 1 & \text{if } \omega = \tilde{\omega} \quad \tilde{\omega} \in \Omega \\ 0 & \text{else} \end{cases}$$

discrete random variable

$$S := \{x_1, x_2, x_3, \dots\} \quad X : \Omega \rightarrow S \quad X(\omega) = x_\omega \in S$$

Random variable

A random variable is a number associated to an event.

binary random variable/ indicator variable/ Bernoulli random variables

$$\mathbb{B} = \{0,1\} \quad I : \Omega \rightarrow \mathbb{B} \quad I(\omega) = \begin{cases} 1 & \text{if } \omega = \tilde{\omega} \quad \tilde{\omega} \in \Omega \\ 0 & \text{else} \end{cases}$$

discrete random variable

$$S := \{x_1, x_2, x_3, \dots\} \quad X : \Omega \rightarrow S \quad X(\omega) = x_\omega \in S$$

continuous random variables

Random variable

A random variable is a number associated to an event.

binary random variable/ indicator variable/ Bernoulli random variables

$$\mathbb{B} = \{0,1\} \quad I : \Omega \rightarrow \mathbb{B} \quad I(\omega) = \begin{cases} 1 & \text{if } \omega = \tilde{\omega} \quad \tilde{\omega} \in \Omega \\ 0 & \text{else} \end{cases}$$

discrete random variable

$$S := \{x_1, x_2, x_3, \dots\} \quad X : \Omega \rightarrow S \quad X(\omega) = x_\omega \in S$$

continuous random variables

$$J \subseteq \mathbb{R}$$

Random variable

A random variable is a number associated to an event.

binary random variable/ indicator variable/ Bernoulli random variables

$$\mathbb{B} = \{0,1\} \quad I : \Omega \rightarrow \mathbb{B} \quad I(\omega) = \begin{cases} 1 & \text{if } \omega = \tilde{\omega} \quad \tilde{\omega} \in \Omega \\ 0 & \text{else} \end{cases}$$

discrete random variable

$$S := \{x_1, x_2, x_3, \dots\} \quad X : \Omega \rightarrow S \quad X(\omega) = x_\omega \in S$$

continuous random variables

$$J \subseteq \mathbb{R}$$

$$X : \Omega \rightarrow J$$

Random variable

A random variable is a number associated to an event.

binary random variable/ indicator variable/ Bernoulli random variables

$$\mathbb{B} = \{0,1\} \quad I : \Omega \rightarrow \mathbb{B} \quad I(\omega) = \begin{cases} 1 & \text{if } \omega = \tilde{\omega} \quad \tilde{\omega} \in \Omega \\ 0 & \text{else} \end{cases}$$

discrete random variable

$$S := \{x_1, x_2, x_3, \dots\} \quad X : \Omega \rightarrow S \quad X(\omega) = x_\omega \in S$$

continuous random variables

$$J \subseteq \mathbb{R}$$

$$P(X = x_i) = 0$$

$$X : \Omega \rightarrow J$$

Random variable

A random variable is a number associated to an event.

binary random variable/ indicator variable/ Bernoulli random variables

$$\mathbb{B} = \{0,1\} \quad I : \Omega \rightarrow \mathbb{B} \quad I(\omega) = \begin{cases} 1 & \text{if } \omega = \tilde{\omega} \quad \tilde{\omega} \in \Omega \\ 0 & \text{else} \end{cases}$$

discrete random variable

$$S := \{x_1, x_2, x_3, \dots\} \quad X : \Omega \rightarrow S \quad X(\omega) = x_{\omega} \in S$$

continuous random variables

$$J \subseteq \mathbb{R}$$

$$X : \Omega \rightarrow J$$

$$P(X = x_i) = 0$$

$$P(X > x_i) = ?$$

end