

The critical $O(N)$ CFT: Methods and conformal data. Updated tables

Johan Henriksson¹

¹*Theoretical Physics Department, CERN, 1211, Geneva, Switzerland*

E-mail: johan.henriksson@cern.ch

ABSTRACT: This note contains up-to-date tables of [1]. Superseded results are displayed in gray, new results in orange.

Please email the author to report any omissions.

Date: September 3, 2026.

Contents

1	Summary	1
1.1	New perturbative studies	1
1.2	New MC studies	1
1.3	New bootstrap studies	1
1.4	New formal studies	1
2	New implementations	2
3	Tables	3
3.1	Families of operators for $N = 1$	7
3.2	Operators in non-traceless-symmetric Lorentz representations	16
3.2.1	Three-row Lorentz Young tableaux	16
3.3	OPE coefficients	18
3.4	Families of operators for general N	21
3.4.1	Spinning singlet operators	26
3.5	Operators in non-traceless-symmetric Lorentz representations	34
3.6	OPE coefficients in the $\varphi \times \varphi$ OPE	35
A	Some extension on representation theory	37
B	Extended character decomposition	38

1 Summary

We give references to papers that appeared after 24 January 2022 (and papers which appeared before that date but which were accidentally omitted).

1.1 New perturbative studies

- We have made a few computations specifically for the preparation of this note and which are not presented elsewhere. They are cited as [2].
- Many high-loop results for operators in the ε -expansion were computed in [3]¹ and [4, 5].²

¹Appeared before January 2022 but they were not include it in [1] due to oversight.

²Complete results of [5] are released in [6], however we only give reference to [5].

- The OPE coefficient involving $\mathbf{Op}[\mathbf{E}, 2, 2]$ and two ϕ was determined exactly in [7].
- The seven-loop results, including $O(\varepsilon^8)$ for Δ_ϕ , have been presented in [8].
- [9]: New result for the $O(N)$ free energy.

1.2 New MC studies

- Conventional MC studies: Ising [10], $O(2)$ [11] $O(3)$ [12].
- Fuzzy sphere studies: Ising [13–17], $O(2)$ [18]³, $O(3)$ [19, 20].

1.3 New bootstrap studies

- Ising [21–24].

1.4 New formal studies

- Status of the $2 + \tilde{\varepsilon}$ expansion [25].

2 New implementations

We have implemented many new operators in the data file `ONdata.m`. This now contains all Ising scalar operators with $\Delta_{4d} \leq 12$, all Ising operators with $\Delta_{4d} \leq 10$, all $O(N)$ scalar operators with $\Delta_{4d} \leq 8$, all $O(N)$ spin- ℓ operators with $\Delta_{4d} \leq 7$, and remaining (Lorentz non-traceless-symmetric) $O(N)$ operators with $\Delta_{4d} \leq 6$. In addition several operators with higher scaling dimension has been included in a non-systematic fashion. On the tables below, operator names are displayed in orange where new results are added.

We have also implemented the critical coupling as

$$\text{ValueE}[\text{criticalCoupling}] = \frac{3\mathbf{e}}{\mathbf{n} + 8} + \frac{9(3\mathbf{n} + 14)\mathbf{e}^2}{(\mathbf{n} + 8)^3} + \dots$$

³https://scgp.stonybrook.edu/video-portal/results.php?event_id=437.

3 Tables

Table 1: Estimates and measurements of the critical exponents η and ν in the three-dimensional Ising CFT.

	η	ν
Free theory	0	0.5
Wilson–Fisher $O(\varepsilon^3)$ truncation	0.037209	0.60750
Wilson–Fisher $O(\varepsilon^7)$ resummation [26]	0.03653(65)	0.62977(22)
Uniaxial magnet FeF ₂ (1987) [27]		0.64(1)
Liquid-gas transition CO ₂ (1998) [28]	0.042(6)	
Liquid-gas transition D ₂ O (2000) [29]		0.62(3)
Liquid mixture* (1988) [30]	0.032(13)	0.629(3)
Liquid mixture [†] (2004) [31]	0.041(5)	0.632(2)
Quantum phase transition (2020) [32]		0.62(4)
High temperature expansion [33]	0.03639(15)	0.63012(16)
MC simulations [34]	0.036284(40)	0.62998(5)
Numerical bootstrap [35]	0.0362978(20)	0.629971(4)
Numerical bootstrap [23]	0.036297612(48)	0.62997097(12)

*Light-scattering measurement in a mixture of water and isobutyric acid.

[†]Turbidity measurement in a mixture of methanol and cyclohexane.

Table 3: A summary of the developments of higher order results for the main observables in the ε -expansion and $1/N$ expansion. The notation is “order[ref.]”.

$d = 4 - \varepsilon$ expansion		$1/N$ expansion	
Δ_φ	ε^4 [36] ε^5 [37, 38]* ε^6 [39] ε^7 [40] ε^8 [41]	Δ_φ	N^{-2} (3d)[42] N^{-2} [43, 44] N^{-3} [45] [†]
$\Delta_{\varphi_S^2}$	ε^4 [46] ε^5 [38] ε^6 [47] ε^7 [40]	Δ_σ	N^{-2} (3d)[48] N^{-2} [49]
$\Delta_{\varphi_S^4}$	ε^4 [46] ε^5 [38] ε^6 [47] ε^7 [40]	Δ_{σ^2}	N^{-1} [50] [‡] N^{-2} [51, 52]
$\Delta_{\varphi_T^2}$	ε^2 [53] ε^3 [54] ε^4 [55] ε^6 [56] [§]	$\Delta_{\varphi_T^2}$	N^{-1} [50] N^{-2} [57]
$\Delta_{\varphi_{T_4}^4}$	ε^5 [58, 59] ε^6 [60] [¶]	$\Delta_{\varphi_{T_4}^4}$	N^{-1} [61] N^{-2} [62]
C_T	ε^2 [63] ε^3 [64] ε^4 [65]	C_T	N^{-1} [66]
C_J	ε^2 [67]** ε^3 [64] ε^4 [65]	C_J	N^{-1} (3d)[68] N^{-1} [69]

*The result in [37] contained several errors, see chapter 15 of [70].

[†]As noted in [71], the expression in [45] contains a misprint in equation (22). See also [72, 73].

[‡]Ref. [74] (page 221) cites [50] together with Bervillier, Girardi, Brezin; Saclay preprint (1974).

[§]Note that their $\zeta_{3,5}$ equals our $\zeta_{5,3}$.

[¶]This result was extracted from the more general results of [60].

^{||,**}Can be extracted from [75], see comment in [67].

Table 5: Results in three dimensions from the conformal bootstrap. The lower part of the table contains the critical exponents, computed from the values in the upper part. For comparison, we have added the exact results at $N = \infty$ (sometimes denoted the spherical model).

Quantity	$N = 1$	$N = 2$	$N = 3$	$N = \infty$
$\Delta_\varphi (\Delta_\sigma)$	0.518148806(24) [23]	0.519088(22) [76]	0.518942(51) [77]	0.5
$\Delta_{\varphi_S^2} (\Delta_\epsilon)$	1.41262528(29) [23]	1.51136(22) [76]	1.59489(59) [77]	2
$\Delta_{\varphi_S^4} (\Delta_{\epsilon'})$	3.82951(61) [78]	3.794(8) [79]	3.7668(100) [80]*	4
$\Delta_{\varphi_T^2} (\Delta_t)$		1.23629(11) [76]	1.20954(32) [77]	1
$\Delta_{\varphi_{T_4}^4} (\Delta_{t_4})$		3.11535(73) [76]	$< 2.99056^\dagger$ [77]	2
$\frac{C_T}{C_{T,\text{free}}}$	0.946543(42) [78] 0.946538675(42) [23]	0.944056(15) [76]	0.944524(28) [77]	1
$\frac{C_J}{C_{J,\text{free}}}$		0.904395(28) [76]	0.90632(16) [77]	1
α	0.11008708(35)	-0.01526(30)	-0.1350(12)	-1
β	0.32641871(6)	0.348699(54)	0.36932(22)	0.5
γ	1.23707551(23)	1.31786(20)	1.39641(81)	2
δ	4.78984254(23)	4.77937(24)	4.78106(75)	5
η	0.036297612(48)	0.038176(44)	0.03787(13)	0
ν	0.62997097(12) [23]	0.671754(99) [76]	0.71168(41) [77]	1
ϕ_c		1.18478(19) [76]	1.27424(83) [77]	2
ω	0.82951(61) [78]	0.794(8) [79]	0.7668(100) [80]	1

*Computed using $\Lambda = 35$ data from 42 sampling points [80].

† This value is a rigorous upper bound and the true operator dimension is expected to lie not far from this value.

Table 6: Results in three dimensions from Monte Carlo simulations. For $N = 0$ the exponent ν and the Hausdorff dimension $D_H = d - \Delta_{\varphi_T^2}$ are directly related.

Quantity	$N = 0$	$N = 1$	$N = 2$	$N = 3$
γ	1.15695300(95) [81]			
η	0.0250(14) [82]	0.036284(40) [34]	0.03810(8) [83]	0.03784(5) [84]
ν	0.58759700(40) [85]	0.62998(5) [34]	0.67169(7) [83]	0.71164(10) [84]
ω	0.9037(56) [86]	0.832(6) [87]	0.789(4) [83]	0.759(2) [84]
d_H	1.7018467(16)* [85]	1.7349(65) [88]	1.7626(66) [88]	1.7906(3) [89]

*Computed from $\Delta_{\varphi_T^2}$.

3.1 Families of operators for $N = 1$

IsingF1[k] Scalar operators of the form ϕ^k , $k \geq 1$, $k \neq 3$. We have ε^2 [90] ε^5 [5, 91],⁴

$$\text{DeltaE}[\text{IsingF1}[k]] = k \left[1 - \frac{\varepsilon}{2}\right] + \frac{k(k-1)}{6}\varepsilon - \frac{k(17k^2 - 67k + 47)}{324}\varepsilon^2 + \dots + O(\varepsilon^6). \quad (3.1)$$

IsingF2[l] Broken currents of the form $\mathcal{J}_\ell$, $\ell \in 2\mathbb{Z}_+$. We have ε^2 [92] ε^4 [93],

$$\text{DeltaE}[\text{IsingF2}[l]] = 2 + l - \varepsilon + \left(\frac{1}{54} - \frac{1}{9l(l+1)}\right)\varepsilon^2 + \dots + O(\varepsilon^5). \quad (3.2)$$

IsingF3[l] Spinning operators of the form $\partial^\ell \phi^3$, $\ell \geq 2$, ε^1 [94] ε^2 [7],

$$\text{DeltaE}[\text{IsingF3}[l]] = 3 + l - \frac{3\varepsilon}{2} + \frac{\varepsilon}{3} \left(1 + \frac{2(-1)^l}{l+1}\right) + \dots + O(\varepsilon^3). \quad (3.3)$$

Starting from $\ell = 6$, there are additional operators of the type $\partial^\ell \phi^3$, however they all have vanishing first-order anomalous dimensions. The family **IsingF3** is identified by the non-vanishing of the anomalous dimension to order ε .

IsingF4[k] Spin-two operators of the form $\partial^2 \phi^k$, $k \geq 2$, ε^1 [95] ε^2 [6, 91],⁵

$$\text{DeltaE}[\text{IsingF4}[k]] = 2 + k \left[1 - \frac{\varepsilon}{2}\right] + \frac{(k-2)(3k+1)}{18}\varepsilon - \frac{(k-2)(102k^2 - 335k - 235)}{1944}\varepsilon^2 + O(\varepsilon^3). \quad (3.4)$$

IsingF5[k] Spin-three operators of the form $\partial^3 \phi^k$, $k \geq 3$, $k \neq 4$, ε^1 [95],

$$\text{DeltaE}[\text{IsingF5}[k]] = 3 + k \left[1 - \frac{\varepsilon}{2}\right] + \frac{k^2 - 2k - 2}{6}\varepsilon + O(\varepsilon^2). \quad (3.5)$$

IsingF6[k] Spin-four operators of the form $\partial^4 \phi^k$, $k \geq 2$, ε^1 [95], ε^2 [96],

$$\text{DeltaE}[\text{IsingF6}[k]] = 4 + k \left[1 - \frac{\varepsilon}{2}\right] + \frac{(k-2)(5k-1)}{30}\varepsilon + O(\varepsilon^2). \quad (3.6)$$

IsingF7[k] Spin-four operators of the form $\partial^4 \phi^k$, $k \geq 4$, ε^1 [95], ε^2 [96],

$$\text{DeltaE}[\text{IsingF7}[k]] = 4 + k \left[1 - \frac{\varepsilon}{2}\right] + \frac{3k^2 - 7k - 12}{18}\varepsilon + O(\varepsilon^2). \quad (3.7)$$

IsingF8[k] Spin-five operators of the form $\partial^5 \phi^k$, $k \geq 3$, $k \neq 4$, ε^1 [95],

$$\text{DeltaE}[\text{IsingF8}[k]] = 5 + k \left[1 - \frac{\varepsilon}{2}\right] + \frac{3k^2 - 7k - 2}{18}\varepsilon + O(\varepsilon^2). \quad (3.8)$$

⁴The formula was first found using results in [5], then independently in [91].

⁵The formula was derived in [6] by combining results from [5] and [91]

IsingF9[k] Spin-five operators of the form $\partial^5 \phi^k$, $k \geq 5$, ε^1 [95],

$$\text{DeltaE}[\text{IsingF9}[k]] = 5 + k \left[1 - \frac{\varepsilon}{2}\right] + \frac{3k^2 - 8k - 20}{18} \varepsilon + O(\varepsilon^2). \quad (3.9)$$

IsingF10[k] Scalar operators of the form $\square^2 \phi^k$, $k \geq 4$, ε^1 [95] ε^2 [6, 91],⁶,

$$\text{DeltaE}[\text{IsingF10}[k]] = 4 + k \left[1 - \frac{\varepsilon}{2}\right] + \frac{k(3k - 7)}{18} \varepsilon - \frac{51k^3 - 338k^2 + 443k + 108}{972} + O(\varepsilon^3). \quad (3.10)$$

IsingF11[k] Spin-one operators of the form $\partial \square^2 \phi^k$, $k \geq 5$, ε^1 [95],

$$\text{DeltaE}[\text{IsingF11}[k]] = 5 + k \left[1 - \frac{\varepsilon}{2}\right] + \frac{3k^2 - 8k - 4}{18} \varepsilon + O(\varepsilon^2) \quad (3.11)$$

IsingF12[k] Spin-two operators of the form $\partial^2 \square \phi^k$, $k \geq 4$, ε^1 [95],

$$\text{DeltaE}[\text{IsingF12}[k]] = 4 + k \left[1 - \frac{\varepsilon}{2}\right] + \frac{3k^2 - 7k - 4}{18} \varepsilon + O(\varepsilon^2). \quad (3.12)$$

IsingF13[k] Spin-three operators of the form $\partial^3 \square \phi^k$, $k \geq 5$, ε^1 [95],

$$\text{DeltaE}[\text{IsingF13}[k]] = 5 + k \left[1 - \frac{\varepsilon}{2}\right] + \frac{3k^2 - 8k - 10}{18} \varepsilon + O(\varepsilon^2). \quad (3.13)$$

⁶The formula was derived in [6] by combining results from [5] and [91]

Table 8: $\mathbb{Z}_2$ even scalar operators for $N = 1$. The table includes operators with $\Delta^{4d} \leq 12$. The fact that there are two operators with anomalous dimension $\frac{8}{3}$ of the form $\square^3 \phi^6$ is not a typo, for instance both operators are reported in table 5 of [95].

$\mathcal{O}$	$\mathcal{O} _{\varepsilon}$	$\Delta_{4-\varepsilon}^{(1)}$	$\Delta(\varepsilon)$	Δ_{3d}	Family
$\text{Op}[E, 0, 0]$	$\mathbf{1}$	0	exact	0	
$\text{Op}[E, 0, 1]$	ϕ^2	$[2 - \varepsilon] + \frac{1}{3}\varepsilon$	$\varepsilon^7[40]$	1.412625(10)[35]	1_2
$\text{Op}[E, 0, 2]$	ϕ^4	$[4 - 2\varepsilon] + 2\varepsilon$	$\varepsilon^7[40]$	3.82951(61)[78]	1_4
$\text{Op}[E, 0, 3]$	ϕ^6	$[6 - 3\varepsilon] + 5\varepsilon$	$\varepsilon^2[90]\varepsilon^5[3]$	6.8956(43)[97]	1_6
$\text{Op}[E, 0, 4]$	$\square^2 \phi^4$	$[8 - 2\varepsilon] + \frac{10}{9}\varepsilon$	$\varepsilon^4[5]$	7.2535(51)[97]	10_4
$\text{Op}[E, 0, 5]$	ϕ^8	$[8 - 4\varepsilon] + \frac{28}{3}\varepsilon$	$\varepsilon^2[90]\varepsilon^5[5]$		1_8
$\text{Op}[E, 0, 6]$	$\square^3 \phi^4$	$[10 - 2\varepsilon] + \frac{1}{3}\varepsilon$	$\varepsilon^2[4]$		
$\text{Op}[E, 0, 7]$	$\square^2 \phi^6$	$[10 - 3\varepsilon] + \frac{11}{3}\varepsilon$	$\varepsilon^1 \varepsilon^2[4]$		10_6
$\text{Op}[E, 0, 8]$	ϕ^{10}	$[10 - 5\varepsilon] + 15\varepsilon$	$\varepsilon^2[90]\varepsilon^5[5]$		1_{10}
$\text{Op}[E, 0, 9]$	$\square^4 \phi^4$	$[12 - 2\varepsilon] + \frac{14}{15}\varepsilon$	ε^1		
$\text{Op}[E, 0, 10]$	$\square^3 \phi^6$	$[12 - 3\varepsilon] + \frac{8}{3}\varepsilon$	ε^1		
$\text{Op}[E, 0, 11]$	$\square^3 \phi^6$	$[12 - 3\varepsilon] + \frac{8}{3}\varepsilon$	ε^1		
$\text{Op}[E, 0, 12]$	$\square^2 \phi^8$	$[12 - 4\varepsilon] + \frac{68}{9}\varepsilon$	$\varepsilon^2[6, 91]$		10_8
$\text{Op}[E, 0, 13]$	ϕ^{12}	$[12 - 6\varepsilon] + 22\varepsilon$	$\varepsilon^2[90]\varepsilon^5[5]$		1_{12}

Table 9: $\mathbb{Z}_2$ odd scalar operators for $N = 1$. The table includes operators with $\Delta^{4d} \leq 11$.

$\mathcal{O}$	$\mathcal{O} \varepsilon$	$\Delta_{4-\varepsilon}^{(1)}$	$\Delta(\varepsilon)$	Δ_{3d}	Family
$0p[0,0,1]$	ϕ	$[1 - \frac{1}{2}\varepsilon] + 0\varepsilon$	$\varepsilon^8[41]$	$0.5181489(10)[35]$	1_1
$0p[0,0,2]$	ϕ^5	$[5 - \frac{5}{2}\varepsilon] + \frac{10}{3}\varepsilon$	$\varepsilon^3[98]\varepsilon^5[5]$	$5.2906(11)[97]^*$	1_5
$0p[0,0,3]$	ϕ^7	$[7 - \frac{7}{2}\varepsilon] + 7\varepsilon$	$\varepsilon^2[90]\varepsilon^5[5]$		1_7
$0p[0,0,4]$	$\square^2\phi^5$	$[9 - \frac{5}{2}\varepsilon] + \frac{20}{9}\varepsilon$	$\varepsilon^3[5]$		10_5
$0p[0,0,5]$	ϕ^9	$[9 - \frac{9}{5}\varepsilon] + 12\varepsilon$	$\varepsilon^2[90]\varepsilon^5[5]$		1_9
$0p[0,0,6]$	$\square^3\phi^5$	$[11 - \frac{5}{2}\varepsilon] + \frac{4}{3}\varepsilon$	ε^1		
$0p[0,0,7]$	$\square^2\phi^7$	$[11 - \frac{7}{2}\varepsilon] + \frac{49}{9}$	$\varepsilon^2[6, 91]$		10_7
$0p[0,0,8]$	ϕ^{11}	$[11 - \frac{11}{2}\varepsilon] + \frac{55}{3}\varepsilon$	$\varepsilon^2[90]\varepsilon^5[5]$		1_{11}

*A value with rigorous error bars, $5.262(89)$, was given in [78].

Table 10: $\mathbb{Z}_2$ even operators of spin two, for $N = 1$. The table includes operators with $\Delta_{4d} \leq 10$.

$\mathcal{O}$	$\mathcal{O} \varepsilon$	$\Delta_{4-\varepsilon}^{(1)}$	$\Delta(\varepsilon)$	Δ_{3d}	Family
$\text{Op}[\text{E}, 2, 1]$	$T^{\mu\nu}$	$[4 - \varepsilon]$	exact	3	$2_2, 4_2$
$\text{Op}[\text{E}, 2, 2]$	$\partial^2 \phi^4$	$[6 - 2\varepsilon] + \frac{13}{9}\varepsilon$	$\varepsilon^5[5]$	$5.50915(44)[97]^*$	4_4
$\text{Op}[\text{E}, 2, 3]$	$\partial^2 \square \phi^4$	$[8 - 2\varepsilon] + \frac{8}{9}\varepsilon$	ε^1	$7.0758(58)[97]$	12_4
$\text{Op}[\text{E}, 2, 4]$	$\partial^2 \phi^6$	$[8 - 3\varepsilon] + \frac{38}{9}\varepsilon$	$\varepsilon^2[6, 91]$		4_6
$\text{Op}[\text{E}, 2, 5]$	$\partial^2 \square^2 \phi^4$	$[10 - 2\varepsilon] + \frac{1}{9}\varepsilon$	ε^1		
$\text{Op}[\text{E}, 2, 6]$	$\partial^2 \square^2 \phi^4$	$[10 - 2\varepsilon] + \frac{14}{15}\varepsilon$	ε^1		
$\text{Op}[\text{E}, 2, 7]$	$\partial^2 \square \phi^6$	$[10 - 3\varepsilon] + \frac{31}{9}\varepsilon$	ε^1		12_6
$\text{Op}[\text{E}, 2, 8]$	$\partial^2 \phi^8$	$[10 - 4\varepsilon] + \frac{25}{3}\varepsilon$	$\varepsilon^2[6, 91]$		4_8

*A value with rigorous error bars, $5.499(17)$, was given in [78].

Table 11: $\mathbb{Z}_2$ even operators of spin four, for $N = 1$. The table includes operators with $\Delta^{4d} \leq 10$.

$\mathcal{O}$	$\mathcal{O} \varepsilon$	$\Delta_{4-\varepsilon}^{(1)}$	$\Delta(\varepsilon)$	Δ_{3d}	Family
$\text{Op}[E, 4, 1]$	$C^{\mu\nu\rho\sigma}$	$[6 - \varepsilon] + 0\varepsilon$	ε^4 [93]	5.022665(28)[97]	$2_4, 6_2$
$\text{Op}[E, 4, 2]$	$\partial^4\phi^4$	$[8 - 2\varepsilon] + \frac{4}{9}\varepsilon$	ε^2 [96]	6.42065(64)[97]	7_4
$\text{Op}[E, 4, 3]$	$\partial^4\phi^4$	$[8 - 2\varepsilon] + \frac{19}{15}\varepsilon$	ε^2 [96]	7.38568(28)[97]	6_4
$\text{Op}[E, 4, 4]$	$\partial^4\Box\phi^4$	$[10 - 2\varepsilon] + \frac{49-\sqrt{697}}{90}\varepsilon$	ε^1	8.9410(99)[97]	
$\text{Op}[E, 4, 5]$	$\partial^4\Box\phi^4$	$[10 - 2\varepsilon] + \frac{49+\sqrt{697}}{90}\varepsilon$	ε^1		
$\text{Op}[E, 4, 6]$	$\partial^4\phi^6$	$[10 - 3\varepsilon] + \frac{58}{15}\varepsilon$	ε^2 [96]		6_6
$\text{Op}[E, 4, 7]$	$\partial^4\phi^6$	$[10 - 3\varepsilon] + 3\varepsilon$	ε^2 [96]		7_6

Table 12: $\mathbb{Z}_2$ even operators for $N = 1$ of leading twist for even spin. The identification with numerical data can only be done for the smallest operator at each spin. For spin 6 and 8 we also include operators of subleading twist $\tau^{4d} = 4$.

$\mathcal{O}$	$\mathcal{O} \varepsilon$	$\Delta_{4-\varepsilon}^{(1)}$	$\Delta(\varepsilon)$	Δ_{3d}	Family
$\text{Op}[\text{E}, 6, 1]$	$\mathcal{J}_6$	$[8 - \varepsilon] + 0\varepsilon$	ε^4 [93]	7.028488(16)[97]	2_7
$\text{Op}[\text{E}, 6, 2]$	$\partial^6 \phi^4$	$[10 - 2\varepsilon] + 0.4966\varepsilon$	ε^1		
$\text{Op}[\text{E}, 6, 3]$	$\partial^6 \phi^4$	$[10 - 2\varepsilon] + 0.6707\varepsilon$	ε^1		
$\text{Op}[\text{E}, 6, 4]$	$\partial^6 \phi^4$	$[10 - 2\varepsilon] + 1.1787\varepsilon$	ε^1		
$\text{Op}[\text{E}, 8, 1]$	$\mathcal{J}_8$	$[10 - \varepsilon] + 0\varepsilon$	ε^4 [93]	9.031023(30)[97]	2_8
$\text{Op}[\text{E}, 8, 2]$	$\partial^8 \phi^4$	$[12 - 2\varepsilon] + 0.1142\varepsilon$	ε^1		
$\text{Op}[\text{E}, 8, 3]$	$\partial^8 \phi^4$	$[12 - 2\varepsilon] + 0.5217\varepsilon$	ε^1		
$\text{Op}[\text{E}, 8, 4]$	$\partial^8 \phi^4$	$[12 - 2\varepsilon] + 0.6752\varepsilon$	ε^1		
$\text{Op}[\text{E}, 8, 5]$	$\partial^8 \phi^4$	$[12 - 2\varepsilon] + 1.1276\varepsilon$	ε^1		
$\text{Op}[\text{E}, 10, 1]$	$\mathcal{J}_{10}$	$[12 - \varepsilon] + 0\varepsilon$	ε^4 [93]	11.0324141(99)[97]	2_{10}
$\text{Op}[\text{E}, 12, 1]$	$\mathcal{J}_{12}$	$[14 - \varepsilon] + 0\varepsilon$	ε^4 [93]	13.033286(12)[97]	2_{12}
$\text{Op}[\text{E}, 14, 1]$	$\mathcal{J}_{14}$	$[16 - \varepsilon] + 0\varepsilon$	ε^4 [93]	15.033838(15)[97]	2_{14}
$\text{Op}[\text{E}, 16, 1]$	$\mathcal{J}_{16}$	$[18 - \varepsilon] + 0\varepsilon$	ε^4 [93]	17.034258(34)[97]	2_{16}

Table 13: $\mathbb{Z}_2$ even operators for $N = 1$ of leading twist for odd spin.

$\mathcal{O}$	$\mathcal{O} _\varepsilon$	$\Delta_{4-\varepsilon}^{(1)}$	$\Delta(\varepsilon)$	Δ_{3d}	Family
$\text{Op}[\text{E}, 1, 1]$	$\partial \square^2 \phi^6$	$[11 - 3\varepsilon] + \frac{28}{9}\varepsilon$	ε^1		11_6
$\text{Op}[\text{E}, 3, 1]$	$\partial^3 \phi^6$	$[9 - 3\varepsilon] + \frac{11}{3}\varepsilon$	ε^2 [96]		5_6
$\text{Op}[\text{E}, 5, 1]$	$\partial^5 \square \phi^4$	$[11 - 2\varepsilon] + \frac{29}{45}\varepsilon$	ε^1		
$\text{Op}[\text{E}, 5, 2]$	$\partial^5 \phi^6$	$[11 - 3\varepsilon] + \frac{20}{9}\varepsilon$	ε^1		9_6
$\text{Op}[\text{E}, 5, 3]$	$\partial^5 \phi^6$	$[11 - 3\varepsilon] + \frac{32}{9}\varepsilon$	ε^1		8_6
$\text{Op}[\text{E}, 7, 1]$	$\partial^7 \phi^4$	$[11 - 2\varepsilon] + \frac{1}{3}\varepsilon$	ε^1		
$\text{Op}[\text{E}, 9, 1]$	$\partial^9 \phi^4$	$[13 - 2\varepsilon] + \frac{1}{3}\varepsilon$	ε^1		
$\text{Op}[\text{E}, 9, 2]$	$\partial^9 \phi^4$	$[13 - 2\varepsilon] + \frac{5}{9}\varepsilon$	ε^1		

Table 14: A selection of spinning $\mathbb{Z}_2$ odd operators for $N = 1$. The table includes all such operators with $\Delta^{4d} \leq 10$.

$\mathcal{O}$	$\mathcal{O} _\varepsilon$	$\Delta_{4-\varepsilon}^{(1)}$	$\Delta(\varepsilon)$	Δ_{3d}	Family
$\text{Op}[0,1,1]$	$\partial \square^2 \phi^5$	$[10 - \frac{5}{2}\varepsilon] + \frac{31}{18}\varepsilon$	ε^1		11_5
$\text{Op}[0,2,1]$	$\partial^2 \phi^3$	$[5 - \frac{3}{2}\varepsilon] + \frac{5}{9}\varepsilon$	$\varepsilon^2[7]\varepsilon^5[5]$	4.180305(18)[97]	$3_2, 4_3$
$\text{Op}[0,2,2]$	$\partial^2 \phi^5$	$[7 - \frac{5}{2}\varepsilon] + \frac{8}{3}\varepsilon$	$\varepsilon^2[6, 91]$	6.9873(53)[97]	4_5
$\text{Op}[0,2,3]$	$\partial^2 \square \phi^5$	$[9 - \frac{5}{2}\varepsilon] + 2\varepsilon$	ε^1		12_5
$\text{Op}[0,2,4]$	$\partial^2 \phi^7$	$[9 - \frac{7}{2}\varepsilon] + \frac{55}{9}\varepsilon$	$\varepsilon^2[6, 91]$		4_7
$\text{Op}[0,3,1]$	$\partial^3 \phi^3$	$[6 - \frac{3}{2}\varepsilon] + \frac{1}{6}\varepsilon$	$\varepsilon^2[7]$	4.63804(88)[97]	$3_3, 5_3$
$\text{Op}[0,3,2]$	$\partial^3 \phi^5$	$[8 - \frac{5}{2}\varepsilon] + \frac{13}{6}\varepsilon$	$\varepsilon^2[96]$		5_5
$\text{Op}[0,3,3]$	$\partial^3 \square \phi^5$	$[10 - \frac{5}{2}\varepsilon] + \frac{25}{18}\varepsilon$	ε^1		13_5
$\text{Op}[0,3,4]$	$\partial^3 \phi^7$	$[10 - \frac{7}{2}\varepsilon] + \frac{11}{2}\varepsilon$	$\varepsilon^2[96]$		5_7
$\text{Op}[0,4,1]$	$\partial^4 \phi^3$	$[7 - \frac{3}{2}\varepsilon] + \frac{7}{15}\varepsilon$	$\varepsilon^2[7]$	6.112674(19)[97]	$3_4, 6_3$
$\text{Op}[0,4,2]$	$\partial^4 \phi^5$	$[9 - \frac{5}{2}\varepsilon] + \frac{14}{9}\varepsilon$	$\varepsilon^2[96]$		7_5
$\text{Op}[0,4,3]$	$\partial^4 \phi^5$	$[9 - \frac{5}{2}\varepsilon] + \frac{12}{5}\varepsilon$	$\varepsilon^2[96]$		6_5
$\text{Op}[0,5,1]$	$\partial^5 \phi^3$	$[8 - \frac{3}{2}\varepsilon] + \frac{2}{9}\varepsilon$	$\varepsilon^2[7]$	6.709778(27)[97]	$3_5, 8_3$
$\text{Op}[0,5,2]$	$\partial^5 \phi^5$	$[10 - \frac{5}{2}\varepsilon] + \frac{5}{6}\varepsilon$	ε^1		9_5
$\text{Op}[0,5,3]$	$\partial^5 \phi^5$	$[10 - \frac{5}{2}\varepsilon] + \frac{19}{9}\varepsilon$	ε^1		8_5
$\text{Op}[0,6,1]$	$\partial^6 \phi^3$	$[9 - \frac{3}{2}\varepsilon] + 0\varepsilon$	$\varepsilon^2[2, 99]$		
$\text{Op}[0,6,2]$	$\partial^6 \phi^3$	$[9 - \frac{3}{2}\varepsilon] + \frac{3}{7}\varepsilon$	$\varepsilon^2[7]$		3_6
$\text{Op}[0,7,1]$	$\partial^7 \phi^3$	$[10 - \frac{3}{2}\varepsilon] + \frac{1}{4}\varepsilon$	$\varepsilon^2[7]$	8.747293(56)[97]	3_7
$\text{Op}[0,8,1]$	$\partial^8 \phi^3$	$[11 - \frac{3}{2}\varepsilon] + 0\varepsilon$	$\varepsilon^2[2, 99]$		
$\text{Op}[0,8,2]$	$\partial^8 \phi^3$	$[11 - \frac{3}{2}\varepsilon] + \frac{11}{27}\varepsilon$	$\varepsilon^2[7]$		3_8
$\text{Op}[0,9,1]$	$\partial^9 \phi^3$	$[12 - \frac{3}{2}\varepsilon] + 0\varepsilon$	$\varepsilon^2[2, 99]$		
$\text{Op}[0,9,2]$	$\partial^9 \phi^3$	$[12 - \frac{3}{2}\varepsilon] + \frac{4}{15}\varepsilon$	$\varepsilon^2[7]$		3_9

New two-loop results for twist-3 operators can be computed by a two-loop dilatation operator, whose form was guessed based on analogy with [100], and confirmed in [99]. For these results we write [2, 99].

Note that this produces negative anomalous dimensions; the first instance is $\text{Op}[0,6,1]$ with

$$\Delta E[\text{Op}[0,6,1]] = \left[9 - \frac{3e}{2}\right] - \frac{e^2}{324} + \text{ord } e^2 \quad (3.14)$$

The data file has implemented these operators up to spin 13.

3.2 Operators in non-traceless-symmetric Lorentz representations

Table 15: Lorentz non-traceless-symmetric operators with $\Delta^{4d} \leq 10$. The 4d notation for the representations is the one used in [95]. In the data file, these operators have now been implemented, e.g. `DeltaE[Op[0,{2,2},1]]`

Lorentz YT	4d representation	Field content	$\Delta_{4-\varepsilon}^{(1)}$
	$(2, 0) + (0, 2)$	$\partial^4 \phi^3$	$[7 - \frac{3}{2}\varepsilon] + 0\varepsilon$
	$(\frac{5}{2}, \frac{3}{2}) + (\frac{3}{2}, \frac{5}{2})$	$\partial^5 \phi^3$	$[8 - \frac{3}{2}\varepsilon] + 0\varepsilon$
	$(2, 0) + (0, 2)$	$\partial^4 \phi^4$	$[8 - 2\varepsilon] + \frac{7}{9}\varepsilon$
	$(3, 1) + (1, 3)$	$\partial^6 \phi^3$	$[9 - \frac{3}{2}\varepsilon] + 0\varepsilon$
	$(\frac{5}{2}, \frac{1}{2}) + (\frac{1}{2}, \frac{5}{2})$	$\partial^5 \phi^4$	$[9 - 2\varepsilon] + \frac{2}{9}\varepsilon$
	$(\frac{3}{2}, \frac{1}{2}) + (\frac{1}{2}, \frac{3}{2})$	$\partial^5 \phi^4$	$[9 - 2\varepsilon] + \frac{1}{2}\varepsilon$
	$(\frac{5}{2}, \frac{3}{2}) + (\frac{3}{2}, \frac{5}{2})$	$\partial^5 \phi^4$	$[9 - 2\varepsilon] + \frac{13}{18}\varepsilon$
	$(2, 0) + (0, 2)$	$\partial^4 \phi^5$	$[9 - \frac{5}{2}\varepsilon] + \frac{17}{9}\varepsilon$
	$(\frac{7}{2}, \frac{3}{2}) + (\frac{3}{2}, \frac{7}{2})$	$\partial^7 \phi^3$	$[10 - \frac{3}{2}\varepsilon] + 0\varepsilon$
	$(\frac{7}{2}, \frac{5}{2}) + (\frac{5}{2}, \frac{7}{2})$	$\partial^7 \phi^3$	$[10 - \frac{3}{2}\varepsilon] + 0\varepsilon$
	$(2, 1) + (1, 2)$	$\partial^6 \phi^4$	$[10 - 2\varepsilon] + \frac{28}{45}\varepsilon$
	$(3, 1) + (1, 3)$	$\partial^6 \phi^4$	$[10 - 2\varepsilon] + \frac{37 - \sqrt{649}}{90}\varepsilon$
	$(3, 1) + (1, 3)$	$\partial^6 \phi^4$	$[10 - 2\varepsilon] + \frac{37 + \sqrt{649}}{90}\varepsilon$
	$(3, 2) + (2, 3)$	$\partial^6 \phi^4$	$[10 - 2\varepsilon] + \frac{13}{45}\varepsilon$
	$(\frac{3}{2}, \frac{1}{2}) + (\frac{1}{2}, \frac{3}{2})$	$\partial^5 \phi^5$	$[10 - \frac{5}{2}\varepsilon] + \frac{14}{9}\varepsilon$
	$(\frac{5}{2}, \frac{3}{2}) + (\frac{3}{2}, \frac{5}{2})$	$\partial^5 \phi^5$	$[10 - \frac{5}{2}\varepsilon] + \frac{16}{9}\varepsilon$
	$(\frac{5}{2}, \frac{1}{2}) + (\frac{1}{2}, \frac{5}{2})$	$\partial^5 \phi^5$	$[10 - \frac{5}{2}\varepsilon] + \frac{23}{18}\varepsilon$
	$(2, 0) + (0, 2)$	$\partial^4 \phi^6$	$[10 - 3\varepsilon] + \frac{10}{3}\varepsilon$

Reference [95] gives the family of operators with k fields and Lorentz representation, for $k \geq 3$. For this family, $\Delta = k[1 - \frac{\varepsilon}{2}] + \frac{1}{3}(1/2k(k-1) - 2/3k - 1)\varepsilon + O(\varepsilon^2)$.

3.2.1 Three-row Lorentz Young tableaux

We only comment on operators in the representation given by the Lorentz Young tableau $y_{1,1,1} = \begin{smallmatrix} \square \\ \square \\ \square \end{smallmatrix}$, since they become parity-odd scalars in three dimensions. For $N = 1$ and $d = 3$, the theory of a free scalar field has an operator the form $\epsilon^{\mu\nu\rho} \partial_\mu \partial_\nu \partial_\rho \square^3 \phi^4$.

For the interacting theory, the corresponding operator is the leading operator in the $y_{1,1,1}$ Lorentz representation, which has dimension $[13 - 2\varepsilon] + 0\varepsilon + O(\varepsilon^2)$. In [101], the numerical conformal bootstrap studied the four-point function of stress-tensors in three dimensions. A

general upper bound for the parity-odd scalar was found: $\Delta_{\mathcal{O}_{\text{odd},\text{min}}} \lesssim 11.5$. When inserting some input to single out the Ising model, a somewhat lower bound was found: $\Delta_{\mathcal{O}_{\text{odd},\text{min}}} < 11.2$. Reference [23] found an upper bound $\Delta_{\mathcal{O}_{\text{odd},\text{min}}} \leq 10.9293$ within the Ising bootstrap island.

3.3 OPE coefficients

See tables 16, 17 and 18 for OPE coefficients in the $\phi \times \phi$, $\phi \times \phi^2$ and $\phi^2 \times \phi^2$ OPEs.

Several OPE coefficients involving the stress tensor were computed in [24].

Table 16: Squared OPE coefficients in $\phi \times \phi$ for operators for $N = 1$. Numerical values are taken from [97] and denote non-squared OPE coefficients.

$\phi, \phi, \mathcal{O}$	$\mathcal{O} \varepsilon$	$\lambda_{\phi\phi\mathcal{O}}^2(\varepsilon)$	$\lambda_{\phi\phi\mathcal{O}}^2(\varepsilon)$	$\lambda_{\phi\phi\mathcal{O}}^{3d}$
$\phi, \phi, \text{Op}[\text{E}, 0, 1]$	ϕ^2	2	$\varepsilon^3[\text{102}]\varepsilon^4(\text{num})[\text{103}]$	1.0518537(41)
$\phi, \phi, \text{Op}[\text{E}, 0, 2]$	ϕ^4	$\frac{1}{54}\varepsilon^2$	$\varepsilon^2[\text{104}]\varepsilon^3[\text{103}]$	0.053012(55)*
$\phi, \phi, \text{Op}[\text{E}, 0, 3]$	ϕ^6	$\frac{5}{104976}\varepsilon^4$	$\varepsilon^4[\text{105}]$	0.0007338(31)
$\phi, \phi, \text{Op}[\text{E}, 0, 4]$	$\square^2\phi^4$			0.000162(12)
$\phi, \phi, \text{Op}[\text{E}, 2, 1]$	$T^{\mu\nu}$	$\frac{1}{3}$	$\varepsilon^3[\text{102}]\varepsilon^4[\text{106}]$	0.32613776(45) [†]
$\phi, \phi, \text{Op}[\text{E}, 2, 2]$	$\partial^2\phi^4$	$\frac{1}{1440}\varepsilon^2$	$\varepsilon^2[\text{106}]\varepsilon^3[\text{7}]$	0.0105745(42) [‡]
$\phi, \phi, \text{Op}[\text{E}, 2, 3]$	$\partial^2\square\phi^4$	$\frac{\varepsilon^4}{2592000}$	$\varepsilon^4[\text{7}]$	0.0004773(62)
$\phi, \phi, \text{Op}[\text{E}, 4, 1]$	$C^{\mu\nu\rho\sigma}$	$\frac{1}{35}$	$\varepsilon^3[\text{102}]\varepsilon^4[\text{106}]$	0.069076(43)
$\phi, \phi, \text{Op}[\text{E}, 4, 2]$	$\partial^4\phi^4$	$\frac{1}{419580}\varepsilon^2$	$\varepsilon^2[\text{1}, \text{106}]$	0.0019552(12)
$\phi, \phi, \text{Op}[\text{E}, 4, 3]$	$\partial^4\phi^4$	$\frac{1}{23976}\varepsilon^2$	$\varepsilon^2[\text{1}, \text{106}]$	0.00237745(44)
$\phi, \phi, \text{Op}[\text{E}, 6, 1]$	$\mathcal{I}_6$	$\frac{1}{462}$	$\varepsilon^3[\text{102}]\varepsilon^4[\text{106}]$	0.0157416(41)
$\phi, \phi, \text{Op}[\text{E}, 6, 2]$	$\partial^6\phi^4$	$O(\varepsilon^2)$	$\varepsilon^2[\text{1}, \text{106}]$	
$\phi, \phi, \text{Op}[\text{E}, 6, 3]$	$\partial^6\phi^4$	$O(\varepsilon^2)$	$\varepsilon^2[\text{1}, \text{106}]$	
$\phi, \phi, \text{Op}[\text{E}, 6, 4]$	$\partial^6\phi^4$	$O(\varepsilon^2)$	$\varepsilon^2[\text{1}, \text{106}]$	
$\phi, \phi, \text{Op}[\text{E}, 8, 1]$	$\mathcal{I}_8$	$\frac{1}{6435}$	$\varepsilon^3[\text{102}]\varepsilon^4[\text{106}]$	0.0036850(54)
$\phi, \phi, \text{Op}[\text{E}, 10, 1]$	$\mathcal{I}_{10}$	$\frac{1}{92378}$	$\varepsilon^3[\text{102}]\varepsilon^4[\text{106}]$	0.00087562(13)

*A value with rigorous error bars, 0.05304(16), was given in [78].

[†]Related to the central charge C_T .

[‡]A value with rigorous error bars, 0.01054(10), was given in [78].

Table 17: Squared OPE coefficients in $\phi \times \phi^2$ for operators for $N = 1$. Numerical values are from [97] and denote non-squared OPE coefficients.

$\phi, \phi^2, \mathcal{O}$	$\mathcal{O} \varepsilon$	$\lambda_{\phi\phi^2\mathcal{O}}^2(\varepsilon)$	$\lambda_{\phi\phi^2\mathcal{O}}^2(\varepsilon)$	$\lambda_{\phi\phi^2\mathcal{O}}^{3d}$
$\phi, \phi^2, \text{Op}[0, 0, 1]$	ϕ	2	$\varepsilon^3[102]\varepsilon^4(\text{num})[103]$	1.0518537(41)
$\phi, \phi^2, \text{Op}[0, 0, 2]$	ϕ^5	$\frac{5}{108}\varepsilon^2$	$\varepsilon^2[105]$	0.057235(20)*
$\phi, \phi^2, \text{Op}[0, 2, 1]$	$\partial^2\phi^3$	$\frac{1}{2}$	$\varepsilon^1[7]$	0.38915941(81)
$\phi, \phi^2, \text{Op}[0, 2, 2]$	$\partial^2\phi^5$	$O(\varepsilon^2)$		0.017413(73)
$\phi, \phi^2, \text{Op}[0, 3, 1]$	$\partial^3\phi^3$	$\frac{2}{35}$	$\varepsilon^1[7]$	0.1385(34)
$\phi, \phi^2, \text{Op}[0, 4, 1]$	$\partial^4\phi^3$	$\frac{1}{18}$	$\varepsilon^1[7]$	0.1077052(16)
$\phi, \phi^2, \text{Op}[0, 5, 1]$	$\partial^5\phi^3$	$\frac{2}{231}$	$\varepsilon^1[7]$	0.04191549(88)
$\phi, \phi^2, \text{Op}[0, 6, 1]$	$\partial^6\phi^3$	$O(\varepsilon^2)$		
$\phi, \phi^2, \text{Op}[0, 6, 2]$	$\partial^6\phi^3$	$\frac{3}{572}$	$\varepsilon^1[7]$	
$\phi, \phi^2, \text{Op}[0, 7, 1]$	$\partial^7\phi^3$	$\frac{2}{2145}$	$\varepsilon^1[7]$	0.01161255(13)
$\phi, \phi^2, \text{Op}[0, 8, 1]$	$\partial^8\phi^3$	$O(\varepsilon^2)$		
$\phi, \phi^2, \text{Op}[0, 8, 2]$	$\partial^8\phi^3$	$\frac{1}{2210}$	$\varepsilon^1[7]$	
$\phi, \phi^2, \text{Op}[0, 9, 1]$	$\partial^9\phi^3$	$O(\varepsilon^2)$		
$\phi, \phi^2, \text{Op}[0, 9, 2]$	$\partial^9\phi^3$	$\frac{4}{46189}$	$\varepsilon^1[7]$	

*A value with rigorous error bars, 0.0565(15), was given in [78].

Table 18: Squared OPE coefficients in $\phi^2 \times \phi^2$ for operators for $N = 1$. Numerical values are from [97] and denote non-squared OPE coefficients.

$\phi^2, \phi^2, \mathcal{O}$	$\mathcal{O} \varepsilon$	$\lambda_{\phi^2\phi^2\mathcal{O}}^2(\varepsilon)$	$\lambda_{\phi^2\phi^2\mathcal{O}}^2(\varepsilon)$	$\lambda_{\phi^2\phi^2\mathcal{O}}^{3d}$
$\phi^2, \phi^2, \mathbf{0p}[E, 0, 0]$	$\mathbb{1}$	1	exact	1
$\phi^2, \phi^2, \mathbf{0p}[E, 0, 1]$	ϕ^2	8	$\varepsilon^1[1, 107]$	1.532435(19)
$\phi^2, \phi^2, \mathbf{0p}[E, 0, 2]$	ϕ^4	6	$\varepsilon^1[1, 107]$	1.5360(16)*
$\phi^2, \phi^2, \mathbf{0p}[E, 0, 3]$	ϕ^6			0.1279(17)
$\phi^2, \phi^2, \mathbf{0p}[E, 0, 4]$	$\square^2\phi^4$	$\frac{1}{6}$	$\varepsilon^1[1, 107]$	0.1874(31)
$\phi^2, \phi^2, \mathbf{0p}[E, 2, 1]$	$T^{\mu\nu}$	$\frac{4}{3}$	$\varepsilon^4[106]$	0.8891471(40)
$\phi^2, \phi^2, \mathbf{0p}[E, 2, 2]$	$\partial^2\phi^4$	$\frac{8}{5}$	$\varepsilon^1[1, 107]$	0.69023(49) [†]
$\phi^2, \phi^2, \mathbf{0p}[E, 2, 3]$	$\partial^2\square\phi^4$	$\frac{1}{5}$	$\varepsilon^1[1, 107]$	0.21882(73)
$\phi^2, \phi^2, \mathbf{0p}[E, 2, 4]$	$\partial^2\phi^6$			
$\phi^2, \phi^2, \mathbf{0p}[E, 2, 5]$	$\partial^2\square^2\phi^4$	$\frac{1}{999}$	$\varepsilon^1[1, 7, 107]$	
$\phi^2, \phi^2, \mathbf{0p}[E, 2, 6]$	$\partial^2\square^2\phi^4$	$\frac{4}{111}$	$\varepsilon^1[1, 7, 107]$	
$\phi^2, \phi^2, \mathbf{0p}[E, 4, 1]$	$C^{\mu\nu\rho\sigma}$	$\frac{4}{35}$	$\varepsilon^1[1, 107]\varepsilon^2[7]$	0.24792(20)
$\phi^2, \phi^2, \mathbf{0p}[E, 4, 2]$	$\partial^4\phi^4$	$\frac{125}{2331}$	$\varepsilon^1[1, 107]$	-0.110247(54)
$\phi^2, \phi^2, \mathbf{0p}[E, 4, 3]$	$\partial^4\phi^4$	$\frac{8}{37}$	$\varepsilon^1[1, 107]$	0.22975(10)
$\phi^2, \phi^2, \mathbf{0p}[E, 4, 4]$	$\partial^4\square\phi^4$	$\frac{2091-71\sqrt{697}}{107338}$	$\varepsilon^1[1, 7, 107]$	0.08635(18)
$\phi^2, \phi^2, \mathbf{0p}[E, 4, 5]$	$\partial^4\square\phi^4$	$\frac{2091+71\sqrt{697}}{107338}$	$\varepsilon^1[1, 7, 107]$	
$\phi^2, \phi^2, \mathbf{0p}[E, 6, 1]$	$\mathcal{J}_6$	$\frac{2}{231}$	$\varepsilon^1[1, 107]\varepsilon^2[7]$	0.066136(36)
$\phi^2, \phi^2, \mathbf{0p}[E, 8, 1]$	$\mathcal{J}_8$	$\frac{4}{6435}$	$\varepsilon^1[1, 107]\varepsilon^2[7]$	0.017318(30)
$\phi^2, \phi^2, \mathbf{0p}[E, 10, 1]$	$\mathcal{J}_{10}$	$\frac{2}{46189}$	$\varepsilon^1[1, 107]\varepsilon^2[7]$	0.0044811(15)

*A value with rigorous error bars, 1.5362(12), was given in [78].

[†]A value with rigorous error bars, 0.678(15), was given in [78].

3.4 Families of operators for general N

For general N we introduce a number of families $\text{ONF}\langle i \rangle$. We express the $1/N$ part of the large N operator dimension in terms of the anomalous dimension of φ : $\Delta_\varphi = \mu - 1 + \frac{\eta_1/2}{N} + O(N^{-2})$.

ONF1 $[k]$ Scalar singlet operators of the form $\varphi_S^{2k} \sim \sigma^k$, $k \geq 1$. We have ε^1 [108] ε^2 [90] ε^3 [109] and N^{-1} [71, 110] N^{-2} [90],

$$\text{DeltaE}[\text{ONF1}[k]] = k[2 - \varepsilon] + \frac{k(6k + N - 4)}{N + 8}\varepsilon + \dots + O(\varepsilon^3), \quad (3.15)$$

$$\text{DeltaN}[\text{ONF1}[k]] = 2k + \frac{2k(2\mu - 1)}{\mu - 2} (k\mu(2\mu - 3) - \mu^2 + \mu + 2) \frac{\eta_1/2}{N} + \dots + O(1/N^3). \quad (3.16)$$

ONF2 $[k]$ Scalar fundamental operators of the form $\varphi_V^{2k+1} \sim \sigma^k \varphi$, $k \geq 0$, $k \neq 1$. We have ε^1 [108] and N^{-1} [66],

$$\text{DeltaE}[\text{ONF2}[k]] = (2k + 1)[1 - \frac{\varepsilon}{2}] + \frac{k(6k + N + 2)}{N + 8}\varepsilon + O(\varepsilon^2), \quad (3.17)$$

$$\text{DeltaN}[\text{ONF2}[k]] = 2k + \mu - 1 + \left(\frac{2k(k - 1)\mu(2\mu - 3)(2\mu - 1)}{2 - \mu} + 1 \right) \frac{\eta_1/2}{N} + O(N^{-2}). \quad (3.18)$$

In fact **ONF1** and **ONF2** have the largest order ε anomalous dimensions for an operator with $2k$ and $2k + 1$ fields respectively [94].

ONF3 $[m]$ Scalar completely symmetric traceless tensors of the form $\varphi_{T_m}^m$, $m \geq 1$. We have ε^2 [111] ε^4 [112] ε^5 [1, 60, 112] ε^6 [113] and N^{-1} [114, 115] N^{-2} [62],

$$\text{DeltaE}[\text{ONF3}[m]] = m \left[1 - \frac{1}{2}\varepsilon \right] + \frac{m(m - 1)}{N + 8}\varepsilon + \dots + O(\varepsilon^6), \quad (3.19)$$

$$\text{DeltaN}[\text{ONF3}[m]] = m(\mu - 1) + \frac{m(m\mu + 2 - 2\mu)}{\mu - 2} \frac{\eta_1/2}{N} + \dots + O(N^{-3}). \quad (3.20)$$

ONF4 $[l]$ Singlet broken currents of the form $\mathcal{J}_{S,\ell} = [\varphi, \varphi]_{S,0,\ell}$, $\ell \in 2\mathbb{Z}_+$. We have ε^2 [92] ε^4 [116] and N^{-1} [110] N^{-2} [116],

$$\text{DeltaE}[\text{ONF4}[l]] = 2 - \varepsilon + l + \frac{N + 2}{2(N + 8)^2} \left(1 - \frac{6}{l(l + 1)} \right) \varepsilon^2 + \dots + O(\varepsilon^5), \quad (3.21)$$

$$\text{DeltaN}[\text{ONF4}[l]] = 2(\mu - 1) + l + \left(2 - \frac{2\mu \left((\mu - 1) + \frac{\Gamma(2\mu - 1)\Gamma(l + 1)}{\Gamma(l + 2\mu - 3)} \right)}{(l + \mu - 1)(l + \mu - 2)} \right) \frac{\eta_1/2}{N} + \dots + O(N^{-3}). \quad (3.22)$$

ONF5 [l] Traceless-symmetric broken currents of the form $\mathcal{J}_{T,\ell} = [\varphi, \varphi]_{T,0,\ell}$, $\ell \in 2\mathbb{Z}_+$. We have ε^2 [92] ε^3 [117] ε^4 [65] and N^{-1} [110] N^{-2} [62],

$$\text{DeltaE}[\text{ONF5}[l]] = 2 - \varepsilon + l + \frac{1}{2(N+8)^2} \left(N + 2 - \frac{2(N+6)}{l(l+1)} \right) \varepsilon^2 + \dots + O(\varepsilon^5), \quad (3.23)$$

$$\text{DeltaN}[\text{ONF5}[l]] = 2(\mu - 1) + l + \left(2 - \frac{2\mu(\mu - 1)}{(l + \mu - 1)(l + \mu - 2)} \right) \frac{\eta_1/2}{N} + \dots + O(N^{-3}). \quad (3.24)$$

ONF6 [l] Antisymmetric broken currents of the form $\mathcal{J}_{A,\ell} = [\varphi, \varphi]_{A,0,\ell}$, $\ell \in 2\mathbb{N} + 1$. We have ε^3 [117] ε^4 [65] and N^{-1} [110] N^{-2} [62],

$$\text{DeltaE}[\text{ONF6}[l]] = 2 - \varepsilon + l + \frac{N+2}{2(N+8)^2} \left(1 - \frac{2}{l(l+1)} \right) \varepsilon^2 + \dots + O(\varepsilon^5), \quad (3.25)$$

$$\text{DeltaN}[\text{ONF6}[l]] = 2(\mu - 1) + l + \left(2 - \frac{2\mu(\mu - 1)}{(l + \mu - 1)(l + \mu - 2)} \right) \frac{\eta_1/2}{N} + \dots + O(N^{-3}). \quad (3.26)$$

ONF7 [l] Spinning singlet operators of the form $[\sigma, \sigma]_{0,\ell}$, $\ell \in 2\mathbb{N}$. We have N^{-1} [110],

$$\begin{aligned} \text{DeltaN}[\text{ONF7}[l]] = & 4 + l - \frac{8}{(2 - \mu)^2} \left((\mu - 1)(2 - \mu)(2\mu - 1) \right. \\ & \left. + \frac{\mu(2\mu - 3)}{(l + 1)(l + 2)} \left((\mu - 1)(2\mu - 3) - \frac{(\mu - l - 3)_l}{(\mu)_l} \right) \right) \frac{\eta_1/2}{N} + O(N^{-2}), \end{aligned} \quad (3.27)$$

where $(a)_k = \frac{\Gamma(a+k)}{\Gamma(a)}$ is the (rising) Pochhammer symbol. There is no simple expression for these operators in the ε -expansion.

ONF8 [l] Spinning fundamental operators of the form $[\varphi, \sigma]_{0,\ell}$, $\ell \geq 1$. We have ε^1 [94] and N^{-1} [110],

$$\text{DeltaE}[\text{ONF8}[l]] = [3 - \frac{3}{2}\varepsilon] + l \quad (3.28)$$

$$+ \frac{4(-1)^l + (N+4)(l+1) + \sqrt{16(l+1)(-1)^l + 48 + 16N + (l+1)^2 N^2}}{2(N+8)(l+1)} \varepsilon + O(\varepsilon^2),$$

$$\begin{aligned} \text{DeltaN}[\text{ONF8}[l]] = & \mu + 1 + l + \left(\frac{4\mu(\mu - 1)(2\mu - 3)}{(l + 1)(l + \mu - 1)(2 - \mu)} \right. \\ & \left. - \frac{8\mu^2 - 11\mu + 2}{2 - \mu} + \frac{2(-1)^l \Gamma(l+1) \Gamma(\mu+1)}{(2 - \mu) \Gamma(l + \mu)} \right) \frac{\eta_1/2}{N} + O(N^{-2}). \end{aligned} \quad (3.29)$$

ONF9 $[k]$ Scalar singlet operators of the form $\Box\varphi_S^{2k} \sim \Box\sigma^k$, $k \geq 2$. We have ε^1 [1] and N^{-1} [71],

$$\text{DeltaE}[\text{ONF9}[k]] = 2 + k(2 - \varepsilon) + \frac{6k^2 + (N - 12)k + 4}{N + 8}\varepsilon + O(\varepsilon^2), \quad (3.30)$$

$$\begin{aligned} \text{DeltaN}[\text{ONF9}[k]] &= 2k + 2 + \frac{2(2\mu - 1)}{3(2 - \mu)}(3k^2\mu(2\mu - 3) - k(10\mu^2 - \mu - 18) \\ &\quad + 2\mu^3 - \mu^2 - 7\mu + 6)\frac{\eta_1/2}{N} + O(N^{-2}). \end{aligned} \quad (3.31)$$

ONF10 $[k, m]$ Scalar operators of the form $\varphi_{T_m}^{2k+m} \sim \sigma^k \varphi_{T_m}^m$, $k, m \geq 0$. We have ε^1 [108] and N^{-1} [66],

$$\text{DeltaE}[\text{ONF10}[k, m]] = (2k + m)[1 - \frac{\varepsilon}{2}] + \frac{6k^2 + k(6m + N - 4) + m(m - 1)}{N + 8}\varepsilon + O(\varepsilon^2), \quad (3.32)$$

$$\begin{aligned} \text{DeltaN}[\text{ONF10}[k, m]] &= 2k + m(\mu - 1) + \frac{1}{2 - \mu}[2k^2\mu(2\mu - 1)(2\mu - 3) + m(m\mu + 2 - 2\mu) \\ &\quad - 2k(2\mu - 1)(2\mu^2 - \mu - 2 - 2m(\mu - 1))]\frac{\eta_1/2}{N} + O(N^{-2}). \end{aligned} \quad (3.33)$$

For $m = 0$ the family reduces to **ONF1** $[k]$, for $m = 1$ to **ONF2** $[k]$ and for $k = 0$ to **ONF3** $[m]$.

ONF11 $[m, l]$ Spin- l operators of the form $\partial^l \varphi_{Y_{m,l}}^{m+l}$ transforming in the $O(N)$ representation $Y_{m,l}$. We have ε^1 [1] and N^{-1} [115].

$$\text{DeltaE}[\text{ONF11}[m, l]] = (m + l)[1 - \frac{\varepsilon}{2}] + l + \frac{3m^2 + 3ml - 3m + l^2 - 4l}{3(N + 8)}\varepsilon + O(\varepsilon^2), \quad (3.34)$$

$$\begin{aligned} \text{DeltaN}[\text{ONF11}[m, l]] &= (m + l)(\mu - 1) + l \\ &\quad + \left[(m + l)(m + l - 2) - 4(l - 1) \left(\frac{m}{2\mu} + \frac{l}{2\mu + 2} \right) \right] \frac{\mu}{2 - \mu} \frac{\eta_1/2}{N} + O(N^{-2}). \end{aligned} \quad (3.35)$$

For $l = 0$ the family reduces to **ONF3** $[m]$. Note that $Y_{m,0} = T_m$, and $Y_{m,1} = H_{m+1}$, and that $Y_{2,2} = B_4$. Note also that operators with $m + l$ fields in the $O(N)$ representation $Y_{m,l}$ and with no contracted derivatives must have spin $\geq l$.

ONF12 $[m]$ Spin-2 operators of the form $\partial^2 \varphi_{T_m}^m$ transforming in the rank- m traceless-symmetric $O(N)$ representation, $m \geq 2$. We have ε^2 [2, 100] and N^{-1} [2, 118]

$$\text{DeltaE}[\text{ONF12}[m]] = m[1 - \frac{\varepsilon}{2}] + 2 + \frac{3m^2 - 5m - 2}{3(N + 8)}\varepsilon + \dots + O(\varepsilon^3), \quad (3.36)$$

$$\text{DeltaN}[\text{ONF12}[m]] = m(\mu - 1) + 2 + \frac{m^2\mu(\mu + 1) - 2m(\mu^2 + 2\mu - 3) - 4}{(\mu + 1)(2 - \mu)} \frac{\eta_1/2}{N} + O(N^{-2}). \quad (3.37)$$

For $l = 0$ the family reduces to $\text{ONF3}[m]$. Note that $Y_{m,0} = T_m$, and $Y_{m,1} = H_{m+1}$, and that $Y_{2,2} = B_4$. Note also that operators with $m + l$ fields in the $O(N)$ representation $Y_{m,l}$ and with no contracted derivatives must have spin $\geq l$.

Apart from the **eleven twelve** families of operators for generic N listed above, [94] gives three additional families in the ε -expansion. These families are only defined by the fact that they acquire an order ε anomalous dimension, which singles them out from additional operators of the same type. This property is not visible in the large N expansion, whence we do not include them in our list of families.

- $\partial^\ell \varphi_V^3$ with non-zero order ε dimension ε^1 [94]. This is the sister family to $\text{ONF8}[\ell]$. Dimension given by (3.28), but with a minus sign in front of the square root.
- $\partial^\ell \varphi_{T_3}^3$ with non-zero order ε dimension ε^1 [94]: , $\ell \neq 1$. By computing many of these operator dimensions using [100], we have **guessed a general two-loop formula** [2]:

$$\begin{aligned} \Delta_\ell = & \left[3 - \frac{3}{2}\varepsilon\right] + \ell + \frac{2}{N+8} \left(1 + \frac{2(-1)^\ell}{\ell+1}\right) \varepsilon + \varepsilon^2 \left[- \frac{N^2 - 46N - 192}{4(N+8)^3} \right. \\ & + \frac{4(N+6)(\ell^3 - (\ell+1)^3 S_1(\ell)) + 2\ell^2(5N+26) + 4\ell(3N+14) + 6N+60}{(N+8)^2(\ell+3)(\ell+1)^3(\ell-1)} \\ & - 2(-1)^\ell \left(\frac{\ell^3(3N^2+28N+92) + \ell^2(7N^2+54N+164) - \ell(2N^2+18N+76) + 48N+204}{(N+8)^3(\ell+3)(\ell+1)^2(\ell-1)} \right. \\ & \left. \left. - \frac{(N+8)(\ell^2+2\ell)+N}{(N+8)^2(\ell+3)(\ell+1)(\ell-1)} S_1(\ell) \right) \right] + O(\varepsilon^3) \end{aligned} \quad (3.38)$$

- $\partial^\ell \varphi_{H_3}^3$ with non-zero order ε dimension ε^1 [94]: $\left[3 - \frac{3}{2}\varepsilon\right] + \ell + \frac{1}{N+8} \left(2 - \frac{2(-1)^\ell}{\ell+1}\right) \varepsilon + O(\varepsilon^2)$, $\ell \neq 0$.

Moreover, in [66], a closed-form expression at large N is given for the anomalous dimension of spin-one operators of the form $\partial\sigma^k\varphi$.

Table 19: Singlet scalar operators for $N > 1$. $N_{\dots}^0$ denotes $N^0 + O(N^{-1})$, where there is an exact expression.

$\mathcal{O}$	$\mathcal{O} \varepsilon$	$\mathcal{O} N$	$\Delta_{4-\varepsilon}^{(1)}$	$\Delta(\varepsilon)$	$\Delta(N)$	Family
$0p[S, 0, 1]$	φ_S^2	σ	$[2 - \varepsilon] + \frac{N+2}{N+8}\varepsilon$	$\varepsilon^7[40]$	$N^{-2}[49]$	$1_2, 10_{1,0}$
$0p[S, 0, 2]$	φ_S^4	σ^2	$[4 - 2\varepsilon] + 2\varepsilon$	$\varepsilon^7[40]$	$N^{-2}[51]$	$1_2, 7_0, 10_{2,0}$
$0p[S, 0, 3]$	$\square\varphi_S^4$	$\square\sigma^2$	$[6 - 2\varepsilon] + \frac{2N+4}{N+8}\varepsilon$	$\varepsilon^2[119]\varepsilon^4[4]\varepsilon^5[5]$	$N^{-1}[71]$	9_2
$0p[S, 0, 4]$	φ_S^6	σ^3	$[6 - 3\varepsilon] + \frac{3N+42}{N+8}\varepsilon$	$\varepsilon^2[90]\varepsilon^4[4]\varepsilon^5[5]$	$N^{-2}[90]$	$1_3, 10_{3,0}$
$0p[S, 0, 5]$	$\square^2\varphi_S^4$	$\square^2\varphi_S^4$	$[8 - 2\varepsilon] + 0\varepsilon N_{\dots}^0$	$\varepsilon^3[4]\varepsilon^4[5]$		
$0p[S, 0, 6]$	$\square^2\varphi_S^4$	$\square^2\sigma^2$	$[8 - 2\varepsilon] + 2\varepsilon N_{\dots}^0$	$\varepsilon^3[4]\varepsilon^4[5]$		
$0p[S, 0, 7]$	$\square\varphi_S^6$	$\square\sigma^3$	$[8 - 3\varepsilon] + \frac{3N+22}{N+8}\varepsilon$	$\varepsilon^3[4]\varepsilon^4[5]$	$N^{-1}[71]$	9_3
$0p[S, 0, 8]$	φ_S^8	σ^4	$[8 - 4\varepsilon] + \frac{4N+80}{N+8}\varepsilon$	$\varepsilon^2[90]\varepsilon^3[4]\varepsilon^4[5]$	$N^{-2}[90]$	$1_4, 10_{4,0}$
$0p[S, 0, 9]$	$\square^3\varphi_S^4$		$[10 - 2\varepsilon] + 0\varepsilon N_{\dots}^0$	$\varepsilon^2[5]$		
$0p[S, 0, 10]$	$\square^2\varphi_S^6$		$[10 - 3\varepsilon] + \varepsilon N_{\dots}^0$	$\varepsilon^2[5]$		
$0p[S, 0, 11]$	$\square^2\varphi_S^6$		$[10 - 3\varepsilon] + 2\varepsilon N_{\dots}^0$	$\varepsilon^2[5]$		
$0p[S, 0, 12]$	$\square^2\varphi_S^6$		$[10 - 3\varepsilon] + 3\varepsilon N_{\dots}^0$	$\varepsilon^2[5]$		
$0p[S, 0, 13]$	$\square^3\varphi_S^4$		$[10 - 2\varepsilon] + 2\varepsilon N_{\dots}^0$	$\varepsilon^2[5]$		
$0p[S, 0, 14]$	$\square^2\varphi_S^6$		$[10 - 3\varepsilon] + 3\varepsilon N_{\dots}^0$	$\varepsilon^2[5]$		
$0p[S, 0, 15]$	$\square\varphi_S^8$	$\square\sigma^4$	$[10 - 4\varepsilon] + \frac{4N+13}{N+8}\varepsilon$	$\varepsilon^2[5]$	$N^{-1}[71]$	9_4
$0p[S, 0, 16]$	φ_S^{10}	σ^5	$[10 - 5\varepsilon] + \frac{5N+130}{N+8}\varepsilon$	$\varepsilon^2[90]$	$N^{-2}[90]$	$1_5, 10_{5,0}$

3.4.1 Spinning singlet operators

Table 20: Singlet spinning operators for $N > 1$.

$\mathcal{O}$	$\mathcal{O} _{\varepsilon}$	$\mathcal{O} _N$	$\Delta_{4-\varepsilon}^{(1)}$	$\Delta(\varepsilon)$	$\Delta(N)$	Family
0p[S, 1, 1]	$\partial\Box\varphi_S^6$	$\partial\Box\sigma^3$	$[9 - 3\varepsilon] + \frac{9N+58}{3(N+8)}\varepsilon$			
0p[S, 2, 1]	$T^{\mu\nu}$	$T^{\mu\nu}$	$[4 - \varepsilon]$	exact	exact	4_2
0p[S, 2, 2]	$\partial^2\varphi_S^4$	$[\varphi, \varphi]_{S,1,2}$	$[6 - 2\varepsilon] + \varepsilon N_{\dots}^0$	$\varepsilon^5[5]$		
0p[S, 2, 3]	$\partial^2\varphi_S^4$	$[\sigma, \sigma]_{0,2}$	$[6 - 2\varepsilon] + 2\varepsilon N_{\dots}^0$	$\varepsilon^5[5]$	$N^{-1}[110]$	7_2
0p[S, 2, 4]	$\partial^2\Box\varphi_S^4$		$[8 - 2\varepsilon] + 0\varepsilon N_{\dots}^0$	ε^1		
0p[S, 2, 5]	$\partial^2\Box\varphi_S^4$		$[8 - 2\varepsilon] + \varepsilon N_{\dots}^0$	ε^1		
0p[S, 2, 6]	$\partial^2\varphi_S^6$		$[8 - 3\varepsilon] + 2\varepsilon N_{\dots}^0$	ε^1		
0p[S, 2, 7]	$\partial^2\Box\varphi_S^4$		$[8 - 2\varepsilon] + 2\varepsilon N_{\dots}^0$	ε^1		
0p[S, 2, 8]	$\partial^2\varphi_S^6$		$[8 - 3\varepsilon] + 3\varepsilon N_{\dots}^0$	ε^1		
0p[S, 3, 1]	$\partial^3\Box\varphi_S^4$		$[9 - 2\varepsilon] + \frac{n+2}{n+8}\varepsilon$	ε^1		
0p[S, 3, 2]	$\partial^3\varphi_S^6$		$[9 - 3\varepsilon] + 2\varepsilon N_{\dots}^0$	ε^1		
0p[S, 3, 3]	$\partial^3\varphi_S^6$		$[9 - 3\varepsilon] + 3\varepsilon N_{\dots}^0$	ε^1		
0p[S, 4, 1]	$\mathcal{I}_{S,4}$	$\mathcal{I}_{S,4}$	$[6 - \varepsilon]$	$\varepsilon^4[116]$	$N^{-2}[116]$	4_4
0p[S, 4, 2]	$\partial^4\varphi_S^4$	$\partial^4\varphi_S^4$	$[8 - 2\varepsilon] + 0\varepsilon N_{\dots}^0$	ε^1		
0p[S, 4, 3]	$\partial^4\varphi_S^4$		$[8 - 2\varepsilon] + \varepsilon N_{\dots}^0$	ε^1		
0p[S, 4, 4]	$\partial^4\varphi_S^4$		$[8 - 2\varepsilon] + \varepsilon N_{\dots}^0$	ε^1		
0p[S, 4, 5]	$\partial^4\varphi_S^4$	$[\sigma, \sigma]_{0,4}$	$[8 - 2\varepsilon] + 2\varepsilon N_{\dots}^0$	ε^1	$N^{-1}[110]$	7_4
0p[S, 5, 1]	$\partial^5\varphi_S^4$		$[9 - 2\varepsilon] + \frac{N+2}{N+8}\varepsilon$	ε^1		
0p[S, 6, 1]	$\mathcal{I}_{S,6}$	$\mathcal{I}_{S,6}$	$[8 - \varepsilon]$	$\varepsilon^4[116]$	$N^{-2}[116]$	4_6
0p[S, 6, 2]	$\partial^6\varphi_S^4$	$\partial^6\varphi_S^4$	$[10 - 2\varepsilon] + 0\varepsilon N_{\dots}^0$	ε^1		
0p[S, 6, 3]	$\partial^6\varphi_S^4$	$\partial^6\varphi_S^4$	$[10 - 2\varepsilon] + 0\varepsilon N_{\dots}^0$	ε^1		
0p[S, 6, 4]	$\partial^6\varphi_S^4$		$[10 - 2\varepsilon] + \varepsilon N_{\dots}^0$	ε^1		
0p[S, 6, 5]	$\partial^6\varphi_S^4$		$[10 - 2\varepsilon] + \varepsilon N_{\dots}^0$	ε^1		
0p[S, 6, 6]	$\partial^6\varphi_S^4$		$[10 - 2\varepsilon] + \varepsilon N_{\dots}^0$	ε^1		
0p[S, 6, 7]	$\partial^6\varphi_S^4$	$[\sigma, \sigma]_{0,6}$	$[10 - 2\varepsilon] + 2\varepsilon N_{\dots}^0$	ε^1	$N^{-1}[110]$	7_6
0p[S, 8, 1]	$\mathcal{I}_{S,8}$	$\mathcal{I}_{S,8}$	$[10 - \varepsilon]$	$\varepsilon^4[116]$	$N^{-2}[116]$	4_8
0p[S, 10, 1]	$\mathcal{I}_{S,10}$	$\mathcal{I}_{S,10}$	$[12 - \varepsilon]$	$\varepsilon^4[116]$	$N^{-2}[116]$	4_{10}

Table 21: Operators in the vector representation V .

$\mathcal{O}$	$\mathcal{O} \varepsilon$	$\mathcal{O} N$	$\Delta_{4-\varepsilon}^{(1)}$	$\Delta(\varepsilon)$	$\Delta(N)$	Family
$\text{Op}[V, 0, 1]$	φ	φ	$[1 - \frac{1}{2}\varepsilon] + 0\varepsilon$	ε^8 [41]	N^{-3} [45]*	$2_0, 3_1, 11_{1,0}, 10_{0,1}$
$\text{Op}[V, 0, 2]$	φ_V^5	$\sigma^2\varphi$	$[5 - \frac{5}{2}\varepsilon] + \frac{2N+28}{N+8}\varepsilon$	ε^5 [5]	N^{-1} [66]	$2_2, 10_{2,1}$
$\text{Op}[V, 0, 3]$	$\square\varphi_V^5$	$\square\sigma^2\varphi$	$[7 - \frac{5}{2}\varepsilon] + \frac{2N+12}{N+8}\varepsilon$	ε^1		
$\text{Op}[V, 0, 4]$	φ_V^7	$\sigma^3\varphi$	$[7 - \frac{7}{2}\varepsilon] + \frac{3N+60}{N+8}\varepsilon$	ε^1	N^{-1} [66]	$2_3, 10_{3,1}$
$\text{Op}[V, 1, 1]$	$\partial\varphi_V^3$	$\partial\sigma\varphi$	$[4 - \frac{3}{2}\varepsilon] + \frac{N+2}{N+8}\varepsilon$	ε^5 [5]	N^{-1} [69]	$8_1, 12_1$
$\text{Op}[V, 1, 2]$	$\partial\varphi_V^5$	$\partial\sigma^2\varphi$	$[6 - \frac{5}{2}\varepsilon] + \frac{2N+18}{N+8}\varepsilon$	ε^5 [5]	N^{-1} [66]	
$\text{Op}[V, 2, 1]$	$\partial^2\varphi_V^3$		$[5 - \frac{3}{2}\varepsilon] + 0\varepsilon N^0$	ε^5 [5]		
$\text{Op}[V, 2, 2]$	$\partial^2\varphi_V^3$	$\partial^2\sigma\varphi$	$[5 - \frac{3}{2}\varepsilon] + \varepsilon N^0$	ε^5 [5]	N^{-1} [110]	8_2
$\text{Op}[V, 2, 3]$	$\partial^2\varphi_V^5$		$[7 - \frac{5}{2}\varepsilon] + \varepsilon N^0$	ε^1		
$\text{Op}[V, 2, 4]$	$\partial^2\varphi_V^5$		$[7 - \frac{5}{2}\varepsilon] + 2\varepsilon N^0$	ε^1		
$\text{Op}[V, 2, 5]$	$\partial^2\varphi_V^5$		$[7 - \frac{5}{2}\varepsilon] + 2\varepsilon N^0$	ε^1		
$\text{Op}[V, 3, 1]$	$\partial^3\varphi_V^3$		$[6 - \frac{3}{2}\varepsilon] + 0\varepsilon N^0$	ε^1		
$\text{Op}[V, 3, 2]$	$\partial^3\varphi_V^3$	$\partial^3\sigma\varphi$	$[6 - \frac{3}{2}\varepsilon] + \varepsilon N^0$	ε^1	N^{-1} [110]	8_3
$\text{Op}[V, 4, 1]$	$\partial^4\varphi_V^3$		$[7 - \frac{3}{2}\varepsilon] + 0\varepsilon$	ε^1		
$\text{Op}[V, 4, 2]$	$\partial^4\varphi_V^3$		$[7 - \frac{3}{2}\varepsilon] + 0\varepsilon N^0$	ε^1		
$\text{Op}[V, 4, 3]$	$\partial^4\varphi_V^3$	$\partial^4\sigma\varphi$	$[7 - \frac{3}{2}\varepsilon] + \varepsilon N^0$	ε^1	N^{-1} [110]	8_4
$\text{Op}[V, 5, 1]$	$\partial^5\varphi_V^3$		$[8 - \frac{3}{2}\varepsilon] + 0\varepsilon$	ε^1		
$\text{Op}[V, 5, 2]$	$\partial^5\varphi_V^3$		$[8 - \frac{3}{2}\varepsilon] + 0\varepsilon N^0$	ε^1		
$\text{Op}[V, 5, 3]$	$\partial^5\varphi_V^3$	$\partial^5\sigma\varphi$	$[8 - \frac{3}{2}\varepsilon] + \varepsilon N^0$	ε^1	N^{-1} [110]	8_5
$\text{Op}[V, 6, 1]$	$\partial^6\varphi_V^3$		$[9 - \frac{3}{2}\varepsilon] + 0\varepsilon$	ε^1		
$\text{Op}[V, 6, 2]$	$\partial^6\varphi_V^3$		$[9 - \frac{3}{2}\varepsilon] + 0\varepsilon$	ε^1		
$\text{Op}[V, 6, 3]$	$\partial^6\varphi_V^3$		$[9 - \frac{3}{2}\varepsilon] + 0\varepsilon N^0$	ε^1		
$\text{Op}[V, 6, 4]$	$\partial^6\varphi_V^3$	$\partial^6\sigma\varphi$	$[9 - \frac{3}{2}\varepsilon] + \varepsilon N^0$	ε^1	N^{-1} [110]	8_6
$\text{Op}[V, 7, 1]$	$\partial^7\varphi_V^3$		$[10 - \frac{3}{2}\varepsilon] + 0\varepsilon$	ε^1		
$\text{Op}[V, 7, 2]$	$\partial^7\varphi_V^3$		$[10 - \frac{3}{2}\varepsilon] + 0\varepsilon$	ε^1		
$\text{Op}[V, 7, 3]$	$\partial^7\varphi_V^3$		$[10 - \frac{3}{2}\varepsilon] + 0\varepsilon N^0$	ε^1		
$\text{Op}[V, 7, 4]$	$\partial^7\varphi_V^3$		$[10 - \frac{3}{2}\varepsilon] + \varepsilon N^0$	ε^1	N^{-1} [110]	8_7

* As noted in [71], the expression in [45] contains a misprint in equation (22).

Table 22: Operators in the T representation. Note the double eigenvalue at $\text{Op}[T,0,7]$ and $\text{Op}[T,0,8]$, *lifted at two-loop*.

$\mathcal{O}$	$\mathcal{O} \varepsilon$	$\mathcal{O} N$	$\Delta_{4-\varepsilon}^{(1)}$	$\Delta(\varepsilon)$	$\Delta(N)$	Family
$\text{Op}[T,0,1]$	φ_T^2	φ_T^2	$[2-\varepsilon] + \frac{2}{N+8}\varepsilon$	$\varepsilon^6[56]$	$N^{-2}[57]$	$3_2, 10_{0,2}, 11_{2,0}$
$\text{Op}[T,0,2]$	φ_T^4	$\sigma\varphi_T^2$	$[4-2\varepsilon] + \frac{N+16}{N+8}\varepsilon$	$\varepsilon^6[60]$	$N^{-1}[50]$	$10_{1,2}$
$\text{Op}[T,0,3]$	$\square\varphi_T^4$		$[6-2\varepsilon] + \frac{N+4}{N+8}\varepsilon$	$\varepsilon^5[5]$		
$\text{Op}[T,0,4]$	φ_T^6	$\sigma^2\varphi_T^2$	$[6-3\varepsilon] + \frac{2N+42}{N+8}\varepsilon$	$\varepsilon^5[5]$	$N^{-1}[66]$	$10_{2,2}$
$\text{Op}[T,0,5]$	$\square^2\varphi_T^4$		$[8-2\varepsilon] + 0\varepsilon N^0$	$\varepsilon^2[5]$		
$\text{Op}[T,0,6]$	$\square^2\varphi_T^4$		$[8-2\varepsilon] + \varepsilon N^0$	$\varepsilon^2[5]$		
$\text{Op}[T,0,7]$	$\square\varphi_T^6$		$[8-3\varepsilon] + \frac{2N+22}{N+8}\varepsilon$	$\varepsilon^2[5]$		
$\text{Op}[T,0,8]$	$\square\varphi_T^6$		$[8-3\varepsilon] + \frac{2N+22}{N+8}\varepsilon$	$\varepsilon^2[5]$		
$\text{Op}[T,0,9]$	φ_T^8	$\sigma^3\varphi_T^2$	$[8-4\varepsilon] + \frac{3N+80}{N+8}\varepsilon$	$\varepsilon^2[5]$	$N^{-1}[66]$	$10_{3,2}$
$\text{Op}[T,1,1]$	$\partial\square\varphi_T^4$		$[7-2\varepsilon] + \frac{3N+16}{3(N+8)}\varepsilon$	ε^1		
$\text{Op}[T,1,2]$	$\partial\varphi_T^6$		$[7-3\varepsilon] + \frac{2N+30}{N+8}\varepsilon$	ε^1		
$\text{Op}[T,2,1]$	$\mathcal{J}_{T,2}$	$\mathcal{J}_{T,2}$	$[4-\varepsilon] + 0\varepsilon$	$\varepsilon^4[65]\varepsilon^5[5]$	$N^{-2}[62]$	$5_2, 12_2$
$\text{Op}[T,2,2]$	$\partial^2\varphi_T^4$	$\partial^2\varphi_T^4$	$[6-2\varepsilon] + 0\varepsilon N^0$	$\varepsilon^5[5]$		
$\text{Op}[T,2,3]$	$\partial^2\varphi_T^4$		$[6-2\varepsilon] + \varepsilon N^0$	$\varepsilon^5[5]$		
$\text{Op}[T,2,4]$	$\partial^2\varphi_T^4$		$[6-2\varepsilon] + \varepsilon N^0$	$\varepsilon^5[5]$		
$\text{Op}[T,3,1]$	$\partial^3\varphi_T^4$		$[7-2\varepsilon] + 0\varepsilon N^0$	ε^1		
$\text{Op}[T,3,2]$	$\partial^3\varphi_T^4$		$[7-2\varepsilon] + \varepsilon N^0$	ε^1		
$\text{Op}[T,4,1]$	$\mathcal{J}_{T,4}$	$\mathcal{J}_{T,4}$	$[6-\varepsilon] + 0\varepsilon$	$\varepsilon^4[65]$	$N^{-2}[62]$	5_4
$\text{Op}[T,6,1]$	$\mathcal{J}_{T,6}$	$\mathcal{J}_{T,6}$	$[8-\varepsilon] + 0\varepsilon$	$\varepsilon^4[65]$	$N^{-2}[62]$	5_6

Table 23: Operators in the A representation.

$\mathcal{O}$	$\mathcal{O} \varepsilon$	$\mathcal{O} N$	$\Delta_{4-\varepsilon}^{(1)}$	$\Delta(\varepsilon)$	$\Delta(N)$	Family
$\text{Op}[A, 1, 1]$	J_{ij}^μ	J_{ij}^μ	$[3 - \varepsilon]$	exact	exact	6_1
$\text{Op}[A, 1, 2]$	$\partial\varphi_A^4$		$[5 - 2\varepsilon] + \frac{N+10}{N+8}\varepsilon$	ε^5 [5]		
$\text{Op}[A, 1, 3]$	$\partial\Box\varphi_A^4$		$[7 - 2\varepsilon] + 0\varepsilon N_{\dots}^0$	ε^1		
$\text{Op}[A, 1, 4]$	$\partial\Box\varphi_A^4$		$[7 - 2\varepsilon] + \varepsilon N_{\dots}^0$	ε^1		
$\text{Op}[A, 1, 5]$	$\partial\varphi_A^6$		$[7 - 3\varepsilon] + \frac{2N+32}{N+8}\varepsilon$	ε^1		
$\text{Op}[A, 3, 1]$	$\mathcal{J}_{A,3}$	$\mathcal{J}_{A,3}$	$[5 - \varepsilon]$	ε^4 [65]	N^{-2} [62]	6_3
$\text{Op}[A, 3, 2]$	$\partial^3\varphi_A^4$		$[7 - 2\varepsilon] + 0\varepsilon N_{\dots}^0$	ε^1		
$\text{Op}[A, 3, 3]$	$\partial^3\varphi_A^4$		$[7 - 2\varepsilon] + \varepsilon N_{\dots}^0$	ε^1		
$\text{Op}[A, 3, 4]$	$\partial^3\varphi_A^4$		$[7 - 2\varepsilon] + \varepsilon N_{\dots}^0$	ε^1		
$\text{Op}[A, 5, 1]$	$\mathcal{J}_{A,5}$	$\mathcal{J}_{A,5}$	$[7 - \varepsilon]$	ε^4 [65]	N^{-2} [62]	6_5

Table 24: Some operators in the rank 3 traceless-symmetric representation.

$\mathcal{O}$	$\mathcal{O} \varepsilon$	$\mathcal{O} N$	$\Delta_{4-\varepsilon}^{(1)}$	$\Delta(\varepsilon)$	$\Delta(N)$	Family
$\text{Op}[\text{Tm}[3], 0, 1]$	$\varphi_{T_3}^3$	$\varphi_{T_3}^3$	$[3 - \frac{3}{2}\varepsilon] + \frac{6}{N+8}\varepsilon$	$\varepsilon^6[60]$	$N^{-2}[62]$	$3_3, 10_{0,3}, 11_{3,0}$
$\text{Op}[\text{Tm}[3], 0, 2]$	$\varphi_{T_3}^5$	$\sigma\varphi_{T_3}^3$	$[5 - \frac{5}{2}\varepsilon] + \frac{N+26}{N+8}\varepsilon$	$\varepsilon^5[5]$	$N^{-1}[66]$	$10_{1,3}$
$\text{Op}[\text{Tm}[3], 0, 3]$	$\square\varphi_{T_3}^5$	$\sigma\square\varphi_{T_3}^5$	$[7 - \frac{5}{2}\varepsilon] + \frac{N+10}{N+8}\varepsilon$	ε^1		
$\text{Op}[\text{Tm}[3], 0, 4]$	$\varphi_{T_3}^7$	$\sigma^2\varphi_{T_3}^3$	$[7 - \frac{7}{2}\varepsilon] + \frac{2N+58}{N+8}\varepsilon$	ε^1	$N^{-1}[66]$	$10_{2,3}$
$\text{Op}[\text{Tm}[3], 1, 1]$	$\partial\varphi_{T_3}^5$	$\partial\sigma\varphi_{T_3}^3$	$[6 - \frac{5}{2}\varepsilon] + \frac{N+16}{N+8}\varepsilon$	$\varepsilon^5[5]$		
$\text{Op}[\text{Tm}[3], 2, 1]$	$\partial^2\varphi_{T_3}^3$	$\partial^2\varphi_{T_3}^3$	$[5 - \frac{3}{2}\varepsilon] + \frac{10}{3(N+8)}\varepsilon$	$\varepsilon^2[100]\varepsilon^5[5]$	$N^{-1}[1, 118]$	12_3
$\text{Op}[\text{Tm}[3], 2, 2]$	$\partial^2\varphi_{T_3}^5$	$\partial^2\varphi_{T_5}^3$	$[7 - \frac{5}{2}\varepsilon] + 0\varepsilon N^0_{\dots}$	ε^1		
$\text{Op}[\text{Tm}[3], 2, 3]$	$\partial^2\varphi_{T_3}^5$	$\partial^2\sigma\varphi_{T_3}^3$	$[7 - \frac{5}{2}\varepsilon] + \varepsilon N^0_{\dots}$	ε^1		
$\text{Op}[\text{Tm}[3], 2, 4]$	$\partial^2\varphi_{T_3}^5$	$\partial^2\sigma\varphi_{T_3}^3$	$[7 - \frac{5}{2}\varepsilon] + \varepsilon N^0_{\dots}$	ε^1		
$\text{Op}[\text{Tm}[3], 3, 1]$	$\partial^3\varphi_{T_3}^3$	$\partial^3\varphi_{T_3}^3$	$[6 - \frac{3}{2}\varepsilon] + \frac{1}{N+8}\varepsilon$	$\varepsilon^2[100]$	$N^{-1}[1, 118]$	
$\text{Op}[\text{Tm}[3], 4, 1]$	$\partial^4\varphi_{T_3}^3$	$\partial^4\varphi_{T_3}^3$	$[7 - \frac{3}{2}\varepsilon] + \frac{14}{5(N+8)}\varepsilon$	$\varepsilon^2[100]$	$N^{-1}[1, 118]$	
$\text{Op}[\text{Tm}[3], 5, 1]$	$\partial^5\varphi_{T_3}^3$	$\partial^5\varphi_{T_3}^3$	$[8 - \frac{3}{2}\varepsilon] + \frac{4}{3(N+8)}\varepsilon$	$\varepsilon^2[100]$	$N^{-1}[1, 118]$	
$\text{Op}[\text{Tm}[3], 6, 1]$	$\partial^6\varphi_{T_3}^3$	$\partial^6\varphi_{T_3}^3$	$[9 - \frac{3}{2}\varepsilon] + 0\varepsilon$	$\varepsilon^2[100]$	$N^{-1}[1, 118]$	
$\text{Op}[\text{Tm}[3], 6, 2]$	$\partial^6\varphi_{T_3}^3$	$\partial^6\varphi_{T_3}^3$	$[9 - \frac{3}{2}\varepsilon] + \frac{18}{7(N+8)}\varepsilon$	$\varepsilon^2[100]$	$N^{-1}[1, 118]$	
$\text{Op}[\text{Tm}[3], 7, 1]$	$\partial^7\varphi_{T_3}^3$	$\partial^7\varphi_{T_3}^3$	$[10 - \frac{3}{2}\varepsilon] + \frac{3}{2(N+8)}\varepsilon$	$\varepsilon^2[1, 100]$	$N^{-1}[1, 118]$	

Table 25: Some operators in the rank 4 traceless-symmetric representation. Writing “[1, 100]” means that this particular result was not listed in [100], but was computed using the method of that paper, and likewise for “[1, 118]”.

$\mathcal{O}$	$\mathcal{O} \varepsilon$	$\mathcal{O} N$	$\Delta_{4-\varepsilon}^{(1)}$	$\Delta(\varepsilon)$	$\Delta(N)$	Family
$\text{Op}[\text{Tm}[4], 0, 1]$	$\varphi_{T_4}^4$	$\varphi_{T_4}^4$	$[4 - 2\varepsilon] + \frac{12}{N+8}\varepsilon$	$\varepsilon^6[60]$	$N^{-2}[62]$	$3_4, 10_{0,4}, 11_{4,0}$
$\text{Op}[\text{Tm}[4], 0, 2]$	$\varphi_{T_4}^6$	$\sigma\varphi_{T_4}^4$	$[6 - 3\varepsilon] + \frac{N+38}{N+8}\varepsilon$	$\varepsilon^5[5]$	$N^{-1}[66]$	$10_{1,4}$
$\text{Op}[\text{Tm}[4], 0, 3]$	$\square^2\varphi_{T_4}^4$		$[8 - 2\varepsilon] + \frac{20}{3(N+8)}\varepsilon$	$\varepsilon^2[5]$		
$\text{Op}[\text{Tm}[4], 0, 4]$	$\square\varphi_{T_4}^6$		$[8 - 3\varepsilon] + \frac{N+18}{N+8}\varepsilon$	$\varepsilon^2[5]$		
$\text{Op}[\text{Tm}[4], 0, 5]$	$\varphi_{T_4}^8$	$\sigma^2\varphi_{T_4}^4$	$[8 - 4\varepsilon] + \frac{2N+76}{N+8}\varepsilon$	$\varepsilon^2[5]$	$N^{-1}[66]$	$10_{2,4}$
$\text{Op}[\text{Tm}[4], 1, 1]$	$\partial\varphi_{T_4}^6$		$[7 - 3\varepsilon] + \frac{N+26}{N+8}\varepsilon$	ε^1		
$\text{Op}[\text{Tm}[4], 2, 1]$	$\partial^2\varphi_{T_4}^4$	$\partial^2\varphi_{T_4}^4$	$[6 - 2\varepsilon] + \frac{26}{3(N+8)}\varepsilon$	$\varepsilon^2[100]\varepsilon^5[5]$	$N^{-1}[1, 118]$	12_4
$\text{Op}[\text{Tm}[4], 3, 1]$	$\partial^3\varphi_{T_4}^4$	$\partial^3\varphi_{T_4}^4$	$[7 - 2\varepsilon] + \frac{6}{N+8}\varepsilon$	$\varepsilon^2[100]$	$N^{-1}[1, 118]$	
$\text{Op}[\text{Tm}[4], 4, 1]$	$\partial^4\varphi_{T_4}^4$	$\partial^4\varphi_{T_4}^4$	$[8 - 2\varepsilon] + \frac{8}{3(N+8)}\varepsilon$	$\varepsilon^2[1, 100]$	$N^{-1}[1, 118]$	
$\text{Op}[\text{Tm}[4], 4, 2]$	$\partial^4\varphi_{T_4}^4$	$\partial^4\varphi_{T_4}^4$	$[8 - 2\varepsilon] + \frac{38}{5(N+8)}\varepsilon$	$\varepsilon^2[1, 100]$	$N^{-1}[1, 118]$	
$\text{Op}[\text{Tm}[4], 5, 1]$	$\partial^5\varphi_{T_4}^4$	$\partial^5\varphi_{T_4}^5$	$[9 - 2\varepsilon] + \frac{6}{N+8}\varepsilon$	$\varepsilon^2[1, 100]$	$N^{-1}[1, 118]$	
$\text{Op}[\text{Tm}[4], 6, 1]$	$\partial^6\varphi_{T_4}^4$	$\partial^6\varphi_{T_4}^4$	$[10 - 2\varepsilon] + 0\varepsilon N_{\dots}^0$	$\varepsilon^2[1, 100]$	$N^{-1}[1, 118]$	
$\text{Op}[\text{Tm}[4], 6, 2]$	$\partial^6\varphi_{T_4}^4$	$\partial^6\varphi_{T_4}^4$	$[10 - 2\varepsilon] + 0\varepsilon N_{\dots}^0$	$\varepsilon^2[1, 100]$	$N^{-1}[1, 118]$	
$\text{Op}[\text{Tm}[4], 6, 3]$	$\partial^6\varphi_{T_4}^4$	$\partial^6\varphi_{T_4}^4$	$[10 - 2\varepsilon] + 0\varepsilon N_{\dots}^0$	$\varepsilon^2[1, 100]$	$N^{-1}[1, 118]$	

For the leading scalar T_3 operator $\text{Op}[\text{Tm}[3], 0, 1]$, we report $\varepsilon^3[120]\varepsilon^4[112]\varepsilon^6[60]$ and $N^{-1}[114]N^{-2}[62]$. Numerical values in $d = 3$ are $\Delta_{\varphi_{T_3}^3} = 2.1086(3)$ for $N = 2$ [76], $\Delta_{\varphi_{T_3}^3} = 1.0384(10)$ for $N = 3$ [89], and $\Delta_{\varphi_{T_3}^3} = 1.9768(10)$ for $N = 4$ [89].

Table 26: Scalar operators in various $O(N)$ representations.

$\mathcal{O}$	$\mathcal{O} _{\varepsilon}$	$\mathcal{O} _N$	$\Delta_{4-\varepsilon}^{(1)}$	$\Delta(\varepsilon)$	$\Delta(N)$	Family
$\text{Op}[\text{Hm}[3], 0, 1]$	$\square\varphi_{H_3}^5$		$[7 - \frac{5}{2}\varepsilon] + \frac{N+13}{N+8}\varepsilon$	ε^1		
$\text{Op}[\text{B4}, 0, 1]$	$\square\varphi_{B_4}^4$	$\square\varphi_{B_4}^4$	$[6 - 2\varepsilon] + \frac{6}{N+8}\varepsilon$	$\varepsilon^5[5]$		
$\text{Op}[\text{B4}, 0, 2]$	$\square^2\varphi_{B_4}^4$		$[8 - 2\varepsilon] + \frac{8}{3(N+8)}\varepsilon$	$\varepsilon^2[5]$		
$\text{Op}[\text{B4}, 0, 3]$	$\square\varphi_{B_4}^6$		$[8 - 3\varepsilon] + \frac{N+24}{N+8}\varepsilon$	$\varepsilon^2[5]$		
$\text{Op}[\text{Hm}[4], 0, 1]$	$\square\varphi_{H_4}^6$		$[8 - 3\varepsilon] + \frac{N+22}{N+8}\varepsilon$	$\varepsilon^2[5]$		
$\text{Op}[\text{Tm}[5], 0, 1]$	$\varphi_{T_5}^5$	$\varphi_{T_5}^5$	$[5 - \frac{5}{2}\varepsilon] + \frac{20}{N+8}\varepsilon$	$\varepsilon^5[1, 60, 112]\varepsilon^6[113]$	$N^{-2}[62]$	$3_5, 10_{0,5}, 11_{5,0}$
$\text{Op}[\text{Tm}[5], 0, 2]$	$\varphi_{T_5}^7$	$\sigma\varphi_{T_7}^5$	$[7 - \frac{7}{2}\varepsilon] + \frac{N+52}{N+8}\varepsilon$	ε^1	$N^{-1}[66]$	$10_{1,6}$
$\text{Op}[\text{Tm}[6], 0, 1]$	$\varphi_{T_6}^6$	$\varphi_{T_6}^6$	$[6 - 3\varepsilon] + \frac{30}{N+8}\varepsilon$	$\varepsilon^5[1, 60, 112]\varepsilon^6[113]$	$N^{-2}[62]$	$3_6, 10_{0,6}, 11_{6,0}$
$\text{Op}[\text{Tm}[6], 0, 2]$	$\varphi_{T_6}^8$	$\sigma\varphi_{T_6}^6$	$[8 - 3\varepsilon] + \frac{N+68}{N+8}\varepsilon$	$\varepsilon^2[5]$	$N^{-1}[66]$	$10_{1,6}$
$\text{Op}[\text{YT}[\{3, 2\}], 0, 1]$	$\square\varphi_{Y_{3,2}}^5$		$[7 - \frac{3}{2}\varepsilon] + \frac{12}{N+8}\varepsilon$	ε^1		
$\text{Op}[\text{YT}[\{4, 2\}], 0, 1]$	$\square\varphi_{Y_{4,2}}^6$		$[8 - 3\varepsilon] + \frac{20}{N+8}\varepsilon$	$\varepsilon^2[5]$		

Table 27: Spinning operators in various $O(N)$ representations.

$\mathcal{O}$	$\mathcal{O} \varepsilon$	$\mathcal{O} N$	$\Delta_{4-\varepsilon}^{(1)}$	$\Delta(\varepsilon)$	$\Delta(N)$	Family
$\text{Op}[\text{Hm}[3], 1, 1]$	$\partial\varphi_{H_3}^3$	$\partial\varphi_{H_3}^3$	$[4 - \frac{3}{2}\varepsilon] + \frac{3}{N+8}\varepsilon$	$\varepsilon^5[5]$	$N^{-1}[115]$	$11_{2,1}$
$\text{Op}[\text{Hm}[3], 1, 2]$	$\partial\varphi_{H_3}^5$	$\partial\sigma\varphi_{H_3}^3$	$[6 - \frac{5}{2}\varepsilon] + \frac{N+19}{N+8}\varepsilon$	$\varepsilon^5[5]$		
$\text{Op}[\text{Hm}[3], 2, 1]$	$\partial^2\varphi_{H_3}^3$	$\partial^2\varphi_{H_3}^3$	$[5 - \frac{3}{2}\varepsilon] + \frac{4}{3(N+8)}\varepsilon$	$\varepsilon^5[5]$		
$\text{Op}[\text{Hm}[3], 2, 2]$	$\partial^2\varphi_{H_3}^5$		$[7 - \frac{5}{2}\varepsilon] + \frac{3N+35}{3(N+8)}\varepsilon$	ε^1		
$\text{Op}[\text{Hm}[3], 2, 3]$	$\partial^2\varphi_{H_3}^5$		$[7 - \frac{5}{2}\varepsilon] + \frac{3N+50}{3(N+8)}\varepsilon$	ε^1		
$\text{Op}[\text{Hm}[3], 3, 1]$	$\partial^3\varphi_{H_3}^3$	$\partial^3\varphi_{H_3}^3$	$[6 - \frac{3}{2}\varepsilon] + \frac{5}{2(N+8)}\varepsilon$	ε^1		
$\text{Op}[\text{Hm}[3], 4, 1]$	$\partial^4\varphi_{H_3}^3$	$\partial^4\varphi_{H_3}^3$	$[7 - \frac{3}{2}\varepsilon] + 0\varepsilon$	ε^1		
$\text{Op}[\text{Hm}[3], 4, 2]$	$\partial^4\varphi_{H_3}^3$	$\partial^4\varphi_{H_3}^3$	$[7 - \frac{3}{2}\varepsilon] + \frac{8}{5(N+8)}\varepsilon$	ε^1		
$\text{Op}[\text{Hm}[3], 5, 1]$	$\partial^5\varphi_{H_3}^3$	$\partial^5\varphi_{H_3}^3$	$[8 - \frac{3}{2}\varepsilon] + 0\varepsilon$	ε^1		
$\text{Op}[\text{Hm}[3], 5, 2]$	$\partial^5\varphi_{H_3}^3$	$\partial^5\varphi_{H_3}^3$	$[8 - \frac{3}{2}\varepsilon] + \frac{7}{3(N+8)}\varepsilon$	ε^1		
$\text{Op}[\text{Hm}[3], 6, 1]$	$\partial^6\varphi_{H_3}^3$	$\partial^6\varphi_{H_3}^3$	$[9 - \frac{3}{2}\varepsilon] + 0\varepsilon$	ε^1		
$\text{Op}[\text{Hm}[3], 6, 2]$	$\partial^6\varphi_{H_3}^3$	$\partial^6\varphi_{H_3}^3$	$[9 - \frac{3}{2}\varepsilon] + \frac{12}{7(N+8)}\varepsilon$	ε^1		
$\text{Op}[\text{A3}, 3, 1]$	$\partial^3\varphi_{A_3}^3$	$\partial^3\varphi_{A_3}^3$	$[6 - \frac{3}{2}\varepsilon] + 0\varepsilon$	ε^1		
$\text{Op}[\text{A3}, 5, 1]$	$\partial^5\varphi_{A_3}^3$	$\partial^5\varphi_{A_3}^3$	$[8 - \frac{3}{2}\varepsilon] + 0\varepsilon$	ε^1		
$\text{Op}[\text{A3}, 6, 1]$	$\partial^6\varphi_{A_3}^3$	$\partial^6\varphi_{A_3}^3$	$[9 - \frac{3}{2}\varepsilon] + 0\varepsilon$	ε^1		
$\text{Op}[\text{Hm}[4], 1, 1]$	$\partial\varphi_{H_4}^4$	$\partial\varphi_{H_4}^4$	$[5 - 2\varepsilon] + \frac{8}{N+8}\varepsilon$	$\varepsilon^5[5]$	$N^{-1}[115]$	$11_{3,1}$
$\text{Op}[\text{Hm}[4], 1, 2]$	$\partial\Box\varphi_{H_4}^4$		$[7 - 2\varepsilon] + \frac{16}{3(N+8)}\varepsilon$	ε^1		
$\text{Op}[\text{Hm}[4], 1, 3]$	$\partial\varphi_{H_4}^6$		$[7 - 3\varepsilon] + \frac{N+30}{N+8}\varepsilon$	ε^1		
$\text{Op}[\text{Hm}[4], 2, 1]$	$\partial^2\varphi_{H_4}^4$	$\partial^2\varphi_{H_4}^4$	$[6 - 2\varepsilon] + \frac{6}{N+8}\varepsilon$	$\varepsilon^5[5]$		
$\text{Op}[\text{Hm}[4], 3, 1]$	$\partial^3\varphi_{H_4}^4$	$\partial^3\varphi_{H_4}^4$	$[7 - 2\varepsilon] + \frac{10}{3(N+8)}\varepsilon$	ε^1		
$\text{Op}[\text{Hm}[4], 3, 2]$	$\partial^3\varphi_{H_4}^4$	$\partial^3\varphi_{H_4}^4$	$[7 - 2\varepsilon] + \frac{7}{N+8}\varepsilon$	ε^1		
$\text{Op}[\text{B4}, 2, 1]$	$\partial^2\varphi_{B_4}^4$	$\partial^2\varphi_{B_4}^4$	$[6 - 2\varepsilon] + \frac{14}{3(N+8)}\varepsilon$	$\varepsilon^5[5]$	$N^{-1}[115]$	$11_{2,2}$
$\text{Op}[\text{Tm}[5], 2, 1]$	$\partial^2\varphi_{T_5}^5$	$\partial^2\varphi_{T_5}^5$	$[7 - \frac{5}{2}\varepsilon] + \frac{16}{N+8}\varepsilon$	$\varepsilon^2[2, 100]$	$N^{-1}[2, 118]$	12_5
$\text{Op}[\text{Hm}[5], 1, 1]$	$\partial\varphi_{H_5}^5$	$\partial\varphi_{H_5}^5$	$[6 - \frac{5}{2}\varepsilon] + \frac{15}{N+8}\varepsilon$	$\varepsilon^5[5]$	$N^{-1}[115]$	$11_{4,1}$
$\text{Op}[\text{Hm}[5], 2, 1]$	$\partial^2\varphi_{H_5}^5$	$\partial^2\varphi_{H_5}^5$	$[7 - \frac{5}{2}\varepsilon] + \frac{38}{3(N+8)}\varepsilon$	ε^1		
$\text{Op}[\text{Hm}[6], 1, 1]$	$\partial\varphi_{H_6}^6$	$\partial\varphi_{H_6}^6$	$[7 - 3\varepsilon] + \frac{24}{N+8}\varepsilon$	ε^1	$N^{-1}[115]$	$11_{5,1}$
$\text{Op}[\text{Tm}[6], 2, 1]$	$\partial^2\varphi_{T_6}^6$	$\partial^2\varphi_{T_6}^6$	$[8 - 3\varepsilon] + \frac{76}{3(N+8)}\varepsilon$	$\varepsilon^2[2, 100]$	$N^{-1}[2, 118]$	12_6
$\text{Op}[\text{YT}[\{3, 2\}], 2, 1]$	$\partial^2\varphi_{Y_{3,2}}^5$	$\partial^2\varphi_{Y_{3,2}}^5$	$[7 - \frac{5}{2}\varepsilon] + \frac{32}{3(N+8)}\varepsilon$	ε^1		
$\text{Op}[\text{YT}[\{2, 1, 1\}], 1, 1]$	$\partial\Box\varphi_{Y_{2,1,1}}^4$		$[7 - 2\varepsilon] + \frac{8}{3(N+8)}\varepsilon$	ε^1		
$\text{Op}[\text{YT}[\{2, 1, 1\}], 3, 1]$	$\partial^3\varphi_{Y_{2,1,1}}^4$		$[7 - 2\varepsilon] + \frac{11}{3(N+8)}\varepsilon$	ε^1		

3.5 Operators in non-traceless-symmetric Lorentz representations

Here we will just mention a few operators in non-traceless-symmetric Lorentz representations, in order to give a complete presentation of the spectrum up to $\Delta^{4d} \leq 6$.

Table 28: $O(N)$ singlet and Lorentz non-traceless-symmetric operators up to 4d dimension 8. The 4d irrep notation is the one used in [95].

$\mathcal{O}$	Lorentz YT	4d irrep	Field content	$\Delta_{4-\varepsilon}^{(1)}$
$\text{Op}[S, \{2, 1\}, 1]$	$\begin{array}{ c c } \hline \square & \square \\ \hline \end{array}$	$(\frac{3}{2}, \frac{1}{2}) + (\frac{1}{2}, \frac{3}{2})$	$\partial^3 \varphi_S^4$	$[7 - 2\varepsilon] + \frac{N+2}{N+8}\varepsilon$
$\text{Op}[S, \{3, 1\}, 1]$	$\begin{array}{ c c c } \hline \square & \square & \square \\ \hline \end{array}$	$(2, 1) + (1, 2)$	$\partial^4 \varphi_S^4$	$[8 - 2\varepsilon] + \frac{N+2}{N+8}\varepsilon$
$\text{Op}[S, \{2, 2\}, 1]$	$\begin{array}{ c c } \hline \square & \square \\ \hline \end{array}$	$(2, 0) + (0, 2)$	$\partial^4 \varphi_S^4$	$[8 - 2\varepsilon] + \frac{3N+20-\sqrt{9N^2+64N+288}}{6(N+8)}\varepsilon$
$\text{Op}[S, \{2, 2\}, 2]$	$\begin{array}{ c c } \hline \square & \square \\ \hline \end{array}$	$(2, 0) + (0, 2)$	$\partial^4 \varphi_S^4$	$[8 - 2\varepsilon] + \frac{3N+20+\sqrt{9N^2+64N+288}}{6(N+8)}\varepsilon$

Dimension $\Delta = 5$ One operator in $O(N)$ irrep A_3 , Lorentz irrep $\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}$: $\varphi^{[i} \partial_{[\mu} \varphi^j \partial_{\nu]} \varphi^{k]}$, dimension $\Delta = [5 - \frac{3}{2}\varepsilon] + \dots + O(\varepsilon^6)$, where we have ε^5 [5]. Implemented as $\text{Op}[A3, \{1, 1\}, 1]$. Note that this anomalous dimensions is negative at $O(\varepsilon^2)$ and $O(\varepsilon^3)$.

Dimension $\Delta = 6$ Two operators constructed out of three fields and three gradients: $O(N)$ irreps V and H_3 , both in Lorentz irrep $\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}$, and both with dimension $\Delta = [6 - \frac{3}{2}\varepsilon] + O(\varepsilon^2)$. Moreover, two operators constructed out of four fields and two gradients, both in Lorentz irrep $\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}$: $O(N)$ irrep A , dimension $\Delta = [6 - 2\varepsilon] + \frac{N+2}{N+8}\varepsilon + \dots + O(\varepsilon^6)$; $O(N)$ irrep $Y_{2,1,1}$, dimension $\Delta = [6 - 2\varepsilon] + \frac{4}{N+8}\varepsilon + \dots + O(\varepsilon^6)$. For the latter two we have ε^5 [5]. These operators are implemented as

$$\text{Op}[V, \{2, 1\}, 1], \text{Op}[Hm[3], \{2, 1\}, 1], \text{Op}[A, \{1, 1\}, 1], \text{Op}[YT[\{2, 1, 1\}], \{1, 1\}, 1]. \quad (3.39)$$

In table 28 we give the leading $O(N)$ singlet operators in non-traceless-symmetric $O(N)$ representations, up to $\Delta^{4d} \leq 8$.

Operators in the Lorentz representation labelled by the Young tableau $\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}$ are related to parity-odd scalars in the three-dimensional theory. In the free three-dimensional $O(N)$ symmetric CFT, one can for generic $N > 1$ construct a parity odd singlet scalar out of four fields and five gradients, giving (in the free theory) dimension 7 [121], equation 3.1. It has the form $\epsilon^{\mu\nu\rho} \partial_\mu \partial_\nu \partial_\rho \square \varphi_S^4$. The corresponding operator in the $4 - \varepsilon$ expansion has dimension $[9 - 2\varepsilon] + 0\varepsilon + O(\varepsilon^2)$.

3.6 OPE coefficients in the $\varphi \times \varphi$ OPE

Table 29: OPE coefficients in the $\varphi \times \varphi$ OPE for operators for general N .

$\mathcal{O}$	$\mathcal{O} \varepsilon$	$\mathcal{O} N$	$\lambda_{\varphi\varphi\mathcal{O}}^2(\varepsilon)$	$\lambda_{\varphi\varphi\mathcal{O}}^2(\varepsilon)$	$\lambda_{\varphi\varphi\mathcal{O}}^2(N)$	$\lambda_{\varphi\varphi\mathcal{O}}^2(N)$
$\varphi, \varphi, \text{Op}[\text{S}, 0, 0]$	$\mathbb{1}$	$\mathbb{1}$	1	exact	1	exact
$\varphi, \varphi, \text{Op}[\text{S}, 0, 1]$	φ_S^2	σ	$O(1)$	ε^3 [64]	$O(N^{-1})$	N^{-2} [66]
$\varphi, \varphi, \text{Op}[\text{S}, 0, 2]$	φ_S^4	σ^2	$O(\varepsilon^2)$	ε^2 [65]	$O(N^{-2})$	N^{-2} [122]
$\varphi, \varphi, \text{Op}[\text{S}, 0, 4]$	φ_S^6	σ^3	$O(\varepsilon^4)$	ε^4 [123]		
$\varphi, \varphi, \text{Op}[\text{S}, 2, 1]$	$T^{\mu\nu}$	$T^{\mu\nu}$	$O(1)$	ε^4 [65]	$O(N^{-1})$	N^{-2} [63]
$\varphi, \varphi, \text{Op}[\text{S}, 2, 2]$	$\partial^2 \varphi_S^4$		$O(\varepsilon^2)$	ε^2 [1, 65]		
$\varphi, \varphi, \text{Op}[\text{S}, 2, 3]$	$\partial^2 \varphi_S^4$	$[\sigma, \sigma]_{0,2}$	$O(\varepsilon^2)$	ε^2 [1, 65]	$O(N^{-2})$	N^{-2} [122]
$\varphi, \varphi, \text{Op}[\text{S}, 4, 1]$	$\mathcal{J}_{S,4}$	$\mathcal{J}_{S,4}$	$O(1)$	ε^4 [65]	$O(N^{-1})$	N^{-2} [122]
$\varphi, \varphi, \text{Op}[\text{T}, 0, 1]$	φ_T^2	φ_T^2	$O(1)$	ε^3 [64]	$O(1)$	N^{-1} [64, 124]
$\varphi, \varphi, \text{Op}[\text{T}, 0, 2]$	φ_T^4		$O(\varepsilon^2)$	ε^2 [65]	$O(N^{-1})$	N^{-1} [1, 124]
$\varphi, \varphi, \text{Op}[\text{T}, 2, 1]$	$\mathcal{J}_{T,2}$	$\mathcal{J}_{T,2}$	$O(1)$	ε^4 [65]	$O(1)$	N^{-1} [64, 124]
$\varphi, \varphi, \text{Op}[\text{T}, 4, 1]$	$\mathcal{J}_{T,4}$	$\mathcal{J}_{T,4}$	$O(1)$	ε^4 [65]	$O(1)$	N^{-1} [64, 124]
$\varphi, \varphi, \text{Op}[\text{A}, 1, 1]$	J^μ	J^μ	$O(1)$	ε^4 [65]	$O(1)$	N^{-1} [69] $N^{-2}(\text{num})$ [122]
$\varphi, \varphi, \text{Op}[\text{A}, 1, 2]$	$\partial \varphi_A^4$		$O(\varepsilon^2)$	ε^2 [65]	$O(N^{-1})$	N^{-1} [1, 124]
$\varphi, \varphi, \text{Op}[\text{A}, 3, 1]$	$\mathcal{J}_{A,3}$	$\mathcal{J}_{A,3}$	$O(1)$	ε^4 [65]	$O(1)$	N^{-1} [64, 124]

Table 30: Other OPE coefficients implemented for general N .

$\mathcal{O}_1, \mathcal{O}_2, \mathcal{O}_3$	$\mathcal{O}_1, \mathcal{O}_2, \mathcal{O}_3 _\varepsilon$	$\mathcal{O}_1, \mathcal{O}_2, \mathcal{O}_3 _N$	$\lambda_{\mathcal{O}_1 \mathcal{O}_2 \mathcal{O}_3}^2 _\varepsilon$	$\lambda_{\mathcal{O}_1 \mathcal{O}_2 \mathcal{O}_3}^2 _N$
$\text{Op}[S, 0, 1], \text{Op}[S, 0, 1], \text{Op}[S, 0, 1]$	$\varphi_S^2, \varphi_S^2, \varphi_S^2$	σ, σ, σ	$\varepsilon^1[107]$	$N^{-2}[125]$
$\text{Op}[S, 0, 1], \text{Op}[S, 0, 1], \text{Op}[S, 0, 2]$	$\varphi_S^2, \varphi_S^2, \varphi_S^4$	σ, σ, σ^2	$\varepsilon^1[7]$	$N^{-1}(3d)[126]$
$\text{Op}[S, 0, 1], \text{Op}[S, 0, 1], \text{Op}[S, 0, 3]$	$\varphi_S^2, \varphi_S^2, \square \varphi_S^4$	$\sigma, \sigma, \square \sigma^2$	$\varepsilon^1[7]$	
$\text{Op}[S, 0, 1], \text{Op}[S, 0, 1], \text{Op}[S, 0, 4]$	$\varphi_S^2, \varphi_S^2, \varphi_S^6$	σ, σ, σ^3		$N^{-1}[127]$
$\text{Op}[T, 0, 1], \text{Op}[T, 0, 1], \text{Op}[S, 0, 1]$	$\varphi_T^2, \varphi_T^2, \varphi_S^2$	$\varphi_T^2, \varphi_T^2, \sigma$	$\varepsilon^1[7]$	
$\text{Op}[T, 0, 1], \text{Op}[T, 0, 1], \text{Op}[S, 0, 2]$	$\varphi_T^2, \varphi_T^2, \varphi_S^4$	$\varphi_T^2, \varphi_T^2, \sigma^2$	$\varepsilon^1[7]$	
$\text{Op}[T, 0, 1], \text{Op}[T, 0, 1], \text{Op}[T, 0, 1]$	$\varphi_T^2, \varphi_T^2, \varphi_T^2$	$\varphi_T^2, \varphi_T^2, \varphi_T^2$	$\varepsilon^1[7]$	
$\text{Op}[T, 0, 1], \text{Op}[T, 0, 1], \text{Op}[T, 0, 2]$	$\varphi_T^2, \varphi_T^2, \varphi_T^4$	$\varphi_T^2, \varphi_T^2, \sigma \varphi_T^2$	$\varepsilon^1[7]$	
$\text{Op}[T, 0, 1], \text{Op}[T, 0, 1], \text{Op}[\text{Tm}[4], 0, 1]$	$\varphi_T^2, \varphi_T^2, \varphi_{T_4}^4$	$\varphi_T^2, \varphi_T^2, \varphi_{T_4}^4$	$\varepsilon^1[7]$	
$\text{Op}[T, 0, 1], \text{Op}[T, 0, 1], \text{Op}[B_4, 0, 1]$	$\varphi_T^2, \varphi_T^2, \square \varphi_{B_4}^4$	$\varphi_T^2, \varphi_T^2, \square \varphi_{B_4}^4$	$\varepsilon^1[7]$	

A Some extension on representation theory

We would like to give some more details on some additional $O(N)$ representations. We reproduce table 7 with some additional irreps displayed

Table 7: Dimensions of the irreducible representations of $O(N)$ symmetry with rank up to 4. For $N = 2$ and $N = 3$ we also give additional symbols for the irreps, corresponding to those used in [128] and [77]. For $N = 3$, the parity corresponds to the parity of the number of boxes in the Young tableau.

Irrep			$N = 2$	$N = 3$	$N = 4$	$N \geq 5$
S	$\bullet$	S	$q^p = 0^+ \quad 1$	$\mathbf{q}^p = 0^+ \quad 1$	1	1
V	$\square$	V	$q = 1 \quad 2$	$\mathbf{q}^p = 1^- \quad 3$	4	N
A	$\begin{array}{ c } \hline \square \\ \hline \end{array}$	A	$q^p = 0^- \quad 1$	$\mathbf{q}^p = 1^+ \quad 3$	$3 + \bar{3}$	$N(N-1)/2$
T	$\begin{array}{ c c } \hline \square & \square \\ \hline \end{array}$	T	$q = 2 \quad 2$	$\mathbf{q}^p = 2^+ \quad 5$	9	$(N-1)(N+2)/2$
A_3	$\begin{array}{ c } \hline \square \\ \square \\ \hline \end{array}$	A_3		$\mathbf{q}^p = 0^- \quad 1$	4^-	$N(N-1)(N-2)/6$
T_3	$\begin{array}{ c c c } \hline \square & \square & \square \\ \hline \end{array}$	$T_m[3]$	$q = 3 \quad 2$	$\mathbf{q}^p = 3^- \quad 7$	16	$N(N+4)(N-1)/6$
H_3	$\begin{array}{ c c } \hline \square & \square \\ \hline \end{array}$	$H_m[3]$		$\mathbf{q}^p = 2^- \quad 5$	$8 + \bar{8}$	$N(N+2)(N-2)/3$
A_4	$\begin{array}{ c } \hline \square \\ \square \\ \square \\ \hline \end{array}$	—			1^-	$N(N-1)(N-2)(N-3)/24$
T_4	$\begin{array}{ c c c c } \hline \square & \square & \square & \square \\ \hline \end{array}$	$T_m[4]$	$q = 4 \quad 2$	$\mathbf{q}^p = 4^+ \quad 9$	25	$N(N+1)(N+6)(N-1)/24$
H_4	$\begin{array}{ c c c } \hline \square & \square & \square \\ \hline \end{array}$	$H_m[4]$		$\mathbf{q}^p = 3^+ \quad 7$	$15 + \bar{15}$	$(N+1)(N+4)(N-1)(N-2)/8$
B_4	$\begin{array}{ c c } \hline \square & \square \\ \hline \end{array}$	B_4			$5 + \bar{5}$	$N(N+1)(N+2)(N-3)/12$
$Y_{2,1,1}$	$\begin{array}{ c c } \hline \square & \square \\ \hline \end{array}$	$YT[\{2,1,1\}]$			9^-	$N(N+2)(N-1)(N-3)/8$
T_5	$\begin{array}{ c c c c c } \hline \square & \square & \square & \square & \square \\ \hline \end{array}$	$T_m[5]$	$q = 5 \quad 2$	$\mathbf{q}^p = 5^- \quad 11$	36	$N(N+1)(N+2)(N+8)(N-1)/120$
T_6	$\begin{array}{ c c c c c c } \hline \square & \square & \square & \square & \square & \square \\ \hline \end{array}$	$T_m[6]$	$q = 6 \quad 2$	$\mathbf{q}^p = 6^+ \quad 13$	48	$N(N+1)(N+2)(N+3)(N+10)(N-1)/720$
H_5	$\begin{array}{ c c c c } \hline \square & \square & \square & \square \\ \hline \end{array}$	$H_m[5]$		$\mathbf{q}^p = 4^- \quad 9$	49	$N(N+2)(N+6)(N-1)(N-2)/30$
$Y_{3,2}$	$\begin{array}{ c c c } \hline \square & \square & \square \\ \hline \end{array}$	$YT[\{3,2\}]$			$12 + \bar{12}$	$N(N+2)(N+4)(N-1)(N-3)/24$
$Y_{4,2}$	$\begin{array}{ c c c c } \hline \square & \square & \square & \square \\ \hline \end{array}$	$YT[\{4,2\}]$			$21 + \bar{21}$	$N^2(N+3)(N+6)(N-1)(N-3)/80$

Note that the dimension of the T_m irrep is $\frac{(2m+n-2)\Gamma(m+n-2)}{\Gamma(m+1)\Gamma(n-1)}$.

B Extended character decomposition

For an $O(N)$ irrep labelled by Young tableau Y_λ , $\lambda = (\lambda_1, \dots, \lambda_r)$, the characters can be computed using equation (A.7) of [129] (based on [130]). The rest of the computation is described in appendix B of [1]. Other relevant references are appendix A of [3], equation (2.57) of [131] (for the extraction of singlets).

We reproduce here the scalar spectrum:

$$\begin{aligned}
Z_0 = & S + \Delta V \phi + \Delta^2(S + T)\phi^2 + \Delta^3(V_{\text{EOM}} + T_3)\phi^3 + \Delta^4(S + T + T_4)\phi^4 \\
& + \Delta^5(V + T_3 + T_5)\phi^5 \\
& + \Delta^6[(S + T + B_4)\phi^4 + (S + T + T_4 + T_6)\phi^6] \\
& + \Delta^7[(V + H_3 + T_3 + Y_{3,2})\phi^5 + (V + T_3 + T_5 + T_7)\phi^7] \\
& + \Delta^8[(2S + 2T + T_4 + B_4)\phi^4 + (S + 2T + H_4 + T_4 + B_4 + Y_{4,2})\phi^6 \\
& \quad + (S + T + T_4 + T_6 + T_8)\phi^8] \\
& + \Delta^9[(4V + 3T_3 + T_5 + 3H_3 + H_5 + Y_{3,2} + Y_{2,2,1})\phi^5 \\
& \quad + (V + 2T_3 + T_5 + H_3 + H_5 + Y_{3,2} + Y_{5,2})\phi^7 + (V + T_3 + T_5 + T_7 + T_9)\phi^9] \\
& + \Delta^{10}[(2S + 2T + T_4 + A_4 + B_4)\phi^4 \\
& \quad + (4S + 8T + 4T_4 + T_6 + 2A + 4B_4 + 4H_4 + H_6 + 2Y_{4,2} + Y_{2,1,1} + Y_{2,2,2} + Y_{3,2,1})\phi^6 \\
& \quad + (S + 2T + 2T_4 + T_6 + H_4 + H_6 + B_4 + Y_{4,2} + Y_{6,2})\phi^8 \\
& \quad + (S + T + T_4 + T_6 + T_8 + T_{10})\phi^{10}] \\
& + O(\Delta^{11})
\end{aligned} \tag{B.1}$$

Finite- N effects can be checked by replacing the general characters by those for an $O(N)$ group with finite rank. In this way, one can reproduce some identifies from [129], and generate new. We have not implemented this systematically. Examples are

$$(2, 2)|_{N=2} = -(2) \tag{B.2}$$

$$(4, 2)|_{N=2} = -(4) \tag{B.3}$$

where the latter was confirmed in private communication with W. Cao and T. Melia.

References

- [1] J. Henriksson, *The critical $O(N)$ CFT: Methods and conformal data*, *Phys. Rept.* **1002** (2023) 1–72, [[2201.09520^{PDF}](#)], [[INSPIRE](#)].
- [2] *Computed for this document.*
- [3] W. Cao, F. Herzog, T. Melia and J. Roosmale Nepveu, *Renormalization and non-renormalization of scalar EFTs at higher orders*, *JHEP* **09** (2021) 014, [[2105.12742^{PDF}](#)], [[INSPIRE](#)].
- [4] J. Roosmale Nepveu, *Renormalization and the Double Copy of Effective Field Theories*. PhD thesis, Humboldt University, Berlin, 2024. [[INSPIRE](#)]. [DOI:10.18452/29107](#).
- [5] J. Henriksson, F. Herzog, S. R. Kousvos and J. Roosmale Nepveu, *Multiloop spectra in general scalar EFTs and CFTs*, *Phys. Lett. B* **874** (2026) 140235, [[2507.12518^{PDF}](#)], [[INSPIRE](#)].
- [6] J. Henriksson, S. R. Kousvos and J. Roosmale Nepveu, *EFT meets CFT: Multiloop renormalization of higher-dimensional operators in general ϕ^4 theories*, *JHEP* **05** (2026) 005, [[2511.16740^{PDF}](#)], [[INSPIRE](#)].
- [7] F. Bertucci, J. Henriksson and B. McPeak, *Analytic bootstrap of mixed correlators in the $O(n)$ CFT*, *JHEP* **10** (2022) 104, [[2205.09132^{PDF}](#)], [[INSPIRE](#)].
- [8] O. Schnetz, *ϕ^4 theory at seven loops*, *Phys. Rev. D* **107** (2023) 036002, [[2212.03663^{PDF}](#)], [[INSPIRE](#)].
- [9] O. Diatlyk, F. K. Popov and Y. Wang, *Beyond $N = \infty$ in large N conformal vector models at finite temperature*, *JHEP* **08** (2024) 219, [[2309.02347^{PDF}](#)], [[INSPIRE](#)].
- [10] B.-X. Lao and S. Rychkov, *3D Ising CFT and exact diagonalization on icosahedron: The power of conformal perturbation theory*, *SciPost Phys.* **15** (2023) 243, [[2307.02540^{PDF}](#)], [[INSPIRE](#)].
- [11] G. Cuomo, J. M. V. P. Lopes, J. Matos, J. Oliveira and J. Penedones, *Numerical tests of the large charge expansion*, *JHEP* **05** (2024) 161, [[2305.00499^{PDF}](#)], [[INSPIRE](#)].
- [12] M. Hasenbusch, *Cubic fixed point in three dimensions: Monte Carlo simulations of the ϕ^4 model on the simple cubic lattice*, *Phys. Rev. B* **107** (2023) 024409, [[2211.16170^{PDF}](#)], [[INSPIRE](#)].
- [13] W. Zhu, C. Han, E. Huffman, J. S. Hofmann and Y.-C. He, *Uncovering conformal symmetry in the 3D Ising transition: State-operator correspondence from a fuzzy sphere regularization*, *Phys. Rev. X* **13** (2023) 021009, [[2210.13482^{PDF}](#)], [[INSPIRE](#)].
- [14] L. Hu, Y.-C. He and W. Zhu, *Operator product expansion coefficients of the 3D Ising criticality via quantum fuzzy spheres*, *Phys. Rev. Lett.* **131** (2023) 031601, [[2303.08844^{PDF}](#)], [[INSPIRE](#)].
- [15] C. Han, L. Hu, W. Zhu and Y.-C. He, *Conformal four-point correlators of the three-dimensional Ising transition via the quantum fuzzy sphere*, *Phys. Rev. B* **108** (2023) 235123, [[2306.04681^{PDF}](#)], [[INSPIRE](#)].
- [16] J. S. Hofmann, F. Goth, W. Zhu, Y.-C. He and E. Huffman, *Quantum Monte Carlo simulation*

- of the 3D Ising transition on the fuzzy sphere, *SciPost Phys. Core* **7** (2024) 028, [2310.19880^{PDF}], [INSPIRE].
- [17] A. M. Läuchli, L. Herviou, P. H. Wilhelm and S. Rychkov, *Exact diagonalization, matrix product states and conformal perturbation theory study of a 3D Ising fuzzy sphere model*, *SciPost Phys.* **19** (2025) 076, [2504.00842^{PDF}], [INSPIRE].
 - [18] A. Läuchli, “Fuzzy Sphere Study of the 3D $O(2)$ CFT: Spectrum, Finite size corrections and some OPE coefficients.” Talk at “Fuzzy Sphere meets Bootstrap,” Simons Center, 7 November 2023, 2023.
 - [19] C. Han, L. Hu and W. Zhu, *Conformal operator content of the Wilson-Fisher transition on fuzzy sphere bilayers*, *Phys. Rev. B* **110** (2024) 115113, [2312.04047^{PDF}], [INSPIRE].
 - [20] A. Dey, L. Herviou, C. Mudry and A. M. Läuchli, *Conformal data for the $O(3)$ Wilson-Fisher CFT from fuzzy sphere realization of quantum rotor model*, *Arxiv preprint* (2025), [2510.09755^{PDF}], [INSPIRE].
 - [21] D. Poland, V. Prilepina and P. Tadić, *The five-point bootstrap*, *JHEP* **10** (2023) 153, [2305.08914^{PDF}], [INSPIRE].
 - [22] D. Poland, V. Prilepina and P. Tadić, *Improving the five-point bootstrap*, *JHEP* **05** (2024) 299, [2312.13344^{PDF}], [INSPIRE].
 - [23] C.-H. Chang, V. Dommes, R. S. Erramilli, A. Homrich, P. Kravchuk, A. Liu et al., *Bootstrapping the 3d Ising stress tensor*, *JHEP* **03** (2025) 136, [2411.15300^{PDF}], [INSPIRE].
 - [24] D. Poland, V. Prilepina and P. Tadić, *Mixed five-point correlators in the 3d Ising model*, *JHEP* **10** (2025) 237, [2507.01223^{PDF}], [INSPIRE].
 - [25] F. De Cesare and S. Rychkov, *Disturbing news about the $d = 2 + \epsilon$ expansion*, *Prog. Theor. Exp. Phys.* **2025** (2025) 093B02, [2505.21611^{PDF}], [INSPIRE].
 - [26] A. M. Shalaby, *Critical exponents of the $O(N)$ -symmetric ϕ^4 model from the ϵ^7 hypergeometric-Meijer resummation*, *Eur. Phys. J. C* **81** (2021) 87, [2005.12714^{PDF}], [INSPIRE].
 - [27] D. P. Belanger and H. Yoshizawa, *Neutron scattering and the critical behavior of the three-dimensional Ising antiferromagnet FeF_2* , *Phys. Rev. B* **35** (1987) 4823–4830, [INSPIRE].
 - [28] P. Damay, F. Leclercq, R. Magli, F. Formisano and P. Lindner, *Universal critical-scattering function: An experimental approach*, *Phys. Rev. B* **58** (1998) 12038–12043.
 - [29] D. M. Sullivan, G. W. Neilson, H. E. Fischer and A. R. Rennie, *Small angle neutron scattering from D_2O in the critical region*, *J. Phys.: Condens. Matter* **12** (2000) 3531–3542.
 - [30] J. V. Sengers and J. G. Shanks, *Experimental critical-exponent values for fluids*, *J. Stat. Phys.* **137** (2009) 857–877.
 - [31] A. Lytle and D. T. Jacobs, *Turbidity determination of the critical exponent η in the liquid–liquid mixture methanol and cyclohexane*, *J. Chem. Phys.* **120** (2004) 5709–5716.

- [32] S. Ebadi et al., *Quantum phases of matter on a 256-atom programmable quantum simulator*, *Nature* **595** (2021) 227, [[2012.12281^{PDF}](#)], [[INSPIRE](#)].
- [33] M. Campostrini, A. Pelissetto, P. Rossi and E. Vicari, *25th order high temperature expansion results for three-dimensional Ising like systems on the simple cubic lattice*, *Phys. Rev. E* **65** (2002) 066127, [[cond-mat/0201180^{PDF}](#)], [[INSPIRE](#)].
- [34] M. Hasenbusch, *Restoring isotropy in a three-dimensional lattice model: The Ising universality class*, *Phys. Rev. B* **104** (2021) 014426, [[2105.09781^{PDF}](#)], [[INSPIRE](#)].
- [35] F. Kos, D. Poland, D. Simmons-Duffin and A. Vichi, *Precision islands in the Ising and $O(N)$ models*, *JHEP* **08** (2016) 036, [[1603.04436^{PDF}](#)], [[INSPIRE](#)].
- [36] E. Brezin, J. L. Guillou, J. Zinn-Justin and B. Nickel, *Higher order contributions to critical exponents*, *Phys. Lett. A* **44** (1973) 227–228, [[INSPIRE](#)].
- [37] K. G. Chetyrkin, A. L. Kataev and F. V. Tkachov, *Five loop calculations in the $g\phi^4$ model and the critical index η* , *Phys. Lett. B* **99** (1981) 147, [[INSPIRE](#)]. [Erratum: *Phys. Lett. B* **101** (1981) 457].
- [38] H. Kleinert, J. Neu, V. Schulte-Frohlinde, K. G. Chetyrkin and S. A. Larin, *Five loop renormalization group functions of $O(n)$ symmetric ϕ^4 theory and epsilon expansions of critical exponents up to ϵ^5* , *Phys. Lett. B* **272** (1991) 39–44, [[hep-th/9503230^{PDF}](#)], [[INSPIRE](#)]. [Erratum: *Phys. Lett. B* **319** (1993) 545].
- [39] D. V. Batkovich, K. G. Chetyrkin and M. V. Kompaniets, *Six loop analytical calculation of the field anomalous dimension and the critical exponent η in $O(n)$ -symmetric φ^4 model*, *Nucl. Phys. B* **906** (2016) 147–167, [[1601.01960^{PDF}](#)], [[INSPIRE](#)].
- [40] O. Schnetz, *Numbers and functions in quantum field theory*, *Phys. Rev. D* **97** (2018) 085018, [[1606.08598^{PDF}](#)], [[INSPIRE](#)].
- [41] O. Schnetz, “Eightloop gamma in ϕ^4 .” Emmy Noether seminar, Erlangen (Germany), April 30 2021.
- [42] R. Abe, *Critical exponent η up to $1/n^2$ for the three-dimensional system with short-range interaction*, *Prog. Theor. Phys.* **49** (1973) 1877–1888.
- [43] K. Symanzik, “Unpublished.” Cited in [44].
- [44] I. Kondor, L. Herenyi and T. Temesvari, *Resummation of the $1/n$ expansion through a selfconsistent approach*, *Phys. Rev. B* **22** (1980) 1451–1458, [[INSPIRE](#)].
- [45] A. N. Vasiliev, Yu. M. Pismak and J. R. Honkonen, *$1/n$ expansion: Calculation of the exponent η in the order $1/n^3$ by the conformal bootstrap method*, *Theor. Math. Phys.* **50** (1982) 127–134, [[INSPIRE](#)]. [*Teor. Mat. Fiz.* **50** (1982) 195].
- [46] D. Kazakov, O. Tarasov and A. Vladimirov, *Calculation of critical exponents by quantum field theory methods*, *Sov. Phys. JETP* **50** (1979) 521, [[INSPIRE](#)]. [*Zh. Eksp. Teor. Fiz.* **77** (1979) 1035].

- [47] M. V. Kompaniets and E. Panzer, *Minimally subtracted six loop renormalization of $O(n)$ -symmetric ϕ^4 theory and critical exponents*, *Phys. Rev. D* **96** (2017) 036016, [[1705.06483^{PDF}](#)], [[INSPIRE](#)].
- [48] Y. Okabe and M. Oku, *$1/n$ expansion up to order $1/n^2$ III. Critical exponents γ and ν for $d = 3$* , *Prog. Theor. Phys.* **60** (1978) 1287–1297, [[INSPIRE](#)].
- [49] A. N. Vasiliev, Yu. M. Pismak and J. R. Honkonen, *$1/n$ expansion: Calculation of the exponents η and ν in the order $1/n^2$ for arbitrary number of dimensions*, *Theor. Math. Phys.* **47** (1981) 465–475, [[INSPIRE](#)]. [*Teor. Mat. Fiz.* **47** (1981) 291].
- [50] S.-k. Ma, *Scaling variables and dimensions*, *Phys. Rev. A* **10** (1974) 1818–1836, [[INSPIRE](#)].
- [51] D. J. Broadhurst, J. Gracey and D. Kreimer, *Beyond the triangle and uniqueness relations: Nonzeta counterterms at large N from positive knots*, *Z. Phys. C* **75** (1997) 559–574, [[hep-th/9607174^{PDF}](#)], [[INSPIRE](#)].
- [52] J. Gracey, *Progress with large N_f β functions*, *Nucl. Instrum. Meth. A* **389** (1997) 361–364, [[hep-ph/9609409^{PDF}](#)], [[INSPIRE](#)].
- [53] K. G. Wilson, *Feynman graph expansion for critical exponents*, *Phys. Rev. Lett.* **28** (1972) 548–551, [[INSPIRE](#)].
- [54] Y. Yamazaki, *The cross-over exponent up to ϵ^3 ($\epsilon = 4 - d$) by using Callan-Symanzik equations*, *Phys. Lett. A* **49** (1974) 215 – 217.
- [55] J. E. Kirkham, *Calculation of crossover exponent from Heisenberg to Ising behaviour using the fourth-order ϵ expansion*, *J. Phys. A* **14** (1981) L437, [[INSPIRE](#)].
- [56] M. Kompaniets and K. J. Wiese, *Fractal dimension of critical curves in the $O(n)$ -symmetric ϕ^4 model and crossover exponent at 6-loop order: Loop-erased random walks, self-avoiding walks, Ising, XY, and Heisenberg models*, *Phys. Rev. E* **101** (2020) 012104, [[1908.07502^{PDF}](#)], [[INSPIRE](#)].
- [57] J. A. Gracey, *Crossover exponent in $O(N)$ ϕ^4 theory at $O(1/N^2)$* , *Phys. Rev. E* **66** (2002) 027102, [[cond-mat/0206098^{PDF}](#)], [[INSPIRE](#)].
- [58] H. Kleinert and V. Schulte-Frohlinde, *Exact five loop renormalization group functions of ϕ^4 theory with $O(N)$ -symmetric and cubic interactions: Critical exponents up to ϵ^5* , *Phys. Lett. B* **342** (1995) 284–296, [[cond-mat/9503038^{PDF}](#)], [[INSPIRE](#)].
- [59] P. Calabrese, A. Pelissetto and E. Vicari, *Multicritical phenomena in $O(n_1) \oplus O(n_2)$ symmetric theories*, *Phys. Rev. B* **67** (2003) 054505, [[cond-mat/0209580^{PDF}](#)], [[INSPIRE](#)].
- [60] A. Bednyakov and A. Pikelner, *Six-loop beta functions in general scalar theory*, *JHEP* **04** (2021) 233, [[2102.12832^{PDF}](#)], [[INSPIRE](#)].
- [61] S.-k. Ma, *The $1/n$ expansion*, in *Phase Transitions and Critical Phenomena*, vol. 6 (C. Domb and M. S. Green, eds.). Academic Press, 1976.
- [62] S. E. Derkachov and A. N. Manashov, *The simple scheme for the calculation of the anomalous*

- dimensions of composite operators in the $1/N$ expansion, *Nucl. Phys. B* **522** (1998) 301–320, [[hep-th/9710015^{PDF}](#)], [[INSPIRE](#)].
- [63] A. Cappelletti, D. Friedan and J. I. Latorre, *c-theorem and spectral representation*, *Nucl. Phys. B* **352** (1991) 616–670, [[INSPIRE](#)].
- [64] P. Dey, A. Kaviraj and A. Sinha, *Mellin space bootstrap for global symmetry*, *JHEP* **07** (2017) 019, [[1612.05032^{PDF}](#)], [[INSPIRE](#)].
- [65] J. Henriksson and M. van Loon, *Critical $O(N)$ model to order ϵ^4 from analytic bootstrap*, *J. Phys. A* **52** (2019) 025401, [[1801.03512^{PDF}](#)], [[INSPIRE](#)].
- [66] K. Lang and W. Rühl, *Critical non-linear $O(N)$ σ -models at $2 < d < 4$: The degeneracy of quasi-primary fields and its resolution*, *Z. Phys. C* **61** (1994) 495–510, [[INSPIRE](#)].
- [67] A. Petkou, *Conserved currents, consistency relations and operator product expansions in the conformally invariant $O(N)$ vector model*, *Annals Phys.* **249** (1996) 180–221, [[hep-th/9410093^{PDF}](#)], [[INSPIRE](#)].
- [68] M.-C. Cha, M. P. A. Fisher, S. M. Girvin, M. Wallin and A. P. Young, *Universal conductivity of two-dimensional films at the superconductor-insulator transition*, *Phys. Rev. B* **44** (1991) 6883–6902.
- [69] K. Lang and W. Rühl, *The critical $O(N)$ sigma model at dimensions $2 < d < 4$: A list of quasi-primary fields*, *Nucl. Phys. B* **402** (1993) 573–603, [[INSPIRE](#)].
- [70] H. Kleinert and V. Schulte-Frohlinde, *Critical Properties of ϕ^4 -Theories*. World Scientific Publishing, 2001, [DOI:10.1142/4733](#), [[INSPIRE](#)].
- [71] A. N. Vasiliev and A. S. Stepanenko, *A method of calculating the critical dimensions of composite operators in the massless nonlinear sigma model*, *Theor. Math. Phys.* **94** (1993) 471–481, [[INSPIRE](#)]. [*Teor. Mat. Fiz.* **95** (1993) 160].
- [72] D. J. Broadhurst and A. V. Kotikov, *Compact analytical form for nonzeta terms in critical exponents at order $1/N^3$* , *Phys. Lett. B* **441** (1998) 345–353, [[hep-th/9612013^{PDF}](#)], [[INSPIRE](#)].
- [73] L. Fei, S. Giombi and I. R. Klebanov, *Critical $O(N)$ models in $6 - \epsilon$ dimensions*, *Phys. Rev. D* **90** (2014) 025018, [[1404.1094^{PDF}](#)], [[INSPIRE](#)].
- [74] E. Brezin, J. C. Le Guillou and J. Zinn-Justin, *Field theoretical approach to critical phenomena*, in *Phase Transitions and Critical Phenomena*, vol. 6 (C. Domb and M. S. Green, eds.). Academic Press, 1976.
- [75] I. Jack and H. Osborn, *Background field calculations in curved space-time 1. General formalism and application to scalar fields*, *Nucl. Phys. B* **234** (1984) 331–364, [[INSPIRE](#)].
- [76] S. M. Chester, W. Landry, J. Liu, D. Poland, D. Simmons-Duffin, N. Su et al., *Carving out OPE space and precise $O(2)$ model critical exponents*, *JHEP* **06** (2020) 142, [[1912.03324^{PDF}](#)], [[INSPIRE](#)].
- [77] S. M. Chester, W. Landry, J. Liu, D. Poland, D. Simmons-Duffin, N. Su et al., *Bootstrapping*

- Heisenberg magnets and their cubic instability*, *Phys. Rev. D* **104** (2021) 105013, [[2011.14647^{PDF}](#)], [[INSPIRE](#)].
- [78] M. Reehorst, *Rigorous bounds on irrelevant operators in the 3d Ising model CFT*, *JHEP* **09** (2022) 177, [[2111.12093^{PDF}](#)], [[INSPIRE](#)].
- [79] J. Liu, D. Meltzer, D. Poland and D. Simmons-Duffin, *The Lorentzian inversion formula and the spectrum of the 3d $O(2)$ CFT*, *JHEP* **09** (2020) 115, [[2007.07914^{PDF}](#)], [[INSPIRE](#)]. [Erratum: *JHEP* **01** (2021) 206].
- [80] S. M. Chester, W. Landry, J. Liu, D. Poland, D. Simmons-Duffin, N. Su et al., *Unpublished*, .
- [81] N. Clisby, *Scale-free Monte Carlo method for calculating the critical exponent γ of self-avoiding walks*, *J. Phys. A* **50** (2017) 264003, [[1701.08415^{PDF}](#)], [[INSPIRE](#)].
- [82] Q.-Q. Liu, Y. Deng, T. M. Garoni and H. W. J. Blöte, *The $O(n)$ loop model on a three-dimensional lattice*, *Nucl. Phys. B* **859** (2012) 107–128, [[1112.5647^{PDF}](#)], [[INSPIRE](#)].
- [83] M. Hasenbusch, *Monte Carlo study of an improved clock model in three dimensions*, *Phys. Rev. B* **100** (2019) 224517, [[1910.05916^{PDF}](#)], [[INSPIRE](#)].
- [84] M. Hasenbusch, *Monte Carlo study of a generalized icosahedral model on the simple cubic lattice*, *Phys. Rev. B* **102** (2020) 024406, [[2005.04448^{PDF}](#)], [[INSPIRE](#)].
- [85] N. Clisby and B. Dünweg, *High-precision estimate of the hydrodynamic radius for self-avoiding walks*, *Phys. Rev. E* **94** (2016) 052102, [[2001.03138^{PDF}](#)].
- [86] P. Belohorec, *Renormalization Group Calculation of the Universal Critical Exponents of a Polymer Molecule*. PhD thesis, University of Guelph, 1997.
- [87] M. Hasenbusch, *Finite size scaling study of lattice models in the three-dimensional Ising universality class*, *Phys. Rev. B* **82** (2010) 174433, [[1004.4486^{PDF}](#)], [[INSPIRE](#)].
- [88] F. Winter, W. Janke and A. M. J. Schakel, *Geometric properties of the three-dimensional Ising and XY models*, *Phys. Rev. E* **77** (2008) 061108, [[0803.2177^{PDF}](#)].
- [89] M. Hasenbusch and E. Vicari, *Anisotropic perturbations in three-dimensional $O(N)$ -symmetric vector models*, *Phys. Rev. B* **84** (2011) 125136, [[1108.0491^{PDF}](#)].
- [90] S. E. Derkachov and A. N. Manashov, *On the stability problem in the $O(N)$ nonlinear sigma model*, *Phys. Rev. Lett.* **79** (1997) 1423–1427, [[hep-th/9705020^{PDF}](#)], [[INSPIRE](#)].
- [91] O. Antipin, J. Bersini, J. Hafjall, G. Muco and F. Sannino, *Exact results for the spectrum of the Ising conformal field theory*, *Arxiv preprint* (2025), [[2511.08276^{PDF}](#)], [[INSPIRE](#)].
- [92] K. G. Wilson and J. B. Kogut, *The renormalization group and the ϵ expansion*, *Phys. Rept.* **12** (1974) 75–199, [[INSPIRE](#)].
- [93] S. E. Derkachov, J. A. Gracey and A. N. Manashov, *Four loop anomalous dimensions of gradient operators in ϕ^4 theory*, *Eur. Phys. J. C* **2** (1998) 569–579, [[hep-ph/9705268^{PDF}](#)], [[INSPIRE](#)].

- [94] S. Kehrein, F. Wegner and Yu. Pismak, *Conformal symmetry and the spectrum of anomalous dimensions in the N vector model in $4 - \epsilon$ dimensions*, *Nucl. Phys. B* **402** (1993) 669–692, [[INSPIRE](#)].
- [95] S. K. Kehrein and F. Wegner, *The structure of the spectrum of anomalous dimensions in the N -vector model in $4 - \epsilon$ dimensions*, *Nucl. Phys. B* **424** (1994) 521–546, [[hep-th/9405123](#)]^{PDF}, [[INSPIRE](#)].
- [96] G. Guedes, J. Henriksson, S. R. Kousvos and J. Roosmale Nepveu, *Anomalous dimensions of operators with any number of fields*, *Work in progress* (2026) .
- [97] D. Simmons-Duffin, *The lightcone bootstrap and the spectrum of the 3d Ising CFT*, *JHEP* **03** (2017) 086, [[1612.08471](#)]^{PDF}, [[INSPIRE](#)].
- [98] F. C. Zhang and R. K. P. Zia, *A correction-to-scaling critical exponent for fluids at order ϵ^3* , *J. Phys. A* **15** (1982) 3303–3305.
- [99] S. Harris, A. Kaviraj, J. A. Mann, L. Quintavalle and V. Schomerus, *Comb channel lightcone bootstrap: triple-twist anomalous dimensions*, *JHEP* **08** (2024) 122, [[2401.10986](#)]^{PDF}, [[INSPIRE](#)].
- [100] S. K. Kehrein, *The spectrum of critical exponents in $(\phi^2)^2$ theory in $d = 4 - \epsilon$ dimensions: Resolution of degeneracies and hierarchical structures*, *Nucl. Phys. B* **453** (1995) 777–806, [[hep-th/9507044](#)]^{PDF}, [[INSPIRE](#)].
- [101] A. Dymarsky, F. Kos, P. Kravchuk, D. Poland and D. Simmons-Duffin, *The 3d stress-tensor bootstrap*, *JHEP* **02** (2018) 164, [[1708.05718](#)]^{PDF}, [[INSPIRE](#)].
- [102] R. Gopakumar, A. Kaviraj, K. Sen and A. Sinha, *A Mellin space approach to the conformal bootstrap*, *JHEP* **05** (2017) 027, [[1611.08407](#)]^{PDF}, [[INSPIRE](#)].
- [103] D. Carmi, J. Penedones, J. A. Silva and A. Zhiboedov, *Applications of dispersive sum rules: ϵ -expansion and holography*, *SciPost Phys.* **10** (2021) 145, [[2009.13506](#)]^{PDF}, [[INSPIRE](#)].
- [104] F. Gliozzi, A. L. Guerrieri, A. C. Petkou and C. Wen, *The analytic structure of conformal blocks and the generalized Wilson-Fisher fixed points*, *JHEP* **04** (2017) 056, [[1702.03938](#)]^{PDF}, [[INSPIRE](#)].
- [105] A. Codello, M. Safari, G. P. Vacca and O. Zanusso, *Leading CFT constraints on multi-critical models in $d > 2$* , *JHEP* **04** (2017) 127, [[1703.04830](#)]^{PDF}, [[INSPIRE](#)].
- [106] L. F. Alday, J. Henriksson and M. van Loon, *Taming the ϵ -expansion with large spin perturbation theory*, *JHEP* **07** (2018) 131, [[1712.02314](#)]^{PDF}, [[INSPIRE](#)].
- [107] J. Henriksson, *Analytic Bootstrap for Perturbative Conformal Field Theories*. DPhil thesis, University of Oxford, 2020. [[2008.12600](#)]^{PDF}.
- [108] F. J. Wegner, *Critical exponents in isotropic spin systems*, *Phys. Rev. B* **6** (1972) 1891–1893, [[INSPIRE](#)].
- [109] O. Antipin, J. Bersini, J. Hafjall, G. Muco and F. Sannino, *Semiclassical canovaccio for composite operators*, *Arxiv preprint* (2025), [[2512.23539](#)]^{PDF}, [[INSPIRE](#)].

- [110] K. Lang and W. Rühl, *The critical $O(N)$ σ -model at dimensions $2 < d < 4$: Fusion coefficients and anomalous dimensions*, *Nucl. Phys. B* **400** (1993) 597–623, [INSPIRE].
- [111] A. Houghton and F. J. Wegner, *Feynman-graph calculation of the $(0, l)$ critical exponents to order ϵ^2* , *Phys. Rev. A* **10** (1974) 435–437.
- [112] O. Antipin, J. Bersini, F. Sannino, Z.-W. Wang and C. Zhang, *Charging the $O(N)$ model*, *Phys. Rev. D* **102** (2020) 045011, [2003.13121^{PDF}], [INSPIRE].
- [113] A. Bednyakov and A. Pikelner, *Six-loop anomalous dimension of the ϕ^Q operator in the $O(N)$ symmetric model*, *Phys. Rev. D* **106** (2022) 076015, [2208.04612^{PDF}], [INSPIRE].
- [114] K. Lang and W. Rühl, *Field algebra for critical $O(N)$ vector nonlinear sigma models at $2 < d < 4$* , *Z. Phys. C* **50** (1991) 285–292, [INSPIRE].
- [115] K. Lang and W. Rühl, *Anomalous dimensions of tensor fields of arbitrary rank for critical nonlinear $O(N)$ sigma models at $2 < d < 4$ to first order in $1/N$* , *Z. Phys. C* **51** (1991) 127–132, [INSPIRE].
- [116] A. N. Manashov, E. D. Skvortsov and M. Strohmaier, *Higher spin currents in the critical $O(N)$ vector model at $1/N^2$* , *JHEP* **08** (2017) 106, [1706.09256^{PDF}], [INSPIRE].
- [117] V. M. Braun and A. N. Manashov, *Evolution equations beyond one loop from conformal symmetry*, *Eur. Phys. J. C* **73** (2013) 2544, [1306.5644^{PDF}], [INSPIRE].
- [118] S. E. Derkachov, S. K. Kehrein and A. N. Manashov, *High-gradient operators in the N -vector model*, *Nucl. Phys. B* **493** (1997) 660–682, [cond-mat/9610106^{PDF}], [INSPIRE].
- [119] E. E. Jenkins, A. V. Manohar, L. Naterop and J. Pagès, *Two loop renormalization of scalar theories using a geometric approach*, *JHEP* **02** (2024) 131, [2310.19883^{PDF}], [INSPIRE].
- [120] D. J. Wallace and R. K. P. Zia, *Harmonic perturbations of generalized Heisenberg spin systems*, *J. Phys. C* **8** (1975) 839.
- [121] A. Dymarsky, J. Penedones, E. Trevisani and A. Vichi, *Charting the space of 3D CFTs with a continuous global symmetry*, *JHEP* **05** (2019) 098, [1705.04278^{PDF}], [INSPIRE].
- [122] L. F. Alday, J. Henriksson and M. van Loon, *An alternative to diagrams for the critical $O(N)$ model: dimensions and structure constants to order $1/N^2$* , *JHEP* **01** (2020) 063, [1907.02445^{PDF}], [INSPIRE].
- [123] J. Padayasi, A. Krishnan, M. A. Metlitski, I. A. Gruzberg and M. Meineri, *The extraordinary boundary transition in the 3d $O(N)$ model via conformal bootstrap*, *SciPost Phys.* **12** (2022) 190, [2111.03071^{PDF}], [INSPIRE].
- [124] K. Lang and W. Rühl, *The critical $O(N)$ σ -model at dimension $2 < d < 4$ and order $1/N^2$: Operator product expansions and renormalization*, *Nucl. Phys. B* **377** (1992) 371–401, [INSPIRE].
- [125] M. Goykhman and M. Smolkin, *Vector model in various dimensions*, *Phys. Rev. D* **102** (2020) 025003, [1911.08298^{PDF}], [INSPIRE].

- [126] N. Chai, M. Goykhman and R. Sinha, *Conformal correlators in the critical $O(N)$ vector model*, *Phys. Rev. D* **105** (2022) 086026, [[2103.10168^{PDF}](#)], [[INSPIRE](#)].
- [127] J. Csipes and P. Vařko, *$O(N)$ BCFT: new data from conformal partial wave expansions*, *Arxiv preprint* (2026), [[2606.14733^{PDF}](#)], [[INSPIRE](#)].
- [128] M. Reehorst, E. Trevisani and A. Vichi, *Mixed scalar-current bootstrap in three dimensions*, *JHEP* **12** (2020) 156, [[1911.05747^{PDF}](#)], [[INSPIRE](#)].
- [129] W. Cao, X. Lu and T. Melia, *Constraints on the spectrum of field theories with non-integer $O(N)$ symmetry from quantum evanescence*, *SciPost Phys. Core* **7** (2024) 055, [[2312.10139^{PDF}](#)], [[INSPIRE](#)].
- [130] K. Koike and I. Terada, *Young-diagrammatic methods for the representation theory of the classical groups of type B_n , C_n , D_n* , *J. Algebra* **107** (1987) 466–511.
- [131] B. Grinstein, X. Lu, L. Merlo and P. Quilez, *Hilbert series for covariants and their applications to minimal flavor violation*, *JHEP* **2024** (2024) 154, [[2312.13349^{PDF}](#)], [[INSPIRE](#)].
[Erratum: *JHEP* **03** (2025) 072].