

Notation Comments

Remark. The notation $L_A : \mathbb{F}^n \rightarrow F^m$ when $A \in M_n(\mathbb{F})$ has the letter L to indicate left multiplication by A on column vectors.

Remark. Given any linear operator $T : V \rightarrow V$, and finite ordered bases B, C for V . The matrix of T with respect to B and C is denoted $[T]_B^C$. In particular,

$$[Id_v]_C^B = \left([Id_v]_b^C\right)^{-1} \quad (1)$$

From this,

$$[T]_C^C = [Id_v]_B^C [T]_B^B [Id_v]_C^B \quad (2)$$

$$= Q^{-1} [T]_B^C Q \quad (3)$$

where $Q = [Id_v]_C^B$ is the change of basis matrix.

Cosets

If U is a subspace of V and $v \in V$ then the left coset of U in V represented by v is

$$v + U = \{v + u | u \in U\} \quad (4)$$

The **set of left cosets** of U in V is

$$V/U = \{v + U | v \in V\} \quad (5)$$

Note that if $v \in V$ and $u \in U$ then $v + U = (v + u) + U$. Naively, we could hope that

$$\begin{aligned} V/U \times V/U &\rightarrow V/U \\ (v_1 + U, v_2 + U) &\mapsto (v_1 + v_2) + U \end{aligned}$$

actually defines a function. We have to be sure that when you choose some v'_1, v'_2 that the resulting coset is the same... i.e. that we need to check that this really is a function for which inputs have exactly one output. That is,

$$(v_1 + U, v_2 + U) \mapsto v_1 + v_2 + U$$

is well-defined, in the sense that the right hand side value is independent of choice of coset representatives of the initial cosets. Here, if $v'_1 = v_1 + u_1$, $v'_2 = v_2 + u_2$ with $u_i \in U$. Now

$$v'_1 + v'_2 + U = \left[(v_1 + u_1) + (v_2 + u_2)\right] + U$$

Thus,

$$\begin{aligned} v'_1 + v'_2 + U &= (v_1 + u_1 + v_2) + (u_2 + U) \\ &= (v_1 + v_2 + u_1) + U \\ &= v_1 + v_2 + U \end{aligned}$$

That is, since addition on V is Abelian, every subgroup U is normal and thus the naive formula does give a well-defined function. We now check if

$$\begin{aligned} \mathbb{F} \times V/U &\rightarrow V/U \\ (\lambda, v + U) &\mapsto \lambda v + U \end{aligned}$$

is a well-defined function (**IT IS**). So the family of cosets of subspace U in vector space V is itself a vector space over \mathbb{F} .

Lemma. *Suppose U is a vector subspace of V , and B is a basis of U . Let $C = B \cup B'$ be any basis of V extending B . Then, $\{v + U | v \in B'\}$ is a basis of our quotient vector space V/U .*

Proof. Suppose $\sum_i \lambda_i(v_i + U) = 0_{V/U}$ for some $\lambda_1, \dots, \lambda_n \in \mathbb{F}$. and $v_1, \dots, v_n \in B'$. Since $0_{V/U} = 0_v + U = U$ is our zero vector, thus

$$\left(\sum_i^n \lambda_i v_i \right) + U = U$$

This holds if and only if

$$\sum_i^n \lambda_i v_i \in U$$

However, the $v_i \in B'$ and hence are linearly independent of the $sp(B)$. Therefore, this linear combination can only be $0_v \in U$. But C is a basis and thus all of the $\lambda_i = 0$. Note if $U = V$ then V/U is only $\{0_v + U\}$ and one uses logical statements. \square