

John Waczak

Mathematica code for SHO based off of Dr. Tate's  
"Shooting Method" used in class 1/10/18

```
In[1]:= SetOptions[Plot,  
PlotStyle → {Blue, AbsoluteThickness[2], Dashed},  
ImageSize → 500, AxesStyle →  
Directive[FontFamily → "Arial", FontSize → 18,  
Black, AbsoluteThickness[0.5], Arrowheads[0.04]]];
```

```
In[11]:= m = 1  
ω = 1  
ħ = 1
```

```
Out[11]= 1
```


```
Out[12]= 1
```

```
Out[13]= 1
```

```
In[14]:= v[x_] := (1/2) * m * ω^2 * x^2;
```

Here we graph the harmonic potential

```
In[28]:= Plot[v[x], {x, -1.5, 1.5},  
          PlotStyle -> {Red, AbsoluteThickness[1]}]
```




Now we choose an energy of 0.5 (because we have sent  $m$ ,  $\omega$ , and  $\hbar$  to 1 we expect the ground state to have an energy of  $\hbar\omega(n + 1/2)$  with  $n=0$ )

```

In[29]:= energy = 0.5;
xMax = 5;
solution = NDSolve[
  {psi''[x] == -2 (energy - v[x]) psi[x], psi[-xMax] == 0,
   psi'[-xMax] == 0.001}, psi, {x, -xMax, xMax}];
Plot[psi[x] /. solution, {x, -xMax, xMax}]

```

Out[32]=




Now we find the second eigenstate to have an energy of 1.5 in our units

```

In[37]:= energy = 1.5;
xMax = 5;
solution = NDSolve[
  {psi''[x] == -2 (energy - v[x]) psi[x], psi[-xMax] == 0,
   psi'[-xMax] == 0.001}, psi, {x, -xMax, xMax}];
Plot[psi[x] /. solution, {x, -xMax, xMax}]

```



Now, going back to the ground state, we can see how sensitive the system is as an increase of one ten thousandth was enough make the solution blow up to infinity.

```

In[53]:= energy = 0.50001;
xMax = 5;
solution = NDSolve[
  {psi''[x] == -2 (energy - v[x]) psi[x], psi[-xMax] == 0,
   psi'[-xMax] == 0.001}, psi, {x, -xMax, xMax}];
Plot[psi[x] /. solution, {x, -xMax, xMax}]

```

