# An Application of the Study Determinant on Skew-Coninvolutory Quaternionic Matrices

University of the Philippines Cebu

John Aaron Q. Alcoseba

 $March\ 9,\ 2017$ 

### Contents

1	Introduction	2
<b>2</b>	Preliminaries	4
3	Results and Discussion	5
4	Summary and Recommendations	6

#### Introduction

Ever since their discovery by William Rowan Hamilton in 1843, Quaternions have found extensive use in solving problems both in theoretical and applied mathematics - particularly on the problem of 3D rotation. Computations regarding the said matter use 4x4 matrices with real entries like the ones shown below.

$$R_x(\alpha) = \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & \cos \alpha & -\sin \alpha & 0 \\ 0 & \sin \alpha & \cos \alpha & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix} \qquad R_y(\beta) = \begin{pmatrix} \cos \beta & 0 & \sin \beta & 0 \\ 0 & 1 & 0 & 0 \\ -\sin \beta & 0 & \cos \beta & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$
$$R_z(\gamma) = \begin{pmatrix} \cos \gamma & -\sin \gamma & 0 & 0 \\ \sin \gamma & \cos \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}$$

Quaternions are used today in robotics, three-dimensional computer graphics, computer vision, crystallographic texture analysis, navigation, and molecular dynamics.

### Preliminaries

#### Results and Discussion

**Summary and Recommendations** 

# Bibliography