


## Lecture 25


Higgs Boson:



47

## b-jet Identification (*b-Tagging*)

Critical as b-jet ubiquitous in higgs final states.



52

# Triggering

- LHC provides orders of magnitude more collisions than we can save to disk.
  - Can only keep 1 out of 40,000 events / Discarded data lost forever
- Interesting physics is incredibly rare:
  - ~1 Higgs per billion events / ~1 Di-Higgs per trillion events

**Triggering:** Process of selecting which collisions to save for further analysis.


## Triggering at the LHC:


- Custom Electronics + Commodity CPU
- Fast processing of images (micro-seconds / seconds)
- Events rate from 40 MHz  $\rightarrow$  1kHz.
- Data rate from 80 TBs (!)  $\rightarrow$  2 GB/s

55

---

To collect data faster, each event has multiple proton collisions. Significantly complicates analysis of events





## Vacuum Fluctuations

QM+Spacetime  $\Rightarrow$  Anti-particles  $\Rightarrow$  Vacuum is interesting place.

Because of QM, need to put in Energy to probe smaller distances.


$$E \cdot t \sim E \cdot x \sim 1 \Rightarrow \text{Small distances} \Rightarrow \text{large } E$$

If  $E \gg 2m_e$  nothing stops you from making  $e^+e^-$  pairs.

So operationally, should think of the vacuum as filled of particle-anti-particle pairs constantly coming in and out of existence:

No meaningful sense in which the vacuum is empty.

### Example 1



### Example 2




diagram gives a correction to the mass of the Z-boson from the top quark