

## JOHN CALABRESE — RESEARCH STATEMENT

**Overview.** I completed my DPhil in 2013 at the University of Oxford, where my advisor was Tom Bridgeland (now at Sheffield). For the remainder of the year, I was an **EPSRC Doctoral Prize Fellow** at Imperial College, mentored by Richard Thomas. In January 2014 I moved to Rice as an RTG Lovett Instructor, mentored by Brendan Hassett (now at Brown). Later that year I was awarded an **AMS-Simons travel grant**. In 2015 I was awarded an **NSF Mathematical Sciences Postdoctoral Research Fellowship**, and I am now also a G.C. Evans Instructor. In 2017 I was awarded an **NSF conference grant** to organize the Texas Algebraic Geometry Symposium. I work in *algebraic geometry*, with a focus on *categories of sheaves* and applications to *Donaldson-Thomas* theory.

**Background.** The classical objects of study in algebraic geometry are (smooth) projective varieties, which are subsets  $X \subset \mathbf{CP}^N$  defined by the vanishing of homogeneous polynomials. On the one hand, a given such  $X$  is a *complex manifold* and can therefore be studied borrowing techniques from *geometry*. On the other hand, since  $X$  is defined by polynomials, one can also employ techniques coming from *algebra*. The overarching theme of my research is to understand the geometry of a variety by means of algebraic *invariants*.

The seminal 1955 paper of Serre [Ser55] introduced the notion of *coherent sheaf*, which is now ubiquitous in the subject. Roughly, a coherent sheaf  $V$  on a variety  $X$  is a collection of vector spaces  $V_x$ , one for each point  $x \in X$ . Unlike vector bundles, the dimension of the fibre  $V_x$  needn't be constant: it is allowed to jump along closed subsets. The collection of all coherent sheaves forms a category,  $\mathbf{Coh}(X)$ . In contrast to vector bundles,  $\mathbf{Coh}(X)$  is a particularly pleasant category, as it is closed under all the operations one wishes to perform: kernels, images, direct sums, etc.

With the advent of *homological algebra*, it was later realized one should enlarge  $\mathbf{Coh}(X) \subset \mathbf{D}(X)$ , passing to its *derived category*. The latter consists of chain complexes of coherent sheaves, providing a more flexible framework for derived functors. For example, derived categories are indispensable to even state the dualities of Grothendieck-Verdier, far-reaching generalizations of *Poincaré duality* (in topology) and *Serre duality* (in algebraic geometry). But it wasn't until Mukai [Muk81], that derived categories became objects of independent study, and were later essential in Kontsevich's *homological mirror symmetry* and Bridgeland's *stability conditions*.

Part of my work is devoted to the study the categories  $\mathbf{Coh}(X)$  and  $\mathbf{D}(X)$ , while another part seeks for applications, specifically to *Donaldson-Thomas* invariants. In his thesis, Thomas introduced virtual counts of sheaves for Calabi-Yau 3-folds (a special class of varieties), marking the birth of Donaldson-Thomas theory. These invariants are closely related to the *curve-counting* invariants of *Gromov-Witten* and to *BPS counts* in mathematical physics.

**Past.** Two varieties  $X, Y$  are called *birational* if there are closed subvarieties  $W \subset X, Z \subset Y$  such that  $X \setminus W$  and  $Y \setminus Z$  are isomorphic. While this notion is somewhat useless in classical topology, it has always played a central role in algebraic geometry (due to the more rigid nature of the Zariski topology). In my thesis, I studied the behavior of Donaldson-Thomas invariants under birational maps [Cal16a].

At its very core, [Cal16a] may be seen as an application of an equivalence of categories of Bridgeland [Bri02], together with the theory of *motivic Hall algebras* of Joyce-Song and Kontsevich-Soibelman [JS12, KS10]. The answer was a simple and explicit formula in the case of *flops* (which

are the building blocks of birational maps between Calabi-Yau 3-folds). The formula and methods of [Cal16a] were applied by Maulik in [Mau16], settling conjectures of Oblomkov-Shende and Diaconescu-Hua-Soibelman on the *HOMFLY polynomial* of an algebraic link.

Related to the flop formula, Bryan-Cadman-Young [BCY12] formulated a *Crepanant Resolution Conjecture* for Donaldson-Thomas invariants. This conjecture predicts a transformation rule for the invariants, in the setting of the *McKay Correspondence* [BKR01]. This conjecture consists of two statements: I proved the first and made progress towards the second in the paper [Cal16b].

A celebrated theorem in algebraic geometry is *Gabriel's theorem*, which says two varieties  $X$  and  $Y$  are isomorphic if and only if the categories  $\mathrm{Coh}(X)$  and  $\mathrm{Coh}(Y)$  are equivalent. In other words,  $\mathrm{Coh}(X)$  is a *complete invariant* of the variety  $X$ . While in Oxford, I started collaborating with Michael Groechenig (now at Freie Universität Berlin), which resulted in the paper [CG15]. We provided a modern approach to Gabriel's theorem, leading to a considerable generalization. The approach of [CG15], later led to [Cal17b, Cal17c, Cal17a] which explore related questions.

As a fellow at Imperial College, I worked with Richard Thomas on applying Kuznetsov's recent theory of *homological projective duality* [Kuz07]. While  $\mathrm{Coh}(X)$  is a complete invariant of  $X$ ,  $\mathrm{D}(X)$  is not. An active area of research aims to understand which varieties  $X, Y$  are *Fourier-Mukai partners*, meaning  $\mathrm{D}(X)$  is equivalent to  $\mathrm{D}(Y)$ . This led to the work [CT16], where we constructed new derived equivalences of Calabi-Yau 3-folds arising from pencils of cubic 4-folds. The survey [Cala] provides an overview of the ideas involved. Work in [CT16] was later shown to be part of a broader picture in [BL16].

**Present.** I am currently collaborating with Roberto Pirisi (UBC), on an exciting variant of Gabriel's reconstruction theorem [CP17]. Using work of Meinhardt-Partsche as a springboard [MP14], we have studied the quotient categories  $\mathrm{Coh}(X)/\mathrm{Coh}_{\leq k}(X)$ . We show that the  $k$ -th quotient controls the isomorphism type of  $X$ , *up to codimension*  $\dim X - k$ . In other words, these quotients interpolate between Gabriel's theorem and *birational geometry*. For example, from the quotient  $\mathrm{Coh}(X)/\mathrm{Coh}_{\leq \dim X - 1}(X)$  one extracts the function field of  $X$ , and hence the birational type of  $X$ . This work is a continuation of the program initiated in [CG15].

A conjecture of Kawamata [Kaw02], stated that a variety has at most finitely many Fourier-Mukai partners. In [AT09], it was shown that a variety admits at most countably many partners. Kawamata's conjecture was recently disproved by Lesieutre [Les15]. However, variations on the theme of Kawamata's conjecture can be explored for (semiorthogonal) *components* of  $\mathrm{D}(X)$ . The study of these components has received renewed attention due to Kuznetsov's theory [Kuz07], mentioned above. Borrowing ideas from my previous work [CT16], I have been working on (infinitesimal) Torelli-type theorems for these components [Calb], akin to [HR16].

**Future.** Continuing from [CP17], I hope to make contact with *non-commutative geometry*, in the sense of Artin, Smith, Van den Bergh et al. Indeed, there are many constructions of *non-commutative projective varieties*, and the work [PVdB16] produces examples of *birational non-commutative surfaces*. The works [CP17, PVdB16] are certainly related, and a close investigation will perhaps also lead to higher dimensional constructions, such as *non-commutative flops*.

Together with Jørgen Rennemo (Oxford) and Sjoerd Beentjes (Edinburgh), we have been working on completing the proof of the crepanant resolution conjecture. The Donaldson-Thomas numbers of a variety are typically organized in a generating series, which a posteriori is the expansion of a rational function. What seems missing from [Cal16b] is a clever use of the Hall algebra, to ensure two different generating series are actually the expansions of the same rational function, after a change of variables. We have recently made a breakthrough, and are able to complete the proof in a special case.

# REFERENCES

- [AT09] Mathieu Anel and Bertrand Toën. Dénombrabilité des classes d'équivalences dérivées de variétés algébriques. *J. Algebraic Geom.*, 18(2):257–277, 2009.
- [BCY12] Jim Bryan, Charles Cadman, and Ben Young. The orbifold topological vertex. *Adv. Math.*, 229(1):531–595, 2012.
- [BKR01] Tom Bridgeland, Alastair King, and Miles Reid. The McKay correspondence as an equivalence of derived categories. *J. Amer. Math. Soc.*, 14(3):535–554, 2001.
- [BL16] L. Borisov and Z. Li. On Clifford double mirrors of toric complete intersections. *ArXiv e-prints*, January 2016.
- [Bri02] Tom Bridgeland. Flops and derived categories. *Invent. Math.*, 147(3):613–632, 2002.
- [Cala] John Calabrese. Chopping up derived categories. available online.
- [Calb] John Calabrese. Torelli theorems for semiorthogonal components. *in preparation*.
- [Cal16a] John Calabrese. Donaldson-Thomas invariants and flops. *J. Reine Angew. Math.*, 716:103–145, 2016.
- [Cal16b] John Calabrese. On the crepant resolution conjecture for Donaldson-Thomas invariants. *J. Algebraic Geom.*, 25(1):1–18, 2016.
- [Cal17a] John Calabrese. A note on derived equivalences and birational geometry. *Bull. Lond. Math. Soc.*, to appear, 2017.
- [Cal17b] John Calabrese. Relative singular twisted Bondal-Orlov. *Math. Res. Lett.*, to appear, 2017.
- [Cal17c] John Calabrese. A remark on generators of  $\mathcal{D}(X)$  and flags. *Manuscripta Math.*, 154(1-2):275–278, 2017.
- [CG15] John Calabrese and Michael Groechenig. Moduli problems in abelian categories and the reconstruction theorem. *Algebr. Geom.*, 2(1):1–18, 2015.
- [CP17] John Calabrese and Roberto Pirisi. Gabriel's theorem and birational geometry. *Draft available upon request*, 2017.
- [CT16] John R. Calabrese and Richard P. Thomas. Derived equivalent Calabi-Yau threefolds from cubic fourfolds. *Math. Ann.*, 365(1-2):155–172, 2016.
- [HR16] D. Huybrechts and J. Rennemo. Hochschild cohomology versus the Jacobian ring, and the Torelli theorem for cubic fourfolds. *ArXiv e-prints*, October 2016.
- [JS12] Dominic Joyce and Yinan Song. A theory of generalized Donaldson-Thomas invariants. *Mem. Amer. Math. Soc.*, 217(1020):iv+199, 2012.
- [Kaw02] Yujiro Kawamata.  $D$ -equivalence and  $K$ -equivalence. *J. Differential Geom.*, 61(1):147–171, 2002.
- [KS10] Maxim Kontsevich and Yan Soibelman. Motivic Donaldson-Thomas invariants: summary of results. In *Mirror symmetry and tropical geometry*, volume 527 of *Contemp. Math.*, pages 55–89. Amer. Math. Soc., Providence, RI, 2010.
- [Kuz07] Alexander Kuznetsov. Homological projective duality. *Publ. Math. Inst. Hautes Études Sci.*, (105):157–220, 2007.
- [Les15] John Lesieutre. Derived-equivalent rational threefolds. *Int. Math. Res. Not. IMRN*, (15):6011–6020, 2015.
- [Mau16] Daves Maulik. Stable pairs and the HOMFLY polynomial. *Invent. Math.*, 204(3):787–831, 2016.
- [MP14] Sven Meinhardt and Holger Partsch. Quotient categories, stability conditions, and birational geometry. *Geom. Dedicata*, 173:365–392, 2014.
- [Muk81] Shigeru Mukai. Duality between  $D(X)$  and  $D(\hat{X})$  with its application to Picard sheaves. *Nagoya Math. J.*, 81:153–175, 1981.
- [PVdB16] Dennis Presotto and Michel Van den Bergh. Noncommutative versions of some classical birational transformations. *J. Noncommut. Geom.*, 10(1):221–244, 2016.
- [Ser55] Jean-Pierre Serre. Faisceaux algébriques cohérents. *Ann. of Math. (2)*, 61:197–278, 1955.