

ECE551 - Homework 2

Khoi-Nguyen Mac

September 11, 2016

1 Frames and Bases

- (a) The synthesis operator associated with $\{\varphi_k\}_{k \in \mathcal{K}}$ in \mathbb{R}^2 is

$$\Phi\alpha = \sum_{k \in \mathcal{K}} \alpha_k \varphi_k$$

Hence,

$$\begin{aligned}\Phi_1 &= \begin{bmatrix} \frac{1}{2} & 0 \\ \frac{\sqrt{3}}{2} & 1 \end{bmatrix} \\ \Phi_2 &= \begin{bmatrix} \frac{1}{2\sqrt{2}} & -\frac{\sqrt{3}}{\sqrt{2}} & \frac{1}{\sqrt{2}} & 0 \\ \frac{\sqrt{3}}{\sqrt{2}} & \frac{1}{2\sqrt{2}} & 0 & \frac{1}{\sqrt{2}} \end{bmatrix} \\ \Phi_3 &= \begin{bmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix} \\ \Phi_4 &= \begin{bmatrix} 1 & \frac{1}{\sqrt{2}} & 0 \\ 0 & \frac{1}{\sqrt{2}} & 1 \end{bmatrix}\end{aligned}$$

(Note that we reuse the notation Φ to represent the matrix representation.)

- (b) For $\Phi \in \mathbb{R}^{M \times N}$, if $M = N$ then it is a basis, if $M > N$ then it is a frame.

Let A be the inverse of the Gram matrix of basis Φ , i.e. $A = (\Phi^* \Phi)^{-1}$. Then $\tilde{\Phi} = \Phi A = \Phi (\Phi^* \Phi)^{-1}$ forms a dual basis with Φ .

Let $B = (\Phi \Phi^*)^{-1}$, where Φ is a frame. Then $\tilde{\Phi} = B \Phi = (\Phi \Phi^*)^{-1} \Phi$ forms the canonical dual frame associated with frame Φ .

For Φ_1 (basis),

$$A_1 = \begin{bmatrix} 1 & \frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & 1 \end{bmatrix} \Rightarrow A_1^{-1} = \begin{bmatrix} 4 & -2\sqrt{3} \\ -2\sqrt{3} & 4 \end{bmatrix}$$

$$\tilde{\Phi}_1 = \begin{bmatrix} 2 & -\sqrt{3} \\ 0 & 1 \end{bmatrix}$$

For Φ_2 (frame),

$$B_2 = \frac{1}{2} \begin{bmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} & 1 & 0 \\ \frac{\sqrt{3}}{2} & \frac{1}{2} & 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \\ -\frac{\sqrt{3}}{2} & \frac{1}{2} \\ 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = B_2^{-1}$$

$$\tilde{\Phi}_2 = \frac{1}{\sqrt{2}} \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} & 1 & 0 \\ \frac{\sqrt{3}}{2} & \frac{1}{2} & 0 & 1 \end{bmatrix} = \frac{1}{\sqrt{2}} \begin{bmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} & 1 & 0 \\ \frac{\sqrt{3}}{2} & \frac{1}{2} & 0 & 1 \end{bmatrix} = \Phi_2$$

For Φ_3 (basis),

$$A_3 = \frac{1}{4} \begin{bmatrix} 1 & \sqrt{3} \\ -\sqrt{3} & 1 \end{bmatrix} \begin{bmatrix} 1 & -\sqrt{3} \\ \sqrt{3} & 1 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = A_3^{-1}$$

$$\Rightarrow \tilde{\Phi}_3 = \Phi_3$$

For Φ_4 (basis),


$$B_4 = \begin{bmatrix} \frac{3}{2} & \frac{1}{2} \\ \frac{1}{2} & \frac{3}{2} \end{bmatrix} \Rightarrow B_4^{-1} = \begin{bmatrix} \frac{3}{4} & -\frac{1}{4} \\ -\frac{1}{4} & \frac{3}{4} \end{bmatrix}$$

$$\tilde{\Phi}_4 = \begin{bmatrix} \frac{3}{4} & \frac{1}{2\sqrt{2}} & -\frac{1}{4} \\ -\frac{1}{4} & \frac{1}{2\sqrt{2}} & \frac{3}{4} \end{bmatrix}$$


Figure 1 shows the sketch of the sets and their duals.

- (c) $\langle \varphi_{1,0}, \varphi_{1,1} \rangle = \frac{\sqrt{3}}{2}$, so the basis Φ_1 is not orthogonal, thus not orthonormal. $B_2 = I$, so the frame Φ_2 is tight (a frame is tight if $\Phi\Phi^* = I$.) $\langle \varphi_{3,0}, \varphi_{3,1} \rangle = 0$ and $\|\varphi_{3,0}\| = \|\varphi_{3,1}\| = 1$, so the basis Φ_3 is orthonormal. $B_4 \neq I$, so the frame Φ_4 is not tight.


- (d) $x = \begin{bmatrix} 2 \\ 0 \end{bmatrix}$ and $\alpha_{i,k} = \langle x, \tilde{\varphi}_{i,k} \rangle$. Therefore,




(a) Φ_1



(b) Φ_2



(c) Φ_3



(d) Φ_4

Figure 1: Original sets and their duals.

For Φ_1 ,

$$\alpha_{1,0} = \left\langle \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 2 \\ 0 \end{bmatrix} \right\rangle = 4 \quad \alpha_{1,1} = \left\langle \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} -\sqrt{3} \\ 1 \end{bmatrix} \right\rangle = -2\sqrt{3}$$

For Φ_2 ,

$$\begin{aligned} \alpha_{2,0} &= \frac{1}{\sqrt{2}} \left\langle \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} \frac{1}{2} \\ \frac{\sqrt{3}}{2} \end{bmatrix} \right\rangle = \frac{1}{\sqrt{2}} & \alpha_{2,1} &= \frac{1}{\sqrt{2}} \left\langle \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} -\frac{\sqrt{3}}{2} \\ \frac{1}{2} \end{bmatrix} \right\rangle = -\sqrt{\frac{3}{2}} \\ \alpha_{2,2} &= \frac{1}{\sqrt{2}} \left\langle \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right\rangle = \frac{2}{\sqrt{2}} & \alpha_{2,3} &= \frac{1}{\sqrt{2}} \left\langle \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \end{bmatrix} \right\rangle = 0 \end{aligned}$$

For Φ_3 ,

$$\alpha_{3,0} = \frac{1}{2} \left\langle \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ \sqrt{3} \end{bmatrix} \right\rangle = 1 \quad \alpha_{3,1} = \frac{1}{2} \left\langle \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} -\sqrt{3} \\ 1 \end{bmatrix} \right\rangle = -\sqrt{3}$$

For Φ_4 ,

$$\begin{aligned} \alpha_{4,0} &= \left\langle \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} \frac{3}{4} \\ -\frac{1}{4} \end{bmatrix} \right\rangle = \frac{3}{2} & \alpha_{4,1} &= \frac{1}{2\sqrt{2}} \left\langle \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \end{bmatrix} \right\rangle = \frac{1}{\sqrt{2}} \\ \alpha_{4,2} &= \left\langle \begin{bmatrix} 2 \\ 0 \end{bmatrix}, \begin{bmatrix} -\frac{1}{4} \\ \frac{3}{4} \end{bmatrix} \right\rangle = -\frac{1}{2} \end{aligned}$$

(e) We check the values for $\alpha_{i,k}$ by verifying the expansion $x = \sum_k \alpha_{i,k} \varphi_{i,k}$.

For Φ_1 ,

$$\sum_k \alpha_{1,k} \varphi_{1,k} = 4 \begin{bmatrix} \frac{1}{2} \\ \frac{\sqrt{3}}{2} \end{bmatrix} - 2\sqrt{3} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \end{bmatrix} = x$$

For Φ_2 ,

$$\sum_k \alpha_{2,k} \varphi_{2,k} = \frac{1}{2} \left(1 \begin{bmatrix} \frac{1}{2} \\ \frac{\sqrt{3}}{2} \end{bmatrix} - \sqrt{3} \begin{bmatrix} -\frac{\sqrt{3}}{2} \\ \frac{1}{2} \end{bmatrix} + 2 \begin{bmatrix} 1 \\ 0 \end{bmatrix} \right) = \begin{bmatrix} 2 \\ 0 \end{bmatrix} = x$$

For Φ_3 ,

$$\sum_k \alpha_{3,k} \varphi_{3,k} = \begin{bmatrix} \frac{1}{2} \\ \frac{\sqrt{3}}{2} \end{bmatrix} - \sqrt{3} \begin{bmatrix} -\frac{\sqrt{3}}{2} \\ \frac{1}{2} \end{bmatrix} = \begin{bmatrix} 2 \\ 0 \end{bmatrix} = x$$

For Φ_4 ,

$$\sum_k \alpha_{4,k} \varphi_{4,k} = \frac{3}{2} \begin{bmatrix} 1 \\ 0 \end{bmatrix} + \frac{1}{\sqrt{2}} \begin{bmatrix} \frac{1}{\sqrt{2}} \\ \frac{1}{\sqrt{2}} \end{bmatrix} - \frac{1}{2} \begin{bmatrix} 0 \\ 1 \end{bmatrix} = x$$

(f) We verify that $\Phi\tilde{\Phi}^\top = I$.

$$\Phi_1\tilde{\Phi}_1^\top = \begin{bmatrix} \frac{1}{2} & 0 \\ \frac{\sqrt{3}}{2} & 1 \end{bmatrix} \begin{bmatrix} 2 & 0 \\ -\sqrt{3} & 1 \end{bmatrix} = I$$

$$\Phi_2\tilde{\Phi}_2^\top = \Phi_2\Phi_2^\top = B_2 = I$$

$$\Phi_3\tilde{\Phi}_3^\top = \Phi_3\Phi_3^\top = \begin{bmatrix} \frac{1}{2} & -\frac{\sqrt{3}}{2} \\ \frac{\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix} \begin{bmatrix} \frac{1}{2} & \frac{\sqrt{3}}{2} \\ -\frac{\sqrt{3}}{2} & \frac{1}{2} \end{bmatrix} = I$$

$$\Phi_4\tilde{\Phi}_4^\top = \begin{bmatrix} 1 & \frac{1}{\sqrt{2}} & 0 \\ 0 & \frac{1}{\sqrt{2}} & 1 \end{bmatrix} \begin{bmatrix} \frac{3}{4} & -\frac{1}{4} \\ \frac{1}{2\sqrt{2}} & \frac{1}{2\sqrt{2}} \\ -\frac{1}{4} & \frac{3}{4} \end{bmatrix} = I$$

(g) We check if $\|x\|^2 = \sum_k |\alpha_{i,k}|^2$.

$$\|x\|^2 = \left\| \begin{bmatrix} 2 \\ 0 \end{bmatrix} \right\|^2 = 4^2 = 16$$

$$\sum_k |\alpha_{1,k}|^2 = 28$$

$$\sum_k |\alpha_{2,k}|^2 = \frac{13}{4}$$

$$\sum_k |\alpha_{3,k}|^2 = 4$$

$$\sum_k |\alpha_{4,k}|^2 = 3$$

Hence, the expansion does not preserve the norm.

(h) Since all of the expansion $\sum_k \alpha_{i,k} \varphi_{i,k} = x$, the expansion is redundant.

2 Linear Least-Squares Approximation

3 Orthogonalization of a Projection

(a) If $P = P^*$,

$$\langle Px, y \rangle = y^*(Px) = (y^*P)x = (y^*P^*)x = (Py)^*x = \langle x, Py \rangle.$$

Hence, P is self-adjoint with respect to $\langle \cdot, \cdot \rangle$ on \mathbb{C}^N if $P = P^*$.

(b)

(c)

4 Approximation by Orthogonal Indicator Tiles