

ECE551 - Homework 1

Khoi-Nguyen Mac

August 29, 2016

1 Geometry of orthogonal transformations in Euclidean spaces

1.1

$U \in \mathbb{R}^{N \times N}$ is an orthogonal matrix, therefore:

$$U^\top U = UU^\top = I.$$

By definition,

$$\begin{aligned}\|x\|^2 &:= x^\top x = \langle x, x \rangle, \quad x \in \mathbb{R}^N \\ \Rightarrow \|Ux\| &= \langle Ux, Ux \rangle = (Ux)^\top Ux = x^\top U^\top Ux = x^\top x = \|x\|^2 \\ \Rightarrow \|Ux\| &= \|x\|, \quad \because \text{norms are non-negative.}\end{aligned}$$

1.2

By definition,

$$\begin{aligned}\langle x, y \rangle &:= y^\top x = \sum_{i=0}^{N-1} x_i y_i, \quad x, y \in \mathbb{R}^N \\ \Rightarrow \langle Ux, Uy \rangle &= (Uy)^\top Ux = y^\top U^\top Ux = y^\top x = \langle x, y \rangle.\end{aligned}$$

1.3

If $M > N$ and $U^\top U = I$, then $\text{rank}(U) = N \Rightarrow U^\top U = I_N$. Since $x \in \mathbb{R}^N$ and $y \in \mathbb{R}^N$,

$$\begin{aligned}x^\top U^\top Ux &= x^\top I_N x = x^\top x \\ y^\top U^\top Ux &= y^\top I_N x = y^\top x.\end{aligned}$$

Hence, 1.1 and 1.2 hold.

1.4

If $M < N \Rightarrow \text{rank}(U)$ is at most M .

$$U^\top U \neq I_N.$$

Hence, 1.1 and 1.2 do not hold.

2 Some basic properties of inner product spaces

Notice that:

$$\langle a, b + c \rangle = \overline{\langle b + c, a \rangle} = \overline{\langle b, a \rangle} + \overline{\langle c, a \rangle} = \langle a, b \rangle + \langle a, c \rangle.$$

2.1 The Cauchy-Schwarz inequality

$$|\langle x, y \rangle| \leq \|x\| \|y\|$$

Proof: If $\langle x, y \rangle = 0$, then theorem holds because $\|x\|$ and $\|y\| \geq 0$.

Suppose that $x \neq 0$ and $y \neq 0$. Let $z \in \mathbb{C}$, such that:

$$z = \frac{\langle x, y \rangle}{\|y\|^2} = \frac{\overline{\langle y, x \rangle}}{\|y\|^2}.$$

We have

$$\begin{aligned} 0 &\leq \|x - zy\|^2 = \langle x - zy, x - zy \rangle = \langle x, x - zy \rangle - \langle zy, x - zy \rangle \\ &= \langle x, x \rangle - \langle x, zy \rangle - \langle zy, x \rangle + \langle zy, zy \rangle \\ &= \|x\|^2 - \bar{z}\langle x, y \rangle - z\langle y, x \rangle + z\bar{z}\|y\|^2 \\ &= \|x\|^2 - \frac{\langle y, x \rangle \langle x, y \rangle}{\|y\|^2} - \frac{\langle x, y \rangle \langle y, x \rangle}{\|y\|^2} + \frac{\langle x, y \rangle \langle y, x \rangle}{\|y\|^2} \|y\|^2 \\ &= \|x\|^2 - \frac{\langle x, y \rangle \langle y, x \rangle}{\|y\|^2} \\ &= \|x\|^2 - \frac{|\langle x, y \rangle|^2}{\|y\|^2}. \end{aligned}$$

Therefore,

$$\begin{aligned} 0 &\leq \|x\|^2 - \frac{|\langle x, y \rangle|^2}{\|y\|^2} \\ \Leftrightarrow \|x\|^2 &\geq \frac{|\langle x, y \rangle|^2}{\|y\|^2} \\ \Leftrightarrow \|x\|^2 \|y\|^2 &\geq |\langle x, y \rangle|^2 \\ \Leftrightarrow \|x\| \|y\| &\geq |\langle x, y \rangle|. \end{aligned}$$

□

The equality occurs iff $x = \alpha y$, for some scalar α .

Proof: By substituting $x = \alpha y$, we have

$$|\langle x, y \rangle| = |\langle \alpha y, y \rangle| = |\alpha| |\langle y, y \rangle| = |\alpha| \|y\|^2 = |\alpha| \|y\| \|y\| = \|x\| \|y\|.$$

□

2.2 The triangle inequality

$\|x + y\|^2 \leq \|x\|^2 + \|y\|^2$ with equality iff $y = \alpha x$

Proof:

$$\begin{aligned} \|x + y\|^2 &= \langle x + y, x + y \rangle \\ &= \langle x, x + y \rangle + \langle y, x + y \rangle \\ &= \langle x, x \rangle + \langle x, y \rangle + \langle y, x \rangle + \langle y, y \rangle \\ &= \|x\|^2 + \|y\|^2 + \langle x, y \rangle + \langle y, x \rangle \end{aligned}$$

By Cauchy-Schwarz inequality:

$$\begin{aligned} \langle x, y \rangle &\leq \|x\| \|y\| \\ \Rightarrow \langle y, x \rangle &\leq \|x\| \|y\| \\ \Rightarrow \langle x, y \rangle + \langle y, x \rangle &\leq 2\|x\| \|y\|. \end{aligned}$$

Therefore,

$$\begin{aligned} \|x + y\|^2 &\leq \|x\|^2 + \|y\|^2 + 2\|x\| \|y\| = (\|x\| + \|y\|)^2 \\ \Rightarrow \|x + y\| &\leq \|x\| + \|y\| \quad \because \text{norms are non-negative.} \end{aligned}$$

If $y = \alpha x$ then the equality of Cauchy-Schwarz theorem occurs. Therefore,

$$\begin{aligned} \langle x, y \rangle + \langle y, x \rangle &= 2\|x\| \|y\| \\ \Rightarrow \|x + y\|^2 &= \|x\|^2 + \|y\|^2 + 2\|x\| \|y\| = (\|x\| + \|y\|)^2 \\ \Rightarrow \|x + y\| &= \|x\| + \|y\|. \end{aligned}$$

□

2.3 Parallelogram law

$$\|x + y\|^2 + \|x - y\|^2 = 2(\|x\|^2 + \|y\|^2)$$

Proof:

$$\begin{aligned}
LHS &= \langle x + y, x + y \rangle + \langle x - y, x - y \rangle \\
&= (\|x\|^2 + \|y\|^2 + \langle x, y \rangle + \langle y, x \rangle) + (\|x\|^2 + \|y\|^2 - \langle x, y \rangle - \langle y, x \rangle) \\
&= 2(\|x\|^2 + \|y\|^2) = RHS.
\end{aligned}$$

□

3 Least-squares approximation with orthonormal bases

3.1

We notice that $\langle x, \varphi_i \rangle \varphi_i$ is the orthogonal projection of x onto φ_i . Since $\varphi_i \in \hat{\mathcal{B}}$ are orthonormal, $\sum_{\varphi_i \in \hat{\mathcal{B}}} \langle x, \varphi_i \rangle \varphi_i$ is the orthogonal projection of x onto the subspace $\text{span}(\hat{\mathcal{B}}) = \hat{V}$.

We need to prove that the Euclidean distance between x and its orthogonal projection on \hat{V} is the shortest among Euclidean distances between x and other vectors in \hat{V} .

Let $z = x - \hat{x}$, where $\hat{x} = \sum_{\varphi_i \in \hat{\mathcal{B}}} \langle x, \varphi_i \rangle \varphi_i$. Thus, $z \perp \hat{V}$.

Since any vectors in \hat{V} can be written as $f(\alpha) = \sum_{\varphi_i \in \hat{\mathcal{B}}} \alpha_i \varphi_i$ (α is a vector and α_i 's are scalars,) we have:

$$\begin{aligned}
\|x - f(\alpha)\|^2 &= \|x - \hat{x} + \hat{x} - f(\alpha)\|^2 = \|z + \hat{x} - f(\alpha)\|^2 \\
&= \langle z + \hat{x} - f(\alpha), z + \hat{x} - f(\alpha) \rangle \\
&= \|z\|^2 + \|\hat{x} - f(\alpha)\|^2 + \langle z, \hat{x} - f(\alpha) \rangle + \langle \hat{x} - f(\alpha), z \rangle
\end{aligned}$$

We can see that $\hat{x} \in \hat{V}$ and $f(\alpha) \in \hat{V}$, therefore $\hat{x} - f(\alpha) \in \hat{V}$. Since $z \perp \hat{V} \Rightarrow z \perp \hat{x} - f(\alpha) \Rightarrow \langle z, \hat{x} - f(\alpha) \rangle = \langle \hat{x} - f(\alpha), z \rangle = 0$.

Therefore, $\|x - f(\alpha)\|^2 = \|z\|^2 + \|\hat{x} - f(\alpha)\|^2$. Since $\|\hat{x} - f(\alpha)\|^2 \geq 0$,

$$\begin{aligned}
\|x - f(\alpha)\|^2 &\geq \|z\|^2 \\
\Leftrightarrow \|x - f(\alpha)\|^2 &\geq \|x - \hat{x}\|^2 \\
\Leftrightarrow \|x - f(\alpha)\| &\geq \|x - \hat{x}\|
\end{aligned}$$

3.2

We know that any inner products can define the valid norm

$$\|x\| = \sqrt{\langle x, x \rangle}.$$

Therefore $\|x - f(\alpha)\|^2 = \langle z + \hat{x} - f(\alpha), z + \hat{x} - f(\alpha) \rangle$, with other inner products. The expansion

$$\langle z + \hat{x} - f(\alpha), z + \hat{x} - f(\alpha) \rangle = \|z\|^2 + \|\hat{x} - f(\alpha)\|^2 + \langle z, \hat{x} - f(\alpha) \rangle + \langle \hat{x} - f(\alpha), z \rangle$$

also uses the properties of general inner products. Hence, the proof holds for other kinds of inner products, instead of only standard Euclidean one.

4 Signal Sets and Spaces