

Abstract

1 Introduction

The characterization of tree canopy structure in wooded ecosystems constitutes a long-standing field of research that has been fundamental to interpreting, modelling, and improving understanding of ecosystem function (Watt, 1947; Whittaker and Woodwell, 1969; Horn, 1971; Maarel, 1996). Canopy structure describes the spatial distribution and density of canopy foliage, comprising the primary interface between trees, the atmosphere and sunlight. It is therefore essential to understand the drivers of variation in canopy structure to improve modelling of earth-atmosphere carbon fluxes and community assembly ().

At continental scales, variation in canopy height and canopy cover, two coarse measures of canopy structure both of which have been shown to affect woody productivity and correlate with woody biomass (), can largely be explained by climate and edaphic data (**SOME-GEDI**). Increased resource availability allows for larger trees and more closed canopies (). At the scale of a single tree community however, where variation in climate and soil may be negligible, variation in canopy structure is thought to be affected principally by an interacting combination of tree canopy species composition (), and disturbance history (). However, empirical testing of these mechanisms thought to drive canopy structure in natural wooded ecosystems remains sparse across many biomes ().

Following established biodiversity-ecosystem function theory, the niche partitioning of canopy space, i.e. the spatial complementarity of individual tree canopies, hereafter referred to as ‘crown complementarity’, is thought to be a key mechanism underlying positive biodiversity-productivity effects in wooded ecosystems (Pretzsch, 2014; Barry et al., 2019). Biodiversity-ecosystem function theory predicts that crown complementarity and thus canopy complexity and foliage density should increase with tree diversity in the local neighbourhood, thus increasing standing biomass and woody productivity, as coexisting species must occupy non-identical niche space to avoid competitive exclusion (**Gadow1993**).

As well as the species diversity of trees in a local neighbourhood, the spatial distribution and relative size dominance of those trees, i.e. stand structure, is also expected to affect crown complementarity and canopy structure. Increased heterogeneity in tree size, whether a result of species diversity or disturbance history, is expected to increase crown complementarity as individuals of different sizes can occupy different layers of the canopy (). Additionally, clustering of individuals in space is expected to increase canopy structural heterogeneity across a stand, but ultimately de-

crease total foliage density due to an increase in competitive interactions (). Clustering may occur as a result of disturbance history, or as a result of strong facilitation effects among individuals in a hostile environment (**Ratcliffe2017**).

While much work in the field of forest management has been done to test biotic drivers of tree canopy structure in temperate () and boreal forests (), similar work in the tropics is comparatively scarce (). In dry tropical woodlands especially, tree canopy structure and its effect on ecosystem function has received little attention, possibly due to the misplaced assumption that woody productivity in these ecosystems does not represent a globally significant carbon flux (), or that tree canopies in these smaller stature woodlands do not interact and compete for resources to the same degree as in large stature forests (). In recent years however, it has been shown that dry tropical woodlands represent the largest uncertainty in our estimates of the terrestrial carbon cycle (Quéré et al., 2018; Ahlstrom et al., 2015). Sitch et al. (2015) demonstrated the dominant role of the dry tropics in driving variability in the terrestrial carbon sink, and showed that the dry tropics are the fastest increasing component of the terrestrial carbon sink. Part of this uncertainty arises from our lacking a nuanced understanding of how species composition and structure affect ecosystem function in these ecosystems, which underpins the Dynamic Global Vegetation Models (DGVMs) fed into global carbon dynamics models. This knowledge gap prompts further research of the biotic drivers of variation in productivity in the dry tropics, of which canopy structure is a key part ().

Canopy structure is multi-dimensional and has previously been explained using a plethora of simple metrics that originated in forest and community ecology (). Assessments of canopy structure in the dry tropical have most often modelled tree canopies as a series of ellipses (2D) or ellipsoids (3D) based on field measurements with measuring tapes (). Measurements of this kind are time consuming and yet are an over-simplification of canopy structure (). Alternatively, canopy cover is often measured using indirect optical methods which partition sky from canopy material, i.e. with hemispherical photography or the commonly used LAI-2000, providing a 2D representation of the canopy but lacking information on vertical canopy structure. In recent years, particularly in temperate and boreal forests, LiDAR (Light Detection And Ranging) has emerged as a suitable technology for rapidly and precisely assessing canopy structure in 3D, conserving information on 3D structure of the calibre that is required to understand it's complexities ().

In this study we applied terrestrial LiDAR techniques to woodland-savanna mosaics at two sites in southern Africa, with the aim of increasing our understanding of how various metrics of tree canopy structure are affected by tree neighbourhood diversity and stand structure. Our overarching contention is that neighbourhoods of greater tree diversity and greater structural diversity allow greater canopy complexity and foliage density, resulting in higher productivity, and ultimately

67 a more ‘forest-like’ community, rather than an open canopy savanna.

68 **2 Materials and methods**

69 **2.1 Study sites**

70 Measurements were conducted at two sites, the first in Bicuar National Park, southwest Angola
71 ($S15.1^{\circ}$, $E14.8^{\circ}$), and the second in and around Mtarure Forest Reserve, southeast Tanzania ($S9.0^{\circ}$,
72 $E39.0^{\circ}$) (Figure 1). At each site, 1 ha plots were sited in areas of miombo woodland vegetation,
73 across a gradient of stem density. In Angola, 15 plots were sampled, while in Tanzania, seven were
74 sampled following the curtailment of fieldwork due to COVID-19 travel restrictions. Fieldwork was
75 conducted between February and April at both sites, during the peak growth period of each site in
76 order to capture the highest foliage volume in the canopy.




Figure 1: Location of study sites within southern Africa (a), and of 1 ha plots within each site. The blue polygons denote the boundaries of protected areas which encompass the majority of study sites, Bicular National Park in Angola (b), and Mtarure Forest Reserve in Tanzania (c). The background of each site map is a re-classified version of the GlobCover global land cover classification (**Globcover**).

2.2 Field measurements

Each plot was further subdivided into nine 10 m diameter circular subplots arranged in a regular grid, with a 15 m buffer from the plot edge and 35 m between subplots. For each subplot, we measured all woody stems >5 cm trunk diameter with canopy material inside the subplot. We identified each stem to species and measured trunk diameter (diameter at breast height - 1.3 m), height to top of canopy material, canopy area calculated as an ellipse of two perpendicular crown diameter measurements, distance and direction of stem from the subplot centre.

2.3 Terrestrial laser scanning

Within each subplot, a variable number of scans were recorded using a Leica HDS6100 phase-shift terrestrial laser scanner (TLS) (**Leica**). The number and position of scans within a subplot was determined by the arrangement of canopy material in the subplot. Scan positions were arranged to minimise shadows within the canopy, and to maximise canopy penetration. The number of scans per subplot ranged between one and five in both Angola and Tanzania. Registration of multiple scans from different locations around each subplot minimised the occlusion effect and improved canopy penetration.

2.4 Data analysis

2.4.1 Scan processing

Point clouds from scans in each subplot were registered and unified using Leica Cyclone (version 9.1). Targets from each scan were aligned using Cyclone's automatic target acquisition.

Point clouds were voxelised to cubic voxel sizes of different sizes depending on the application of the data. For subplot height profile estimation and gap fraction we used 5 cm³ voxels, and for whole plot canopy rugosity we used 10 cm³ voxels. Variation in voxel size reflects the spatial scale of each analysis, and is bounded by the beam divergence of the scanner (). Choosing voxels that are too small can result in pock-marked representations of surfaces that are especially problematic when estimating canopy structure at a larger scale, such as when estimating canopy top roughness, while voxels that are too large can result in an over-estimation of plant volume when estimating canopy foliage density (Cifuentes et al., 2014). Voxels were classed as filled if they intersected with one or more points.

Partial object interceptions caused by phase-shift laser scanners can produce erroneous results and must be corrected for to accurately estimate canopy height (). We used a noise reduction algorithm from to discard points that appeared far from other points. This effectively removed ghost points produced by partial interceptions and also removed many erroneous returns caused by airborne dust particles, which was common in our study site.

Ground points were classified using the Progressive Morphological Filter (PMF) from Zhang et al. (2003). Point cloud height was reclassified height based on this revised ground layer by measuring the vertical distance between the nearest ground point and each point.

Raw points clouds for each subplot had $\sim 2.9 \times 10^8$ points, $\sim 4.5 \times 10^7$ points after voxelisation, and $\sim 2.1 \times 10^7$ points after noise reduction.

We used ray-tracing to calculate gap fraction from TLS scans at the centre of each subplot. Hemi-

spherical images were created using the POV-ray software (). Voxels were converted to matt black cubes filling the voxel volume, with a white sky box and no light source. A ‘camera’ with a 180° fisheye lens was placed at the subplot centre at a height of 1.8 m pointing directly upwards. The images produced by POV-ray were analysed using Hemiphot (**Steege**) to estimate canopy gap fraction.

2.5 Stand structure

For each subplot, we calculated an adapted version of the Hegyi index to estimate crowding (**Hegyi1974**). At the plot level, we estimated the regularity of species distribution using the spatial mingling index (**Gadow**). We also measured whole plot stand structure using the Winkelmass ().

2.6 Statistical analysis

Linear mixed effects models tested the effects of diversity and stand structural metrics on canopy structure.

128 3 Results

129 3.1 Vertical canopy complexity



Figure 2: Bivariate relationships between subplot canopy structure metrics (y axis) and diversity and stand structure metrics (x axis). Points and linear model lines of best fit are coloured by site. The black line of best fit is a linear model including both sites.

130 Species diversity, measured by species richness of the tree neighbourhood around each 10 m diam-
 131 eter subplot, appears to have weak but positive effects on canopy layer diversity and total canopy
 132 cover, with possibly weak negative effects on the height of peak canopy density. Stand structural
 133 metrics have much stronger positive effects on canopy structure, particularly canopy cover and

134 layer diversity. (Figure 2).

135 Linear mixed effects models showed that species richness of the subplot neighbourhood had vari-
 136 able effects across the measures of canopy structure, but the effect sizes were not significant for
 137 any model (Figure 3). On the other hand, stand structural metrics, the Hegyi index, Coefficient of
 138 Variation of stem diameter and total stem basal area, had a much greater effect on canopy struc-
 139 ture variables. The Hegyi index had a positive significant effect on canopy cover, while basal area
 140 had positive significant effects on total foliage density, layer diversity, and canopy height, with
 141 strong non-significant effects on non-uniformity of vertical foliage distribution and the height of
 142 peak foliage density.




Figure 3: Standardized fixed effect slopes for each model of a canopy structure metric. Slope estimates are ± 1 standard error. Slope estimates where the interval (standard error) does not overlap zero are considered to be significant effects.

143 3.2 Canopy rugosity

144 Similar to subplot analyses, at the whole-plot level, tree species diversity tended to have weak pos-
 145 itive effects on canopy complexity metrics, while stand structural diversity metrics had stronger
 146 positive effects (Figure 5). Strong positive relationships of basal area on canopy complexity are
 147 driven mostly by two plots with particularly low basal area in Mtarure. These plots are sparse
 148 thorny savanna, dominated by *Senegalia* spp. (Figure 4). Indeed, linear models using only Bicuar
 149 plots show divergent relationships. These two plots also have particularly low canopy cover, canopy
 150 height, and canopy top roughness, despite having similar tree species richness and spatial distribu-
 151 tion of trees (Winkelmass) as other plots.




Figure 4: The first two axes of a Non-metric Multi-Dimensional Scaling (NMDS) analysis of tree species diversity in each plot. Species scores are labelled as black text, while plot scores are labelled as coloured points. Plots can be split into three principal groups: 1) ABG_9, ABG_13 and ABG_15, dominated by *Baikiaea plurijuga*; 2) the other Bicar plots plus TKW_9, TKW_13, TKW_19 and TKW_21, dominated by *Julbernardia* spp., *Brachystegia* spp. and *Ochna* spp.; 3) TKW_7, TKW_10 and TKW_25, dominated by *Senegalia* spp. and *Vachellia* spp..




Figure 5: Bivariate relationships between whole-plot canopy structure metrics (y axis) and diversity and stand structure metrics (x axis). Points and linear model lines of best fit are coloured by site. The black line of best fit is a linear model including both sites.

152 Maximum canopy height at the plot-level appears to be positively affected by both tree species di-
 153 versity and total stem basal area. Basal area has a positive effect on canopy top roughness. Varia-
 154 tion in canopy height was increased both by variation in stem diameter and total stem basal area.
 155 The only significant fixed effect on canopy cover at the plot level was the Winkelmass, which mea-
 156 sures the spatial clustering of trees.




Figure 6: Standardized fixed effect slopes for whole-plot canopy rugosity. Slope estimates are ± 1 standard error. Slope estimates where the interval (standard error) does not overlap zero are considered to be significant effects.

3.3 Comparing subplot and plot measures of canopy structure

Plot-level and subplot-level canopy structure metrics were highly correlated in many cases (Figure 6). Notably, as canopy top rugosity increases, various subplot canopy complexity and density metrics decrease. Additionally, as canopy top roughness increases, many subplot canopy complexity and density metrics increase.




Figure 7: Bivariate plots of canopy structural metrics at the subplot and plot-level. Each point represents the mean values of a single plot. Points and linear model fits are coloured according to site. The black linear model combines both sites. Error bars on points are the standard deviation of subplot metrics.

4 Discussion

We investigated the effects of tree species diversity and structural diversity on several metrics of canopy structure that are hypothesised to affect plot productivity.

Species diversity didn't have strong effects on canopy structure, but stand structure did.

With the whole plot effect of tree species diversity on canopy height, what we might be seeing is actually an increased canopy height allowing greater species diversity, rather than the other way round.

4.1 Scaling up from subplots

5 Conclusion

References

- Ahlstrom, A. et al. (2015). "The dominant role of semi-arid ecosystems in the trend and variability of the land CO₂ sink". In: *Science* 348.6237, pp. 895–899. DOI: 10.1126/science.aaa1668.
- Barry, Kathryn E. et al. (2019). "The Future of Complementarity: Disentangling Causes from Consequences". In: *Trends in Ecology & Evolution* 34.2, pp. 167–180. DOI: 10.1016/j.tree.2018.10.013.
- Cifuentes, Renato et al. (2014). "Effects of voxel size and sampling setup on the estimation of forest canopy gap fraction from terrestrial laser scanning data". In: *Agricultural and Forest Meteorology* 194, pp. 230–240. DOI: 10.1016/j.agrformet.2014.04.013.
- Horn, H. S. (1971). *The adaptive geometry of trees*. Princeton NJ, USA: Princeton University Press. ISBN: 0691080895.
- Maarel, Eddy van der (1996). "Pattern and process in the plant community: Fifty years after A.S. Watt". In: *Journal of Vegetation Science* 7.1, pp. 19–28. DOI: 10.2307/3236412.
- Pretzsch, H. (2014). "Canopy space filling and tree crown morphology in mixed-species stands compared with monocultures". In: *Forest Ecology and Management* 327, pp. 251–264. DOI: <http://dx.doi.org/10.1016/j.foreco.2014.04.027>.
- Quére, Corinne Le et al. (2018). "Global Carbon Budget 2018". In: *Earth System Science Data* 10.4, pp. 2141–2194. DOI: 10.5194/essd-10-2141-2018.
- Sitch, S. et al. (2015). "Recent trends and drivers of regional sources and sinks of carbon dioxide". In: *Biogeosciences* 12.3, pp. 653–679. DOI: 10.5194/bg-12-653-2015.
- Watt, Alex S. (1947). "Pattern and Process in the Plant Community". In: *The Journal of Ecology* 35.1/2, p. 1. DOI: 10.2307/2256497.

- 193 Whittaker, R. H. and G. M. Woodwell (1969). “Structure, Production and Diversity of the Oak-
194 Pine Forest at Brookhaven, New York”. In: *The Journal of Ecology* 57.1, p. 155. DOI: 10.2307/
195 2258214.
- 196 Zhang, Keqi et al. (2003). “A progressive morphological filter for removing nonground measure-
197 ments from airborne LIDAR data”. In: *IEEE Transactions on Geoscience and Remote Sensing*
198 41.4, pp. 872–882. DOI: 10.1109/tgrs.2003.810682.