

In this note we explore different attempts at expressing the  $B_2 \wedge B_2$  component of  $R_8^{(2)}$  with cluster polylogarithms.

First let us define “good”  $\mathcal{X}$ -coordinates: an  $\mathcal{X}$ -coordinate  $R$  is good if  $R$  and  $1 + R$  can be expressed as ratios of  $\mathcal{A}$ -coordinates of the form

$$\begin{aligned} \langle i \ i+1 \ jk \rangle, \quad \langle i(i-1 \ i+1)(j \ j+1)(k \ k+1) \rangle, \\ \langle i \ i+1 \ \bar{j} \cap \bar{k} \rangle, \quad \langle i(i-2 \ i-1)(i+1 \ i+2)(j \ j+1) \rangle. \end{aligned} \tag{1}$$

An algebra is good if it contains only good  $\mathcal{X}$ -coordinates.

### Fitting with $A_3$ ’s

There are 1600 good  $A_3$ ’s in  $\text{Gr}(4, 8)$ , with only 513 actual degrees of freedom in  $B_2 \wedge B_2$ . An ansatz of these functions can be fit to  $R_8^{(2)}$ . A zero-thought choice of free parameters leaves us with an expression involving 223  $A_3$ ’s.

### Fitting with $A_4$ ’s

There are 496 good  $A_4$ ’s.

### Fitting with $D_4$ ’s

There are 24 good  $D_4$ ’s.

### Fitting with $A_5$ ’s

There are 56 good  $A_5$ ’s. No choice of  $f_{A_5}$  can fit  $R_8^{(2)}$ .

### Fitting with $D_5$ ’s

There are no good  $D_5$ ’s. Furthermore, no  $D_5$  contains enough good  $A_3$ ’s to allow us to tune the free parameters in  $f_{D_5}$  in order to have only good  $A_3$ ’s appear in  $f_{D_5}$ . In other words, any representation of  $R_8^{(2)}$  in terms of  $f_{D_5}$  will necessarily involve “bad” ratios inside the  $f_{D_5}$ ’s which must cancel in the full sum. This is upsetting!

### Fitting with $E_6$ ’s

There are no good  $E_6$ ’s. There are 216  $E_6$ ’s which have seed clusters involving only good  $\mathcal{X}$ -coordinates. None of these 216  $E_6$ ’s have sufficient  $\{v, z\}$ -squares to constitute a completely good  $\sim f_{E_6} \sim R_7^{(2)}$ . The most  $\{v, z\}$ -squares inside one of these  $E_6$ ’s is 33, and there are 64  $E_6$ ’s which have at least 30 (out of 42) good  $\{v, z\}$  squares.