

In this section we determine the space of weight-4 cluster polylogarithms associated with the A_3 cluster algebra.

A comment on notation: when we refer to simply “ A_3 functions”, we are implicitly referring to $A_2 \subset A_3$ functions, i.e. using the A_2 subalgebras to construct a function. This is the weakest form of subalgebra constructibility since A_2 functions, evaluated across all A_2 subalgebras of a larger algebra, form a basis for all cluster polylogarithms on the larger algebra. This is why we don’t explicitly refer to A_2 . However in the future we will look at, for example, $A_3 \subset D_5$ functions, and in that case we will explicitly refer to the fact that we are using A_3 functions as building blocks (although they are of course already built out of A_2 ’s).

Begin with the six A_2 subalgebras in A_3 – as discussed in a previous section, there are multiple ways to think of these subalgebras. The simplest is that these are the $\binom{6}{5}$ pentagons embeddable in a hexagon. These are also present as the six pentagonal faces in the A_3 associahedron, fig. (ref). In practice it is most practical to list the six distinct A_2 seeds that appear in the 14 cluster generated by mutating on the standard A_3 seed, $x_1 \rightarrow x_2 \rightarrow x_3$:

$$x_1 \rightarrow x_2, \quad x_2 \rightarrow x_3, \quad \frac{x_2}{1+x_{12}} \rightarrow \frac{(1+x_1)x_3}{1+x_{123}}, \quad (1)$$

$$\frac{x_1x_2}{1+x_1} \rightarrow x_3, \quad x_1(1+x_2) \rightarrow \frac{x_2x_3}{1+x_2}, \quad x_1 \rightarrow x_2(1+x_3).$$

We now construct an ansatz for f_{A_3} as a sum of f_{A_2} evaluated on these six subalgebras:

$$f_{A_3} = c_1 f_{A_2}(x_1 \rightarrow x_2) + \dots + c_6 f_{A_2}(x_1 \rightarrow x_2(1+x_3)). \quad (2)$$

Due to the inherited properties of f_{A_2} , the above ansatz already has a cluster-y cobracket, satisfies cluster adjacency, and is smooth and real-valued in the positive domain. The remaining problem to solve is to find values for the c_i such that f_{A_3} is invariant (up to an overall sign) under the automorphisms of A_3 . These were discussed in a previous section (ref), we write them down explicitly here:

$$\sigma_{A_3} : \quad x_1 \rightarrow \frac{x_2}{1+x_1+x_1x_2}, \quad x_2 \rightarrow \frac{x_3(1+x_1)}{1+x_1+x_1x_2+x_1x_2x_3}, \quad x_3 \rightarrow \frac{1+x_1+x_1x_2}{x_1x_2x_3}, \quad (3)$$

$$\tau_{A_3} : \quad x_1 \rightarrow \frac{1}{x_3}, \quad x_2 \rightarrow \frac{1}{x_2}, \quad x_3 \rightarrow \frac{1}{x_1}.$$

Recall that σ is essentially the cyclic automorphism on the hexagon, and τ is the dihedral flip.

The first thing to try is to make f_{A_3} completely invariant under the A_3 automorphisms:

$$\sigma_{A_3}(f_{A_3}) = f_{A_3}, \quad \tau_{A_3}(f_{A_3}) = f_{A_3}. \quad (4)$$

However it turns out there is no collection of c_i that solves such a constraint. If instead we impose

$$\sigma_{A_3}(f_{A_3}) = f_{A_3}, \quad \tau_{A_3}(f_{A_3}) = -f_{A_3}, \quad (5)$$

Then we find the solution

$$c_i = \text{constant}. \quad (6)$$

We label this particular solution by

$$f_{A_3}^{+-} = f_{A_2}(x_1 \rightarrow x_2) + \dots + f_{A_2}(x_1 \rightarrow x_2(1 + x_3)) = \sum_{i=1}^6 \sigma_{A_3}^i(f_{A_2}(x_1 \rightarrow x_2)), \quad (7)$$

where the signs indicate behavior under σ and τ , respectively. Similarly, based on the lack of a solution for eq. (4) we say that $f_{A_3}^{++} = 0$. There are two remaining sign choices to check, $f_{A_3}^{-+}$ and $f_{A_3}^{--}$, and we find $f_{A_3}^{-+} = 0$ and

$$f_{A_3}^{--} = \sum_{i=1}^6 (-1)^i \sigma_{A_3}^i(f_{A_2}(x_1 \rightarrow x_2)). \quad (8)$$

In conclusion, there are choices for f_{A_3} , which we denote by their behavior under the dihedral cycle and flip automorphisms: $f_{A_3}^{+-}$ and $f_{A_3}^{--}$. These functions arise purely from the interplay between the overall symmetries of the A_3 cluster algebra and the structure of the A_2 subalgebras in A_3 , i.e. there has been no physics input so far. We can tabulate our results in the following table:

	$\sigma^+\tau^+$	$\sigma^+\tau^-$	$\sigma^-\tau^+$	$\sigma^-\tau^-$
A_3	0	1	0	1

(9)

This displays the total number of weight-4 polylogarithm functions associated with the A_3 cluster algebra with the given sign changes under the σ and τ automorphisms.